

Radial Velocity Upper Limits on Planets around GJ 1151

BENJAMIN J. S. POPE,^{1,2} MEGAN BEDELL,³ JOSEPH R. CALLINGHAM,⁴
HARISH VEDANTHAM,⁴ TIMOTHY W. SHIMWELL,⁴ AND FRIENDS

¹*Center for Cosmology and Particle Physics, Department of Physics, New York University, 726
Broadway, New York, NY 10003, USA*

²*NASA Sagan Fellow*

³*Center for Computational Astrophysics, Flatiron Institute, 162 Fifth Ave, New York, NY 10010,
USA*

⁴*ASTRON, Netherlands Institute for Radio Astronomy, Oude Hoogeveensedijk 4, Dwingeloo, 7991
PD, The Netherlands*

(Received January 1, 2019; Revised January 7, 2019; Accepted June 17, 2019)

Submitted to ApJ

ABSTRACT

The recent detection of circularly polarized, variable low-frequency radio emission from the M dwarf GJ 1151 has been interpreted as a Jupiter-like electron cyclotron maser instability arising from star-planet interaction. The existence or parameters of the proposed planets have not been determined. In order to search for any planets or binary companions, we have obtained new HARPS-N observations

Using these data we do not detect a clear planetary signal, and we put 99th-percentile upper limits on the mass of any companion to GJ 1151 at 5.1 M_⊕.

We infer interpolated, high-SNR, high-resolution spectra of GJ 1151. 

1. INTRODUCTION

Exoplanet science has flourished over the last three decades to be one of the major topics in optical astronomy. The number of known planets has doubled nearly every two years since 1995 (Mamajek 2016) and this accelerating rate of discovery is projected to continue for at least the next decade if current and upcoming space-based surveys deliver their expected results. However despite extensive searches (Bastian et al. 2000; Lecavelier des Etangs et al. 2013; Lynch et al. 2018) neither exoplanets nor their host stars have been detected at radio frequencies, as the quiescent emission of such systems has been too faint for current telescopes. But as the Square Kilometre Array (SKA; Dewdney et al. 2009) and its precursors, in particular LOFAR (the LOw-Frequency ARray: van Haarlem et al. 2013) come online, the orders-of-

magnitude increase in sensitivity and survey speed they provide makes the detection of nearby stars and planets a realistic prospect (Pope et al. 2019).

Rather than explicitly searching for radio emission from known exoplanet hosts, Callingham et al. (2019) cross-matched sources identified by the LOFAR Two-meter Sky Survey (LoTSS: Shimwell et al. 2019) with nearby stellar sources found by *Gaia*, finding the great majority of matches to be chance associations. But by restricting the survey to variable and circularly-polarized emission, the rate of chance associations with background radio galaxies is dramatically reduced. Based on this restricted cross-match, Vedantham et al. (submitted) have detected the M4.5 dwarf GJ 1151 at low radio frequencies with LOFAR, finding a high degree of circular polarization consistent with a possible star-planet magnetic interaction analogous to a scaled-up Jupiter and Io.

In this Letter we analyze new HARPS-N (High Accuracy Radial velocity Planet Searcher: Cosentino et al. 2012) observations of GJ 1151 in order to search for radial velocity signals of the proposed planets. We do not detect any planets, but place strong upper limits of a few Earth masses on the $M \sin i$ of any possible companions, ruling out any short-period massive objects or close binary companion.

2. RV DATA

We obtained 20 epochs of observations of GJ 1151 with HARPS-N from 2018-12-20 to 2019-02-27, as a Director’s Discretionary Time program. While RVs were extracted from these using the standard HARPS pipeline, its performance on this M4.5 dwarf was very poor, resulting in a spurious RV scatter of several km/s. We therefore reprocessed these data using *wobble* (Bedell et al. 2019), a data-driven package which simultaneously non-parametrically constructs a stellar spectral template and telluric spectral components and uses these, rather than model spectral masks, to extract radial velocities.

We found that the second epoch (2018-12-22) had a significantly higher extracted RV uncertainty than the others, and accordingly excluded this from the global *wobble* model. In order to assess template-dependent systematic errors, we conducted a ‘leave-one-out’ cross-validation, excluding one additional epoch at a time and rerunning *wobble* to search for consistency between the outputs. As seen in Figure 1, the different resulting time series are broadly consistent in their directions of deviation from the mean, with a scatter between them of order \sim the quoted uncertainties. We therefore believe the uncertainties on the *wobble* RVs are realistic but that they are also model-dependent systematics, and therefore likely correlated.

3. KEPLERIAN INFERENCE

We use the *Joker* pipeline (Price-Whelan et al. 2017), which is optimized for small numbers of irregularly-spaced observations, to fit Keplerian signals and infer posterior planet parameters. We allow an additional astrophysical jitter (white noise) term to vary with a lognormal prior on jitter $\ln(s/(\text{m/s}))^2 \sim N(1, 2)$ for GJ 1151.




Figure 1. Leave-one-out cross-validation of *wobble* RVs. One epoch at a time is left out of the global model, and the results of processing the remaining epochs are shown in different colours. There is overall consistency between the different time series, with a diversity of order \sim the quoted uncertainty between the individual realizations.

4. DISCUSSION

5. CONCLUSIONS

The results of this analysis conclusively rule out binary companions or planets on short orbits more massive than Neptune as the origin of the radio signal from GJ 1151. We nevertheless cannot rule out in either case planets less massive than a few Earth masses. Given that Vedantham et al. (submitted) find that even a planet with a mass around that of \sim Mercury could explain these signals, there is a very large region of parameter space for these planets that cannot be excluded and the planet hypothesis remains reasonable. Because our target is so red, to achieve significantly higher precision than attained by HARPS would require moving to the infrared, using one of the IR precision RV instruments currently being commissioned, such as the Habitable-zone Planet Finder (HPF: Mahadevan et al. 2012) or CARMENES (Quirrenbach et al. 2010), by which GJ 1151 is already subject to monitoring (Alonso-Floriano et al. 2015).

ACKNOWLEDGEMENTS

This work was performed in part under contract with the Jet Propulsion Laboratory (JPL) funded by NASA through the Sagan Fellowship Program executed by the NASA Exoplanet Science Institute. TRW acknowledges the support of the Australian




Figure 2. *The Joker* posterior samples for GJ 1151.

Research Council (grant DP150100250) and the Villum Foundation (research grant 10118).

BJSP acknowledges being on the traditional territory of the Lenape Nations and recognizes that Manhattan continues to be the home to many Algonkian peoples. We give blessings and thanks to the Lenape people and Lenape Nations in recognition that we are carrying out this work on their indigenous homelands.

This research made use of NASA’s Astrophysics Data System; the SIMBAD database, operated at CDS, Strasbourg, France; the IPython package (Pérez & Granger 2007); SciPy (Jones et al. 2001); and Astropy, a community-developed core Python package for Astronomy (Astropy Collaboration et al. 2013). Some of the data presented in this paper were obtained from the Mikulski Archive for Space Telescopes (MAST). STScI is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS5-26555. Support for MAST for non-HST data is provided by the NASA Office of Space Science via grant NNX13AC07G and by other grants and contracts.

REFERENCES

- | | |
|---|---|
| <p>Alonso-Floriano, F. J., Morales, J. C., Caballero, J. A., et al. 2015, <i>A&A</i>, 577, A128, doi: 10.1051/0004-6361/201525803</p> | <p>Astropy Collaboration, Robitaille, T. P., Tollerud, E. J., et al. 2013, <i>A&A</i>, 558, A33, doi: 10.1051/0004-6361/201322068</p> |
|---|---|




Figure 3. Cornerplot of posterior samples from *The Joker* for GJ 1151.

- Bastian, T. S., Dulk, G. A., & Leblanc, Y. 2000, *ApJ*, 545, 1058, doi: [10.1086/317864](https://doi.org/10.1086/317864)
- Bedell, M., Hogg, D. W., Foreman-Mackey, D., Montet, B. T., & Luger, R. 2019, arXiv e-prints. <https://arxiv.org/abs/1901.00503>
- Callingham, J. R., Vedantham, H. K., Pope, B. J. S., & and, T. W. S. 2019, *Research Notes of the AAS*, 3, 37, doi: [10.3847/2515-5172/ab07c3](https://doi.org/10.3847/2515-5172/ab07c3)
- Cosentino, R., Lovis, C., Pepe, F., et al. 2012, in *Proc. SPIE*, Vol. 8446, Ground-based and Airborne Instrumentation for Astronomy IV, 84461V
- Dewdney, P. E., Hall, P. J., Schilizzi, R. T., & Lazio, T. J. L. W. 2009, *IEEE Proceedings*, 97, 1482, doi: [10.1109/JPROC.2009.2021005](https://doi.org/10.1109/JPROC.2009.2021005)
- Jones, E., Oliphant, T., Peterson, P., & Others. 2001, *SciPy: Open source scientific tools for Python*. <http://www.scipy.org/>

- Lecavelier des Etangs, A., Sirothia, S. K., Gopal-Krishna, & Zarka, P. 2013, *A&A*, 552, A65, doi: [10.1051/0004-6361/201219789](https://doi.org/10.1051/0004-6361/201219789)
- Lynch, C. R., Murphy, T., Lenc, E., & Kaplan, D. L. 2018, *MNRAS*, doi: [10.1093/mnras/sty1138](https://doi.org/10.1093/mnras/sty1138)
- Mahadevan, S., Ramsey, L., Bender, C., et al. 2012, in *Proc. SPIE*, Vol. 8446, Ground-based and Airborne Instrumentation for Astronomy IV, 84461S
- Mamajek, E. 2016, doi: [10.6084/m9.figshare.4057704.v1](https://doi.org/10.6084/m9.figshare.4057704.v1)
- Pérez, F., & Granger, B. E. 2007, *Computing in Science and Engineering*, 9, 21, doi: [10.1109/MCSE.2007.53](https://doi.org/10.1109/MCSE.2007.53)
- Pope, B. J. S., Withers, P., Callingham, J. R., & Vogt, M. F. 2019, *MNRAS*, 484, 648, doi: [10.1093/mnras/sty3512](https://doi.org/10.1093/mnras/sty3512)
- Price-Whelan, A. M., Hogg, D. W., Foreman-Mackey, D., & Rix, H.-W. 2017, *ApJ*, 837, 20, doi: [10.3847/1538-4357/aa5e50](https://doi.org/10.3847/1538-4357/aa5e50)
- Quirrenbach, A., Amado, P. J., Mandel, H., et al. 2010, in *Proc. SPIE*, Vol. 7735, Ground-based and Airborne Instrumentation for Astronomy III, 773513
- Shimwell, T. W., Tasse, C., Hardcastle, M. J., et al. 2019, *A&A*, 622, A1, doi: [10.1051/0004-6361/201833559](https://doi.org/10.1051/0004-6361/201833559)
- van Haarlem, M. P., Wise, M. W., Gunst, A. W., et al. 2013, *A&A*, 556, A2, doi: [10.1051/0004-6361/201220873](https://doi.org/10.1051/0004-6361/201220873)