

E214 The ATLAS Experiment

Alperen Esen
s39aesen@uni-bonn.de

Benjamin Brandt
s08bbran@uni-bonn.de

Date of experiment: 08-09.05.2025

Handed in: 30.05.2025

University Bonn – Physics Faculty

Contents

1. W-mass	3
1.1. Electron Calibration Verification	3
1.2. QCD scale factor	4
1.2.1. Kinematic Variables	4
1.3. Cut selection	4
1.4. Gauge curves	4
1.5. W-mass	4
A. Appendix	4

1. W-mass

With the finished calibration, the mass of the W-boson can now be measured. This is done using the Jacobi peak of the electron transverse momentum distribution. In order to determine the W-mass, we use a data set of actual ATLAS data containing $W \rightarrow e\nu$ events, as well as several simulated data sets also containing $W \rightarrow e\nu$ events. There is also a $Z^0 \rightarrow e^+e^-$ data set to check the validity of the previous calibration. Finally there are data sets for QCD- and non-QCD background events.

1.1. Electron Calibration Verification




Figure 1: $Z_{mass_check_e\ell_p}t-cut.pdf$

cut selection	$M_{Z^0, meas} / \text{GeV}$	$\left \frac{M_{Z^0, meas} - M_{Z^0, lit}}{M_{Z^0, meas}} \right $
$p_{T, e^\pm} > 40 \text{ GeV}$	91.71 ± 0.02	
$p_{T, e^\pm} < 40 \text{ GeV}$	90.5 ± 0.0	
$35 < p_{T, e^\pm} < 55 \text{ GeV}$	91.43 ± 0.02	
$\eta > 2$	89.89 ± 0.05	
$\eta < 0.5 \ \& \ p_{T, e^\pm} > 40$	91.69 ± 0.02	
$\eta < 0.5 \ \& \ p_{T, e^\pm} < 40$	90.56 ± 0.03	
$\phi < 0$	91.14 ± 0.02	
$\phi > 0$	91.13 ± 0.02	

Table 1: Measured Z^0 mass for different cut selections

1.2. QCD scale factor

In order to achieve an accurate measurement of the W-mass, the background processes at ATLAS have to be taken into consideration. At LHC, proton proton collisions cause the measured events, therefore the background coming from QCD processes is large. This causes a problem, since the final products of the $W \rightarrow e\nu$ events we use to measure the W-mass are leptons which do not interact with the strong interaction. This makes the simulation of a QCD background for the $W \rightarrow e\nu$ events difficult. In order to solve this problem, the QCD background is extracted from the data. In order to scale the data correctly, a QCD scale factor is used, since the integrated luminosity of the background is not measured and therefore unknown. In order to get an understanding of the QCD scale factor and its effect on the data, different kinematic values are plotted for different QCD scale factors.

1.2.1. Kinematic Variables

In order to get a feeling for the QCD scale factor and estimate its optimal value, different kinematic Variables are analyzed. The chosen kinematic variables are el_{pt} (the transverse momentum of the electron), et_{mis} (missing transverse energy/momentum), n_{jet} (number of jets in the event) and pt_w (transverse momentum of the W boson). The distributions of these different variables for a scale factor of 1 (meaning no scaling) can be seen in figure 2

el_{pt}

1.3. Cut selection

1.4. Gauge curves

1.5. W-mass

A. Appendix




Figure 2: Distributions of the kinematic variables ptw , $njet$, el_{pt} and $etmis$ for a QCD scale factor of 1