

Qualitative polymerase chain reactions

Polymerase chain reaction (PCR) can reveal the presence of specific DNA sequences or structural variations such as insertions and deletions when combined with gel-based detection of the resulting amplicon. Even single-nucleotide polymorphisms (SNPs) can be detected in a qualitative manner with allele-specific primer design or restriction fragment analysis. This approach is routinely used to screen bacterial or plasmid clones, or to distinguish known genetic variants in research or diagnostic development contexts.

Two common protocols are described: a Taq polymerase-based method for rapid, low-cost colony screening (*Alternative A*), and a high-fidelity protocol with lower error rates and improved specificity, particularly for GC-rich or long templates. The high-fidelity approach is also compatible with downstream amplicon sequencing (*Alternative B*). Note that relative band intensities cannot be interpreted quantitatively due to variability in template input, enzyme activity, and amplification efficiency.

Buffers and cycling conditions may require optimization depending on the application.

Risk assessment

▷ Wear gloves, safety glasses, lab coat



Reviewed: May 22, 2026

Procedures

» Primer design

(1.) Design primers to flank the region of interest, ideally producing an amplicon between 100–1 500 bp.

- Avoid placing a T at the 3' end if possible; it has the weakest stability during extension.
- Avoid long stretches of G or C at the 3' end.

This is why: G/C runs of 2bp or longer often result in mispriming and primer dimer formation.

- Aim for a melting temperature of 55–65 °C, with both primers within 2 °C of each other.
- Check for the absence of significant primer-dimer formation or strong hairpin structures.

Hint: For colony screening, aim for an amplicon size between 200–600 bp to aid amplification.

Hint: For large insertions, use primer pairs where one primer binds within the altered region and the other outside. This can result in the presence or absence of a band, or a size shift on the gel.

(2.) *Optional:* For SNP detection, design allele-specific primers with the variable base positioned at the 3' end of the forward or reverse primer. ⊕

Hint: To improve allele discrimination, introduce an additional deliberate mismatch 1–3 bases upstream of the 3' end to further destabilize off-target amplification.

(3.) *Optional:* For multiplex PCR, design multiple primer pairs with similar annealing temperatures. Avoid cross-reactivity or overlapping product sizes. ⊕

Critical: Always validate each primer pair in single-plex reactions before combining them into a multiplex reaction. ←

🔗 [NGH+89]

A > Colony PCR using Taq DNA polymerase

- | | |
|--|--|
| ☐ PCR vessel
☐ Taq DNA polymerase | ☐ 10 × Long incubation buffer, pH 8.8 ⚡ R0193
– or: Supplied reaction buffer |
|--|--|

- (1.) Pick a small amount of a bacterial (or yeast) colony using a sterile pipette tip or toothpick and resuspend it in 40 μL aqueous buffer. Alternatively, dilute a saturated culture about 5- to 10-fold. Using the same tip, transfer a small amount of the cell suspension into the PCR reaction mix.

Critical: Avoid transferring excess amounts of biomass. Overloading the reaction with DNA may inhibit polymerase activity.

This is why: The cell suspension can be stored short-term to recover positive clones after PCR screening.

- (2.) Prepare 10 μL PCR master mix for each sample plus one negative control:

	2 rxn	8 rxn	Ingredient	Stock	Final
	2.0 μL	8.0 μL	Long incubation buffer	10 ×	1 ×
	2.0 μL	8.0 μL	Betaine	5 M	500 mM
	1.0 μL	4.0 μL	MgCl_2	50 mM	2.5 mM
	1.0 μL	4.0 μL	Forward primer	10 μM	500 nM
	1.0 μL	4.0 μL	Reverse primer	10 μM	500 nM
	0.8 μL	3.2 μL	dNTP mix	10 mM	400 μM
	0.4 μL	1.6 μL	Taq DNA polymerase	5 U/ μL	0.1 U/ μL
	2.0 μL	8.0 μL	Cell suspension, dilute		
	9.8 μL	39.2 μL	Water, reagent-grade		
	To 20.0 μL		To 80.0 μL		

Critical: Taq DNA polymerase lacks hot-start inhibition and may begin extending misprimed or partially annealed products if the reaction is assembled at room temperature. Prepare master mixes on ice and begin thermocycling promptly after setup.

Hint: Commercial Taq polymerase master mixes or reaction buffers may be substituted. The long incubation buffer provided here is broadly compatible with both self-made and commercial Taq in most applications.

This is why: Betaine reduces secondary structure formation, particularly in GC-rich templates, and can improve amplification robustness.

- (3.) **Critical:** Include a background or parental strain as a negative control in each PCR run.

This is why: This helps detect non-specific amplification from homologous genomic DNA regions or carryover contamination.

- (4.) Spin down the reactions briefly and place directly in the thermocycler. Run the following program:

Cycles	Temperature	Duration	Name	Remarks
1 ×	95 °C	3:00 min	Initial denaturation	Set to 7:00 min to lyse yeast or thick bacterial cell walls
20–25 ×	95 °C	0:30 min	Denaturation	
	50–60 °C	0:30 min	Annealing	Set to 5 °C below the primer melting temperature
	72 °C	1:00 min/kbp	Extension	Adjust for target length
1 ×	72 °C	5:00 min	Final extension	
1 ×	12 °C	∞	Hold	

This is why: Taq DNA polymerase has a half-life of 1 h at 95 °C, supporting up to 30 cycles with minimal degradation in most protocols. Avoid long extension times and excess cycling if possible.

- (5.) Proceed with analysis of PCR products as described below.

🔗 [GS12]

B > **High-fidelity PCR for sequence validation**

☐ PCR vessel	☐ 5 × High-fidelity PCR buffer, pH 8.8 
☐ High-fidelity DNA polymerase	– or: Supplied reaction buffer

(1.) Dilute DNA polymerase to 0.08 U/μL in 50% glycerol with 1 × reaction buffer (1:25 from a 2 U/μL stock).

(2.) Prepare 10 μL PCR master mix for each sample plus one negative control:

	2 rxn	8 rxn	Ingredient	Stock	Final
	4 μL	16 μL	High-fidelity PCR buffer	5 ×	1 ×
	2 μL	8 μL	Betaine 	5 M	500 mM
	1 μL	4 μL	MgCl ₂	50 mM	2.5 mM
	1 μL	4 μL	Forward primer	10 μM	500 nM
	1 μL	4 μL	Reverse primer	10 μM	500 nM
	2 μL	8 μL	dNTP mix	10 mM	1.0 mM
	2 μL	8 μL	DNA polymerase, 1:25 dilution	0.08 U/μL	0.008 U/μL
	4 μL	16 μL	Template	5 ng/μL	1 ng/μL
	3 μL	12 μL	Water, reagent-grade		
	To 20 μL	To 80 μL			

Critical: Avoid exceeding 2 ng of a 5 kbp plasmid per reaction. Excess template can inhibit the polymerase and promote nonspecific amplification. High-fidelity enzymes are effective with as little as 1 pg/μL DNA. 

Hint: Titrate polymerase to find the lowest effective concentration. Lower enzyme levels often yield similar results, reduce cost, and often improve specificity. Dilute in 50% glycerol with 1 × reaction buffer.

Hint: Commercial high-fidelity polymerase master mixes or reaction buffers may be substituted. The included high-fidelity buffer recipe performs reliably across a wide range of polymerases and applications. For GC-rich templates, add 1% DMSO or 500 mM betaine. Thermostability and processivity may be altered in modified buffers.

(3.) **Critical:** Include a reaction without the template as negative control in each PCR run. 

(4.) Spin down the reactions briefly and place in the thermocycler. Run the following program:

Cycles	Temperature	Duration	Name	Remarks
1 ×	97 °C	1:00 min	Initial denaturation	Set to 2:00 min for GC-rich templates
25–30 ×	97 °C	0:30 min	Denaturation	
	65–68 °C	0:30 min	Annealing	Set to the primer melting temperature
	72 °C	0:15 min/kbp	Extension	Adjust for target length
1 ×	72 °C	5:00 min	Final extension	
1 ×	12 °C	∞	Hold	

Critical: Consult the manufacturer's recommended cycling conditions. 

This is why: High-fidelity DNA polymerases such as KOD, VELOCITY, or Phusion remain active for hours at denaturing temperatures, enabling robust amplification of long or GC-rich targets. The error rate is typically 0.8×10^{-6} – 1.2×10^{-6} bp⁻¹.

(5.) Proceed with analysis of PCR products as described below. 

(6.) If the amplicon will be used for restriction-ligation cloning  SOP0033, purify over a silica column  SOP0021 or by gel extraction  SOP0022 to remove primers, dNTPs, and polymerase before proceeding with restriction digest.

 [RQW+06]

Analyses

- **Critical:** Avoid opening PCR tubes after thermocycling in the same area used for PCR setup. 
- **Optional:** Digest PCR product with a restriction enzyme that distinguishes between alleles. 

Note: Primers must span a polymorphic restriction site.

- Analyze 5–10 µL of the PCR mix on an agarose gel  SOP0032.

Note: If the reaction contains betaine, it can be directly loaded into the wells.

- When amplified with a high-fidelity DNA polymerase, purify the reaction for DNA sequencing.

Reporting standards

Parameter	E/D	Description	Standard
Primer sequences	E	◦ Nucleotide sequences * Database reference or reference to synthesis batch documentation	MIQE
Polymerase and buffer	E	◦ Enzyme name and vendor * Lot number, buffer formulation, and enzyme units used	MIQE
Thermal cycling program	E	◦ Cycle number, temperature, and duration * Cyclor model and heated lid settings	MIQE
Template DNA	E	◦ Source (colony, plasmid) * DNA concentration, estimated size, and preparation method	
Thermal cycler	D	◦ Brand and model * Calibration date or error margin	
Gel electrophoresis	E	◦ Agarose percentage, ladder, staining method * Voltage, run time, and gel image reference	

Document the essential (E) and desirable (D) parameters where feasible to meet the reproducibility standards expected by journals and data/research repositories — at the minimal (◦) or extended (*) level.

Materials

Colony PCR using Taq DNA polymerase

PCR vessel, thin-walled

Taq DNA polymerase

- *e.g.* OneTaq® DNA Polymerase, New England Biolabs
- *e.g.* Taq DNA polymerase, self-made

Taq reaction buffer

- Long incubation buffer  R0193, 10 ×, pH 8.8, Store at −20 °C.
- *or* Supplied reaction buffer

High-fidelity PCR for sequence validation

PCR vessel, thin-walled

High-fidelity DNA polymerase

- *e.g.* Q5® High-Fidelity DNA Polymerase, New England Biolabs
- *e.g.* PrimeSTAR® HS DNA Polymerase, Takara Bio
- *e.g.* KOD Hot Start DNA Polymerase, TOYOBO

High-fidelity reaction buffer

- High-fidelity PCR buffer  R0194, 5 ×, pH 8.8, Store at −20 °C.
- *or* Supplied reaction buffer

Recipes

Long incubation buffer, pH 8.8, 10 ×

Amount	Ingredient	Stock	Final
9.0 mL	Tris hydrochloride (Tris-Cl), pH 8.0 ◇ R0055	1 M	200 mM
1.5 mL	Potassium chloride (KCl) ◇ R0038	3 M	100 mM
595 mg	Ammonium sulfate ((NH ₄) ₂ SO ₄) [7783-20-2]	132.14 g/mol	100 mM
To 45 mL	Water, reagent-grade		

Adjust pH with potassium hydroxide. Filter sterilize. Dispense into 1 mL aliquots. Store at −20 °C.

10 × Long incubation buffer

20 mM Tris-Cl, 10 mM KCl, 10 mM (NH₄)₂SO₄, pH 8.8



No GHS hazards at the indicated concentrations

- ☐ Dilute below the discharge limit
- ☐ Flush to drain with excess water

Date: Sign: R0193

High-fidelity PCR buffer, pH 8.8, 5 ×

Amount	Ingredient	Stock	Final
20 mL	Long incubation buffer, pH 8.8 ◇ R0193	10 ×	5 ×
2 mL	Triton™ X-100 ◇ R0057	10%	0.5%
	Skin irritation (H315); Serious eye damage (H318); Aquatic toxicity (long-term) (H411)		
0.2 mL	Bovine serum albumin (BSA) ◇ R0005	10%	0.05%
To 40 mL	Water, reagent-grade		

Filter sterilize and aliquot. Store at −20 °C.

5 × High-fidelity PCR buffer

1 × Long incubation buffer, 0.1% Triton™ X-100, 0.01% BSA, pH 8.8



Aquatic toxicity (long-term)

- ☐ Collect as HAZARDOUS WASTE
- ☐ DO NOT wash into sewer

Date: Sign: R0194

Troubleshooting

Colony PCR using Taq DNA polymerase

In Step 1:

- Smearing or low-intensity band at correct size
 - Use a smaller colony. Excess cell material in colony PCR inhibits polymerase. Prepare a dilute lysate or use heat-treated extract as template to reduce inhibitory substances.

In Step 2:

- Non-specific bands
 - Decrease the magnesium chloride content to improve specificity. Typical final concentrations range from 1.5–3.0 mM.
 - Optimize primer design, target length, and template quality.
- Faint bands, weak amplification
 - Increase the magnesium chloride content to increase yield.

High-fidelity PCR for sequence validation

In Step 5:

- Amplification product in water-only control
 - Check primer sequences for dimer formation.
 - Prepare new primers or aliquot stocks into small volumes.
 - Clean pipettes and use filtered tips.
 - Set up reactions in a clean area physically separate from post-PCR analysis to avoid aerosol contamination of primers or template.

List of references

- M.R. Green and J. Sambrook, *Molecular cloning : a laboratory manual*, 4th ed. (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y, 2012).
- M. Ralser, R. Querfurth, H.-J. Warnatz, H. Lehrach, M.-L. Yaspo, and S. Krobitsch, *Biochem. Biophys. Res. Commun.* **347**(3), 747—751 (2006).
- C. Newton, A. Graham, L. Heptinstall, S. Powell, C. Summers, N. Kalsheker, J. Smith, and A. Markham, *Nucleic Acids Res.* **17**(7), 2503—2516 (1989).

Change log

2011-11-08	Matthias Meurer; Michael Knop lab	Colony PCR from yeast cells with self-made Taq DNA polymerase.
2014-08-08	Matthias Meurer; Michael Knop lab	Generic buffers for high-fidelity DNA polymerases (VELOCITY).
2025-06-08	Benjamin C. Buchmuller	Adaptation as SOP.
2026-05-22	Benjamin C. Buchmuller	Fix Long incubation buffer Tris-Cl amount (20 → 9.0 mL) to deliver the intended 200 mM in the 10× buffer (20 mM in 1× working buffer), matching KCl and ammonium sulfate which were already correct and the standard NEB ThermoPol / Promega long-PCR composition.

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see <https://creativecommons.org/licenses/by-sa/4.0/> for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies. Where a brand name, manufacturer, or specific source is cited, it is given as an example and does not imply that other suitable equivalents are excluded.

Current when printed. Visit <https://benjbuch.github.io/check/> or scan the QR code to check for updates.



2bf6493