Non-viral transfection of animal tissue culture cells

Non-viral transfection introduces DNA or RNA into animal tissue culture cells using chemical delivery methods. The approach avoids viral vectors and enables rapid, scalable manipulation of a wide range of cell types to conduct transient expressions, reporter assays, and genetic perturbation screens.

Lipofectamine® reagents (*Alternative A*) are mixtures of cationic lipids (DOSPA) and neutral helper lipids (DOPE) that form liposomes, which fuse with the plasma membrane to release cargo into the cytoplasm. They are broadly effective and widely used in routine applications. FuGENE® (*Alternative B*) and polyethyleneimine (PEI; *Alternative C*) act via electrostatic condensation of nucleic acids into transfectable particles. FuGENE® is serum-tolerant and gentle on cells, while PEI is inexpensive and scalable but more cytotoxic and often less robust. Calcium chloride co-precipitation (*Alternative D*) remains a simple, economical option, though its success depends critically on pH, temperature, and cell type.

Risk assessment

– Work with human-derived material or transgenic cell lines (BSL-2)

▷ Wear gloves, safety glasses, lab coat

□ Dispose waste after inactivation as REGULATED MEDICAL WASTE



Reviewed: Jun 11, 2026

Procedures

›› *Common provisions*

(1.) Prepare high-quality, endotoxin-free DNA for transfection.

> ***This is why:*** Use endonuclease-negative strains such as DH10B, DH5α, JM109, Stbl2, or TOP10™ to propagate plasmids. If using endA$^+$ strains like HB101 or Stbl3, include an additional wash step during purification.

(2.) One day prior to transfection, seed cells to reach 30–40% confluency on the day of transfection. Use the table below to estimate DNA and medium volumes for each format.

Reagent		Format	Flasks	Dishes			Plates (per well)			
			T-175	15 cm	10 cm	6 cm	6-well	12-well	24-well	96-well
		Surface area	175 cm^2	145 cm^2	56.7 cm^2	21.5 cm^2	9.6 cm^2	3.5 cm^2	1.9 cm^2	1.1 cm^2
Cells to seed		20–30%	8.2×10^6	7.0×10^6	3.0×10^6	1.1×10^6	4.2×10^5	1.8×10^5	8.4×10^4	1.4×10^4
Opti-MEM™		10 µL/cm^2	1.8 mL	1.4 mL	550 µL	200 µL	150 µL	100 µL	50 µL	25 µL
DNA	Lipofect.®	250 ng/cm^2	44.0 µg	36.0 µg	14.0 µg	5.4 µg	2.4 µg	875 ng	475 ng	275 ng
	FuGENE®	200 ng/cm^2	35.0 µg	29.0 µg	11.3 µg	4.3 µg	1.9 µg	700 ng	380 ng	220 ng
	PEI	150 ng/cm^2	26.3 µg	21.8 µg	8.5 µg	3.2 µg	1.4 µg	525 ng	285 ng	165 ng
	$CaCl_2$	450 ng/cm^2	78.8 µg	65.3 µg	25.5 µg	9.7 µg	4.3 µg	1.58 µg	855 ng	495 ng

> ***Critical:*** Adjustments may be necessary based on transfection method, cell type, or well geometry. If your format is not listed, calculate DNA and medium volume based on surface area and initial seeding density.

(3.) *Critical:* Consult the manufacturer's protocols for suspension, primary, or difficult-to-transfect cells.

> ***Hint:*** Lipofectamine® 3000 is formulated for use with stem cells and suspension lines, but may require different reagent volumes or co-factors such as P3000 reagent. Lipofectamine® 3000 uses similar amounts of DNA as Lipofectamine® 2000.

A › *Transfection with Lipofectamine® 2000*

- ☐ Transfer DNA, endotoxin-free plasmid; supercoiled preferred
- ☐ Lipofectamine® 2000 Transfection Reagent
- ☐ Opti-MEM™ I Reduced Serum Medium

(1.) Dilute endotoxin-free DNA in Opti-MEM™. Use the volumes given in the planning table above.

(2.) In a separate tube, dilute 3.6 µL Lipofectamine® 2000 per microgram DNA with the same volume of Opti-MEM™ as the DNA solution. Mix gently.

> ***Critical:*** Lipofectamine® amounts should be optimized for each cell line. Adjust by increasing or decreasing 40% to balance efficiency and toxicity. Do not change the Opti-MEM™ volume.

> ***This is why:*** Serum inhibits complex formation by binding cationic lipids or interfering with particle stability. Therefore, Lipofectamine–DNA complexes are preformed in serum-free medium such as Opti-MEM™. Once complexes are added, serum can usually be tolerated.

(3.) Combine equal volumes of DNA and Lipofectamine® 2000 mixtures. Gently pipette or vortex briefly. Incubate at room temperature for 5–20 min, but not longer. 20 min

(4.) In the meantime, reduce or replace the cell culture medium volume to 50%.

(5.) Dispense the DNA–lipid complexes dropwise to each well. Gently rock to distribute; avoid swirling.

(6.) *Optional:* After 6–24 h, replace the medium.

[YH98]

B › *Transfection with FuGENE®*

- ☐ Transfer DNA, endotoxin-free plasmid
- ☐ FuGENE® 6 Transfection Reagent
- ☐ Opti-MEM™ I Reduced Serum Medium

(1.) Dilute DNA in Opti-MEM™ as above. Add FuGENE® at 3 µL per microgram DNA. Vortex briefly.

> ***Note:*** FuGENE® is serum-compatible and often requires no medium change post-transfection.

(2.) Incubate DNA–FuGENE® complexes at room temperature for 10–15 min. 15 min

(3.) Apply complexes dropwise to cells in standard culture medium. Gently rock to distribute.

C › *Transfection with polyethyleneimine (PEI)*

- ☐ Transfer DNA, endotoxin-free plasmid
- ☐ 1 g/L PEI, pH 7.0 R0131
- ☐ Opti-MEM™ I Reduced Serum Medium

(1.) Dilute DNA in Opti-MEM™ as above.

(2.) In a separate tube, dilute 3.0 µL polyethyleneimine (PEI) per microgram DNA with the same volume of Opti-MEM™ as the DNA solution.

(3.) Combine. Vortex briefly. Incubate DNA–PEI complexes for 10–30 min at room temperature. 10 min

(4.) Dilute 6-fold with Opti-MEM™. Apply dropwise to cells. Rock gently; avoid swirling.

> ***Hint:*** For sensitive cell lines, gently pipette the solution down the side of the well and not on top of the cells, so as not to disrupt the adherent cells.

(5.) *Optional:* After 6–24 h, replace the medium.

[BLZ+95]

D › ***Transfection with calcium chloride***

- ☐ Transfer DNA, endotoxin-free plasmid; supercoiled preferred
- ☐ 2 × HEPES-buffered saline, pH 7.1 ⟋ R0132
- ☐ PBS, pH 7.4 ⟋ R0037
- ☐ 1 M $CaCl_2$ ⟋ R0012

(1.) Seed cells at 20–30% confluency already 4–6 h before transfection.

(2.) Dilute DNA in water and supplement $CaCl_2$ to a final concentration of 250 mM.

> ***Note:*** Supercoiled DNA (plasmids) yields best results.

(3.) *Critical:* Drop by drop, add an equal volume of 2 × HEPES-buffered saline. Vortex after each drop. Do not let the calcium complex precipitate!

(4.) Incubate at room temperature for 15 min. 15 min

(5.) Apply dropwise to cells. Rock gently; avoid swirling.

> ***Critical:*** Calcium-mediated uptake depends on temperature and pH. Keep medium pH stable and avoid cell overgrowth.

(6.) After 6–12 h, remove the medium. Wash cells with PBS and feed fresh medium.

[KCO03]

Troubleshooting

Transfection with Lipofectamine® 2000

In Step 6:

– High cytotoxicity with low transfection efficiency
  - Replace or supplement serum-free medium with full medium after 3–9 h.
  - Reduce Lipofectamine volume by 40%. Excessive cationic lipid damages membranes without improving DNA delivery.
  - Ensure cells are 30–40% confluent at transfection. Over-confluent cells are less permissive to lipoplex uptake.

Materials

Transfection with Lipofectamine® 2000

- ☐ **Transfer DNA**, endotoxin-free plasmid; supercoiled preferred
- ☐ **Opti-MEM™ I Reduced Serum Medium**
  Thermo Fisher Scientific, 31985070
- ☐ **Lipofectamine® 2000 Transfection Reagent**
  Thermo Fisher Scientific, 11668019

Transfection with FuGENE®

- ☐ **Transfer DNA**, endotoxin-free plasmid
- ☐ **Opti-MEM™ I Reduced Serum Medium**
  Thermo Fisher Scientific, 31985070
- ☐ **FuGENE® 6 Transfection Reagent**
  Promega, E2691

Transfection with polyethyleneimine (PEI)

- ☐ **Transfer DNA**, endotoxin-free plasmid
- ☐ **Opti-MEM™ I Reduced Serum Medium**
  Thermo Fisher Scientific, 31985070
- ☐ **Polyethyleneimine (PEI)**, 1 g/L, pH 7.0 ⟋ R0131
  Store at −80 °C. Stable for two months at 4 °C.

Transfection with calcium chloride

- ☐ **Transfer DNA**, endotoxin-free plasmid; supercoiled preferred
- ☐ **Phosphate-buffered saline (PBS)**, pH 7.4 ⟋ R0037
- ☐ **HEPES-buffered saline**, 2 ×, pH 7.1 ⟋ R0132
  Stable for 1 year at room temperature. Alternatively, store at −20 °C.
- ☐ **Calcium chloride ($CaCl_2$)**, 1 M ⟋ R0012

Recipes

Polyethyleneimine (PEI), pH 7.0, 1 g/L

Amount	Ingredient	Stock	Final
100 mg	Polyethyleneimine (PEI), linear, transfection-grade [26913-06-4]	25 000 g/mol	
To 100 mL	Water, reagent-grade		

Bring the stirred solution to neutral pH. Be patient, this may take hours! Recheck pH after 15 min to ensure it has not drifted. Filter-sterilize through a 0.22 µm polyethersulfone (PES) membrane to remove undissolved particles. Dispense into 500 µL aliquots. Store at −80 °C. Stable for two months at 4 °C. ***Note:*** Purchase PEI from Polysciences (Cat. no. 23966).

1 g/L Polyethyleneimine (PEI)
pH 7.0



WARNING



Skin sensitization

☐ Hold for EHS review before disposal

Date: Sign: R0131

HEPES-buffered saline, pH 7.1, 2 ×

Amount	Ingredient		Stock	Final
25 mL	HEPES, pH 7.5	R0024	1 M	50 mM
28 mL	NaCl	R0046	5 M	280 mM
0.106 g	Sodium phosphate dibasic (Na_2HPO_4)	[7558-79-4]	141.96 g/mol	1.5 mM
5.4 mL	Glucose, sterile-filtered	R0021	20%	12 mM
To 500 mL	Water, reagent-grade			

Filter sterilize. Stable for 1 year at room temperature. Alternatively, store at −20 °C. ***Critical:*** The pH is extremely critical for a successful transfection; even 0.1 pH units deviation reduces efficiency!

2 × HEPES-buffered saline

25 mM HEPES, 140 mM NaCl, 750 µM Na_2HPO_4, 6 mM Glucose, pH 7.1

No GHS hazards at the indicated concentrations

- ☐ Dilute below the discharge limit
- ☐ Flush to drain with excess water

Date: Sign: R0132

List of references

O. Boussif, F. Lezoualc'h, M. Zanta, M. Mergny, D. Scherman, B. Demeneix, and J. Behr, *Proc. Natl. Acad. Sci. U.S.A.* **92**(16), 7297—7301 (1995).

R.E. Kingston, C.A. Chen, and H. Okayama, *Curr. Protocol. Cell Biol.* **Chapter 20** Unit 20.3 (2003).

J. Yang and L. Huang, *Gene Ther.* **5**(3), 380—387 (1998).

Change log

2017-05-15	Moritz Pott	Initial calcium chloride protocol; Daniel Summerer lab.
2020-05-20	Yael David	Initial Lipofectamine® protocol; Tom Muir lab.
⋮	⋮	⋮
2023-04-04	Benjamin C. Buchmuller	Updated Lipofectamine® volumes.
2025-04-05	Benjamin C. Buchmuller	Harmonized provisions. Updated safety information for PEI.
2026-06-11	Benjamin C. Buchmuller	HBS: switch sodium phosphate dibasic from heptahydrate (efflorescent, so its effective MW drifts) to anhydrous.

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see https://creativecommons.org/licenses/by-sa/4.0/ for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies. Where a brand name, manufacturer, or specific source is cited, it is given as an example and does not imply that other suitable equivalents are excluded.

Current when printed. Visit https://benjbuch.github.io/check/ or scan the QR code to check for updates.

e425fc7