

Histone octamer, tetramer, or dimer formation

Risk assessment

▷ Wear gloves, safety glasses, lab coat



Reviewed: Mar 27, 2025

This is a bench card. Full protocol available online.

>> Nucleosome core particle reconstitution

- ☐ Dialysis tubing, 3.5 kDa MWCO
- ☐ R0099 Unfolding buffer, 10 mL
- ☐ R0100 2 × Refolding buffer, 800 mL

- (1.) Reconstitute 50 nmol of each lyophilized histone in 200 μ L unfolding buffer to 250 μ M (2.5–3.0 g/L).
- (2.) Bring histones into solution by sonication in an ice-cooled water bath for 5 min. Then, nutate the stock solutions in the cold room for 30 min.
- (3.) Clarify the solution by centrifugation at 17 000 $\times g$ and 4 °C for 1 min to remove any insoluble material.
- (4.) Determine the protein concentration of each histone by UV/Vis spectrophotometry at 280 nm. Take multiple measurements to minimize error.

Quality assurance: Use matched quartz cuvettes or a liquid column instrument (NanoDrop 2000). Blank with fresh unfolding buffer. For liquid column measurements, increase drop size to 2 μ L as surface tension is low due to high guanidine content.

- (5.) Prepare sample to a final concentration of 10 μ M particle concentration (octamer, tetramer, or dimer):

Histone	Stock	Ratio	Octamer	Tetramer	Dimer
H2A	250 μ M	1.08	190 μ L	Not applicable	176 μ L
H2B	250 μ M	1.08	190 μ L	Not applicable	176 μ L
H3	250 μ M	1.00	176 μ L	176 μ L	Not applicable
H4	250 μ M	1.00	176 μ L	176 μ L	Not applicable
Unfolding buffer	1 $\times$		1 468 μ L	1 848 μ L	1 848 μ L

- (6.) Transfer the histone mixture to a 3.5 kDa MWCO dialysis cassette using a 18 G or 21 G needle. Dialyze against at least 100 mL 1 $\times$ refolding buffer for 2 h at 4 °C. Stir the dialysis buffer gently.

Critical: This step removes guanidine and allows monovalent salt to drive refolding of the histone complexes. Use at least a 50 vol excess of buffer relative to sample volume to ensure efficient buffer exchange.

- (7.) Repeat the dialysis step two more times.

>> Sample preparation for size exclusion chromatography (SEC)

- (1.) Clarify the dialyzed sample by centrifugation at 17 000 $\times g$ for 5 min at 4 °C.

Critical: Do not skip this step. Aggregates may clog the ultrafiltration device or interfere with SEC.

- (2.) Equilibrate a disposable ultrafiltration device with 1 $\times$ refolding buffer at 12 000 $\times g$ for 5 min at 4 °C.
- (3.) Transfer the cleared sample into the ultrafiltration device. Concentrate at 12 000 $\times g$ for 5 min intervals to reach 250–500 μ L. Between spins, gently pipette or flick the concentrate to prevent edge-aggregation.

Histone octamer, tetramer, or dimer formation

Critical: Do not exceed the manufacturer's recommended centrifugal force, typically $14\,000 \times g$, as this can rupture the membrane and result in sample loss.

- (4.) Equilibrate a $0.45\ \mu\text{m}$ spin filter with refolding buffer and filter the concentrated sample to remove aggregates or particulate matter.

Quality assurance: The overall recovery after concentration and filtration is typically 70–90%, depending on histone quality, aggregation, and labeling. Highly concentrated or dye-labeled samples may require gentler handling or multiple clarifications to avoid loss. Avoid concentrating samples beyond 100–150 μM unless needed for low-volume injections.

>> Size exclusion chromatography of histone complexes

- (1.) Prepare histone samples for injection by clearing, concentrating, and filtering as described above.
- (2.) Inject the sample onto a preparative Superdex™ 200 column equilibrated with 2.5 column volumes refolding buffer.

Critical: Never flow refolding buffer without equilibrating the FPLC system in water! The system is stored in 20% ethanol and salt will crash out and ruin the column.

- (3.) Elute the particles by flowing 1.4 column volumes refolding buffer.

Quality assurance: For octamers, expect a sharp, symmetric peak eluting at 13.5–14.0 mL depending on exact conditions and sample quality. Aggregates or subcomplexes appear earlier or later, respectively.

- (4.) Collect and analyze fractions around the main peak by SDS-PAGE to confirm stoichiometry and purity.

Quality assurance: Octamer-containing fractions typically show near-equal band intensities for all four histones.

Storage of histone dimers, tetramers, and octamers

- (1.) Combine desired fractions, quantify by UV absorbance at 280 nm using the extinction coefficients below. Concentrate over an ultrafiltration device to 72 μM for octamers or 144 μM for dimers as before.
- (2.) Add an equal volume of 100% glycerol to make up final stocks to 72 μM (36 μM for dimers or tetramers). Nutate at 4 °C for 30 min.

Critical: Before reconstituting nucleosomes from these 50% glycerol stocks, dialyze against refolding buffer overnight at 4 °C.

- (3.) Aliquot and store at –20 °C.

 Recipe (available online)  Resources (available online)  Troubleshooting (available online)  Notes (available online)

Lab Protocols — © B. C. Buchmuller. Licensed CC-BY-SA-4.0. Research use; provided as-is, without warranty. Current when printed. Visit <https://benjbuch.github.io/check/?q=dd33c6d> or scan the QR code to read the full protocol or to check for updates.

