


Resources

Choosing a nuclear extraction and fractionation workflow



In Step 1: Salt extraction alone releases soluble nuclear proteins (S2). Partial MNase digestion improves solubilization of chromatin-bound proteins during salt fractionation. After complete digestion, 20% of total histone H3 and H4 may already appear in the first extract (S2), and up to 70% after the second extract (S3), regardless of salt concentration. Remaining histones in the pellet (P4) require acid extraction.

List of references

- | | |
|--|---|
| W.A. Bickmore and H.G.E. Sutherland, <i>EMBO J.</i> 21 (6), 1248—1254 (2002). | K.A. Khan, M.K. Ng, and P. Cheung, <i>Front. Cell Dev. Biol.</i> 8 331 (2020). |
| C. Herrmann, D.C. Avgousti, and M.D. Weitzman, <i>Bio-protocol</i> 7 (6), e2175 (2017). | S.S. Teves and S. Henikoff, <i>Methods Mol. Biol.</i> 833 421—432 (2012). |

Change log

- | | | |
|------------|------------------------|--|
| 2023-10-18 | Benjamin C. Buchmuller | Adaptation as SOP. |
| 2023-12-10 | Benjamin C. Buchmuller | Revised figures; replaced washing step in isosmotic sucrose with commonly used TEK buffer; included useful hints from PMID:32457909. |

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see <https://creativecommons.org/licenses/by-sa/4.0/> for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies.

Current when printed. Visit <https://benjbuch.github.io/check/> or scan the QR code to check for updates.



ece3127

