Isolation of nuclei from tissue culture cells

The nucleus is the largest and densest organelle of a mammalian cell. It can be easily purified by sedimentation in aqueous and non-aqueous solutions. The purified nuclei can be directly used in downstream assays or to isolate nuclear protein complexes, chromatin, histones, nuclear RNA, or nucleoli.

In *Alternative A*, cells are swollen and ruptured by applying gentle mechanical stress. In *Alternative B*, the cellular membrane is solubilized with a mild detergent such as Triton™ X-100 or IGEPAL® CA-630 which serves as a substitute for Shell's discontinued Nonidet P-40. The pore does not impose a sharp size limit: passive exchange falls away gradually with size rather than stopping at a threshold (Timney et al., 2016), and 30–40 kDa is where retention becomes practical to rely on. Even mild treatment can disrupt nuclear membranes, raising diffusion and aggregating nuclei.

Risk assessment

– Work with human-derived material or transgenic cell lines (BSL-2)

▷ Wear gloves, safety glasses, lab coat

☐ Dispose waste after inactivation as REGULATED MEDICAL WASTE



Reviewed: Oct 26, 2026

Procedures

Preparation of cells

☐ PBS, pH 7.5 (at 1×) R0037 ☐ 100× Inhibitor cocktail R0090

(1.) Start with 1×10^6–1×10^9 cells. When working with frozen cell pellets, thaw on ice. Approximate the packed cell volume; this is 1 vol. Typically, 1×10^9 cells amount to 1–2 mL cell volume.

Critical: Keep buffer volumes to a minimum to maintain high protein concentrations.

(2.) Wash cells twice with 5 vol ice-cold PBS with inhibitors. Try to obtain a suspension free of visible clumps. Collect the cells at 400 × *g* for 5 min at 4 °C. 10 min

A › ***Detergent-free isolation of nuclei***

☐ Dounce homogenizer, loose (type A) pestle ☐ Swelling buffer, pH 7.5 R0091

(1.) Resuspend the cell pellet in 5 vol ice-cold 1 × hypotonic swelling buffer with inhibitors. Ensure the suspension is free of clumps. Incubate for 10 min on ice to swell the cells. 10 min

(2.) Collect the cells by centrifugation at 400 × *g* for 10 min at 4 °C, discard the supernatant. 10 min

(3.) Resuspend in 2 vol (at least 0.4 mL) swelling buffer. Transfer to a pre-chilled Dounce homogenizer.

Note: When choosing the proper size for the homogenizer, be sure the total volume of the cell suspension does not exceed one-third of the volume of the lower chamber of the glass cylinder.

Critical: Do not vortex! Great care must be exercised throughout this procedure. Repeated pipetting and centrifugation will result in compromised nuclei which are very susceptible to leaking and disruption.

(4.) Using a type A (or "loose") pestle, lyse the cells by 10–15 strokes. Apply even pressure and deliberate strokes for uniform lysis. Avoid the formation of bubbles and foam. Repeat until lysis is complete.

Quality assurance: Monitor lysis by removing 5–10 µL lysate. Mix with an equal volume of 0.4% Trypan blue; view under a microscope. Stop homogenization when 95% of cells lost their cytoplasmic membrane and nuclei still remain intact.

(5.) Transfer to a centrifuge tube, pellet broken cells and nuclei at 400 × *g* for 5 min at 4 °C. Keep the supernatant as cytoplasmic fraction if desired.

Quality assurance: The nuclear pellet should be white as opposed to the off-white/beige cellular pellet.

> ***Note:*** 400–600 × *g* is sufficient to pellet nuclei and will not pellet most other organelles.

(6.) Wash once in 1.5 vol hypotonic swelling buffer to remove cell membranes.

(7.) Resuspend the nuclei in the desired buffer for downstream experiments.

[DK19]

B › ***Isolation of nuclei in isosmotic sucrose***

- ☐ Sucrose buffer, pH 7.5 R0092
- ☐ 10% Triton™ X-100 R0057

(1.) Wash the cell pellet twice in 2–5 vol ice-cold isosmotic sucrose buffer with inhibitors.

(2.) Resuspend in 1–2 vol ice-cold sucrose buffer with inhibitors. Ensure the suspension is free of clumps. Admix an equal volume of sucrose buffer containing 0.6% Triton X-100. Incubate for 10 min on ice. 10 min

> ***Critical:*** If no single cell suspension is obtained at this point, cell lysis will not be homogenous. Avoid shearing membranes and organelles by pipetting very gently.

(3.) Pellet at 400 × *g* for 5 min at 4 °C. If desired, keep the supernatant as cytoplasmic fraction.

> ***Quality assurance:*** The nuclear pellet should be white as opposed to the off-white/beige cellular pellet.

(4.) Wash twice with 2–5 vol sucrose buffer to remove cell membranes.

(5.) Resuspend the nuclei in the desired buffer for downstream experiments.

[QZW+23]

Storage of isolated nuclei

(1.) Resuspend the nuclei in isosmotic sucrose buffer; snap-freeze in liquid nitrogen or a dry ice–ethanol bath. Store at −80 °C for up to two years. Do not refreeze.

Analyses

- ○ Mix a sample with an equal volume of Trypan blue and view under the microscope. Nuclei take up the dye and should be round, single, and free of chromatin strands; unstained bright cells are unlysed and mark incomplete lysis.
- ○ Determine the nucleic acid content spectrophotometrically. Dilute a sample 100 × in 2 M NaCl, blank against buffer diluted the same way, and record the absorption at 260 nm and 280 nm; A_{260}/A_{280} should range 1.5–1.7. Calculate the DNA concentration as A_{260} × 50 ng/µL × the dilution factor, and adjust it to 0.8–1 µg/µL with buffer, or pellet the nuclei and resuspend them in less.

> ***This is why:*** The high salt concentration will disintegrate the nuclear membrane, releasing the chromatin, and disrupt protein–nucleic acid complexes for the absorption measurement.

- ○ Prepare samples for SDS-PAGE corresponding to $2{\times}10^5$ cells (or 20 µg protein) for the total, cytoplasmic, and nuclear fraction. Immunostain for tubulin (55 kDa; cytoplasm), and/or histone H3 (15 kDa; chromatin) to demonstrate successful fractionation.

Troubleshooting

Preparation of cells

In Step 2:

– Cells clump together.

  ◦ If unable to disrupt the cell pellet, a short pulse of sonication can be attempted to separate aggregates.

Detergent-free isolation of nuclei

In Step 4:

– Incomplete cell lysis after repeated douncing.

  ◦ Use type B (or "tight") pestle instead.

In Step 5:

– No nuclear pellet.

  ◦ Nuclear pellets isolated from small cell numbers (less than $1{\times}10^6$ cells), are often very hard or impossible to see. Centrifuge for 30 min to increase visibility. Proceeding without a visible pellet often also yields decent results.

Isolation of nuclei in isosmotic sucrose

In Step 2:

– Incomplete cell lysis or leaking nuclei in the presence of Triton™ X-100.

  ◦ Vary detergent concentration, typically within 0.1–0.5% Triton™ X-100. Try alternative detergents such as IGEPAL® CA-630. Avoid Tergitol-type NP-40 (CAS 9016-45-9), a nonylphenol ethoxylate of much higher hydrophilicity that is not a substitute for Nonidet P-40 and behaves differently towards membranes.

Materials

Preparation of cells

- ☐ **Phosphate-buffered saline (PBS)**, pH 7.5 (at 1 ×) ⊘ R0037
- ☐ **Inhibitor cocktail**, 100 × ⊘ R0090

Detergent-free isolation of nuclei

- ☐ **Dounce homogenizer**, loose (type A) pestle
  - *e.g.* **Dura-Grind® Stainless Steel Dounce Tissue Grinder** Wheaton, 357546
- ☐ **Swelling buffer**, pH 7.5 ⊘ R0091
  Store at 4 °C.

Isolation of nuclei in isosmotic sucrose

- ☐ **Sucrose buffer**, pH 7.5 ⊘ R0092
  Store at 4 °C.
- ☐ **Triton™ X-100**, 10% ⊘ R0057

Recipes

Swelling buffer, pH 7.5

Amount	Ingredient	Stock	Final
100 µL	Tris hydrochloride (Tris-Cl), pH 7.5 R0055	1 M	10 mM
33 µL	KCl R0038	3 M	10 mM
15 µL	$MgCl_2$ R0031	1 M	1.5 mM
5 µL	Dithiothreitol (DTT) ☐ R0015 Skin irritation (H315); Serious eye irritation (H319)	1 M	0.5 mM
100 µL	Inhibitor cocktail ☐ R0090 Flammable liquid (H225); Serious eye irritation (H319); Reproductive toxicity (H361)	100 ×	1 ×
To 10 mL	Water, reagent-grade		

Add DTT and inhibitor cocktail immediately before use. Store at 4 °C. ***Hint:*** Tris-Cl replaces HEPES-KOH. Add 10 mM β-glycerophosphate R0164 and 1 mM orthovanadate R0075 fresh from stock for phosphatase inhibition. Add 1 µM trichostatin A R0165 and 5 mM nicotinamide R0166 fresh from stock for HDAC inhibition. The R0090 protease cocktail can be replaced with commercial EDTA-free protease inhibitor cocktail R0072. [DLR83]

Swelling buffer

10 mM Tris-Cl, 10 mM KCl, 1.5 mM $MgCl_2$,
☐ 0.5 mM DTT, ☐ 1 × Inhibitor cocktail, pH 7.5



WARNING



Reproductive toxicity

☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0091

Sucrose buffer, pH 7.5

Amount	Ingredient	Stock	Final
150 µL	Tris hydrochloride (Tris-Cl), pH 7.5 R0055	1 M	15 mM
1 000 µL	Sucrose R0066	2.5 M	250 mM
200 µL	KCl R0038	3 M	60 mM
30 µL	NaCl R0046	5 M	15 mM
50 µL	$MgCl_2$ R0031	1 M	5 mM
10 µL	Dithiothreitol (DTT) ☐ R0015 Skin irritation (H315); Serious eye irritation (H319)	1 M	1 mM
100 µL	Inhibitor cocktail ☐ R0090 Flammable liquid (H225); Serious eye irritation (H319); Reproductive toxicity (H361)	100 ×	1 ×
To 10 mL	Water, reagent-grade		

Add DTT and inhibitor cocktail immediately before use. Store at 4 °C. ***Critical:*** Sucrose-containing solutions must be made up fresh on the day. ***Hint:*** See R0091 for inhibitor suggestions. [BRC+08]

Sucrose buffer

15 mM Tris-Cl, 250 mM Sucrose, 60 mM KCl,
15 mM NaCl, 5 mM $MgCl_2$, ☐ 1 mM DTT,
☐ 1 × Inhibitor cocktail, pH 7.5



WARNING



Reproductive toxicity

☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0092

List of references

M. Brand, S. Rampalli, C.-P. Chaturvedi, and F.J. Dilworth, *Nat. Protoc.* **3**(3), 398-409 (2008).

M.P. Czubryt, B. Ramjiawan, and G.N. Pierce, *Mol. Cell. Biochem.* **172**(1-2), 97-102 (1997).

J. DeCaprio and T.O. Kohl, *Cold Spring Harbor Protoc.* **2019**(7), (2019).

J.D. Dignam, R.M. Lebovitz, and R.G. Roeder, *Nucleic Acids Res.* **11**(5), 1475-1489 (1983).

Y. Qin, Y. Zhou, K. Wang, J. Gu, Z. Xiong, W. Zhang, and Y. Chen, *Biol. Res.* **56**(1), 18 (2023).

B.L. Timney, B. Raveh, R. Mironska, J.M. Trivedi, S.J. Kim, D. Russel, S.R. Wente, A. Sali, and M.P. Rout, *J. Cell Biol.* **215**(1), 57-76 (2016).

Change log

2020-03-22	**Shany Koren-Hauer**	Initial version (Detergent-free isolation of nuclei).
2022-07-09	**Benjamin C. Buchmuller**	Adaptation as SOP. Initial version (Isolation in isosmotic sucrose).
⋮	⋮	
2025-02-22	**Benjamin C. Buchmuller**	Corrected references.
2026-05-22	**Benjamin C. Buchmuller**	Cleaned up inhibitor recipes; cocktail definition moved to collection.
2026-10-26	**Benjamin C. Buchmuller**	Credited Dignam et al. (1983) for the swelling buffer, Brand et al. (2008) for alternative B, and Czubryt et al. (1997) for storage; corrected the nuclei concentration target to 0.8 to 1 μg/μL DNA. The nuclear pore imposes no sharp mass cutoff, after Timney and colleagues; the 30 to 40 kDa figure is kept as the practical one.

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see https://creativecommons.org/licenses/by-sa/4.0/ for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies. Where a brand name, manufacturer, or specific source is cited, it is given as an example and does not imply that other suitable equivalents are excluded.

Current when printed. Visit https://benjbuch.github.io/check/ or scan the QR code to check for updates.

3e26da5