

Histone octamer, tetramer, or dimer formation

Histones are inherently predisposed to assemble into their native complexes — H2A·H2B dimers, H3·H4 tetramers, and ultimately the histone octamer — even in complex mixtures. The formation of these complexes can be steered by the physicochemical environment. For example, at low pH, dimers and tetramers remain separate, while at neutral to slightly basic pH, electrostatic repulsion is sufficiently reduced to allow octamer formation (Sperling and Wachtel, 1981).

Under the conditions described here, monovalent ions such as chloride mimic the electrostatic shielding normally provided by DNA and maintain solubility during refolding from guanidine hydrochloride (Luger et al., 1999). This strategy tolerates crude inputs such as lyophilized or unfractionated histone mixtures, and the resulting particles can be separated by size exclusion chromatography to isolate well-defined species for downstream nucleosome reconstitution.

Risk assessment

▷ Wear gloves, safety glasses, lab coat



Reviewed: Mar 27, 2025

Procedures

» Nucleosome core particle reconstitution

- ☐ Dialysis tubing
- or: Dialysis cassette

- ☐ Unfolding buffer, pH 7.4 ⚡ R0099
- ☐ 2 × Refolding buffer, pH 7.4 ⚡ R0100

- (1.) Reconstitute 50 nmol of each lyophilized histone in 200 μ L unfolding buffer to 250 μ M (2.5–3.0 g/L). 🔗

Note: Extinction coefficients to determine histone concentrations are provided as Supplementary Resources.

- (2.) Bring histones into solution by sonication in an ice-cooled water bath for 5 min. Then, nutate the stock solutions in the cold room for 30 min. ⌚ 30 min

This is why: Although the histones are denatured by guanidine, cooling during sonication minimizes heat-induced aggregation and preserves reducing conditions.

- (3.) Clarify the solution by centrifugation at 17 000 $\times$ g and 4 $^{\circ}$ C for 1 min to remove any insoluble material.
- (4.) Determine the protein concentration of each histone by UV/Vis spectrophotometry at 280 nm. Take multiple measurements to minimize error.

Quality assurance: Use matched quartz cuvettes or a liquid column instrument (NanoDrop 2000). Blank with fresh unfolding buffer. For liquid column measurements, increase drop size to 2 μ L as surface tension is low due to high guanidine content. 💎

This is why: Although histones lack tryptophan, tyrosine and phenylalanine residues contribute enough absorbance for reliable quantification under denaturing conditions. Guanidine has minimal absorbance at 280 nm.

- (5.) Prepare sample to a final concentration of 10 μ M particle concentration (octamer, tetramer, or dimer):

Histone	Stock	Ratio	Octamer	Tetramer	Dimer
H2A	250 μ M	1.08	190 μ L	Not applicable	176 μ L
H2B	250 μ M	1.08	190 μ L	Not applicable	176 μ L
H3	250 μ M	1.00	176 μ L	176 μ L	Not applicable
H4	250 μ M	1.00	176 μ L	176 μ L	Not applicable
Unfolding buffer	1 $\times$		1 468 μ L	1 848 μ L	1 848 μ L

Hint: The table gives examples for 22 nmol octamer and 48 nmol dimer (or tetramer) formation in 2.2 mL denaturing buffer, assuming 250 μ M stock solutions. Most likely, one of the histones will limit the total amount that can be assembled. Scale accordingly.

This is why: A small excess of H2A and H2B improves octamer formation and minimizes tetramer contamination in the octamer peak when the sample is polished by size exclusion chromatography.

- (6.) Transfer the histone mixture to a 3.5 kDa MWCO dialysis cassette using a 18 G or 21 G needle. Dialyze against at least 100 mL 1 × refolding buffer for 2 h at 4 °C. Stir the dialysis buffer gently. ⌚ 6h

Critical: This step removes guanidine and allows monovalent salt to drive refolding of the histone complexes. Use at least a 50 vol excess of buffer relative to sample volume to ensure efficient buffer exchange. ←

- (7.) Repeat the dialysis step two more times.

» Sample preparation for size exclusion chromatography (SEC)

- | | |
|--|---|
| ☐ Ultrafiltration device, PES | ☐ Spin filter, CA, 0.45 µm pore size |
|--|---|

- (1.) Clarify the dialyzed sample by centrifugation at 17 000 × g for 5 min at 4 °C.

Critical: Do not skip this step. Aggregates may clog the ultrafiltration device or interfere with SEC. ←

- (2.) Equilibrate a disposable ultrafiltration device with 1 × refolding buffer at 12 000 × g for 5 min at 4 °C.

- (3.) Transfer the cleared sample into the ultrafiltration device. Concentrate at 12 000 × g for 5 min intervals to reach 250–500 µL. Between spins, gently pipette or flick the concentrate to prevent edge-aggregation.

Critical: Do not exceed the manufacturer's recommended centrifugal force, typically 14 000 × g, as this can rupture the membrane and result in sample loss. ←

- (4.) Equilibrate a 0.45 µm spin filter with refolding buffer and filter the concentrated sample to remove aggregates or particulate matter. ⚠

Hint: Use a low protein-binding membrane such as cellulose acetate (CA). For particles containing fluorophore-labeled histones, check compatibility with the filter material.

Quality assurance: The overall recovery after concentration and filtration is typically 70–90%, depending on histone quality, aggregation, and labeling. Highly concentrated or dye-labeled samples may require gentler handling or multiple clarifications to avoid loss. Avoid concentrating samples beyond 100–150 µM unless needed for low-volume injections. 💎

» Size exclusion chromatography of histone complexes

- | | |
|--------------------------------------|---|
| ☐ FPLC system | ☐ Preparative SEC column |
|--------------------------------------|---|

- (1.) Prepare histone samples for injection by clearing, concentrating, and filtering as described above.

Note: Typical injection volumes range from 100–500 µL. Do not exceed the loop size of the column.

- (2.) Inject the sample onto a preparative Superdex™ 200 column equilibrated with 2.5 vol refolding buffer, where 1 vol is one column volume.

Critical: Never flow refolding buffer without equilibrating the FPLC system in water! The system is stored in 20% ethanol and salt will crash out and ruin the column. ←

- (3.) Elute the particles by flowing 1.4 vol refolding buffer. 🔗

Quality assurance: For octamers, expect a sharp, symmetric peak eluting at 13.5–14.0 mL depending on exact conditions and sample quality. Aggregates or subcomplexes appear earlier or later, respectively. 💎

- (4.) Collect and analyze fractions around the main peak by SDS-PAGE to confirm stoichiometry and purity.

Hint: On a preparative Superdex™ 200 column and at a flow rate of 0.2 mL/min, collect 250 µL fractions. Dilute 3 µL of each fraction in 6 µL water to reduce salts, then add 3 µL 4 × SDS loading dye. Analyze 10 µL per lane on a 16% acrylamide gel.

Quality assurance: Octamer-containing fractions typically show near-equal band intensities for all four histones. 💎

Storage of histone dimers, tetramers, and octamers

☐ Glycerol, molecular biology grade

- (1.) Combine desired fractions, quantify by UV absorbance at 280 nm using the extinction coefficients below. Concentrate over an ultrafiltration device to 72 μ M for octamers or 144 μ M for dimers as before.
- (2.) Add an equal volume of 100% glycerol to make up final stocks to 72 μ M (36 μ M for dimers or tetramers). Nutate at 4 °C for 30 min.

Critical: Before reconstituting nucleosomes from these 50% glycerol stocks, dialyze against refolding buffer overnight at 4 °C.
- (3.) Aliquot and store at –20 °C.

Materials**Nucleosome core particle reconstitution**

Dialysis device, 3.5 kDa MWCO

- Dialysis tubing
- or Dialysis cassette

Unfolding buffer $\diamond$ R0099, pH 7.4

Refolding buffer $\diamond$ R0100, 2 $\times$, pH 7.4, Store at 4 °C.

Sample preparation for size exclusion chromatography (SEC)

Ultrafiltration device, PES, 30 kDa MWCO (for octamers), 10 kDa MWCO (for dimers or tetramers)

- Vivaspin® 500 Centrifugal Concentrator, Cytiva Life Sciences

Spin filter, CA, 0.45 μ m pore size

- Corning® Costar® Spin-X® centrifuge tube filters, Sigma-Aldrich

Size exclusion chromatography of histone complexes

FPLC system

Preparative SEC column

- Superdex™ 200 prep grade, Cytiva Life Sciences, cross-linked agarose/dextran composite, 10–600 kDa globular fractionation range

Storage of histone dimers, tetramers, and octamers

Glycerol, CAS 56-81-5, molecular biology grade

Recipes**Unfolding buffer, pH 7.4**

Amount	Ingredient	Stock	Final
200 μ L	Tris hydrochloride (Tris-Cl), pH 7.4 $\diamond$ R0055	1 M	20 mM
5.73 g	Guanidine $\cdot$ HCl, ultrapure [50-01-1]	95.53 g/mol	6 M
100 μ L	Dithiothreitol (DTT) $\square$ $\diamond$ R0015 Skin irritation (H315); Serious eye irritation (H319)	1 M	10 mM
10 μ L	Ethylenediaminetetraacetate (EDTA), pH 8.0 $\diamond$ R0017	0.5 M	0.5 mM
To 10 mL	Water, reagent-grade		

Prepare fresh. **Quality assurance:** Use ultrapure guanidine hydrochloride.

Unfolding buffer

20 mM Tris-Cl, 6 M Guanidine, $\square$ 10 mM DTT, 0.5 mM EDTA, pH 7.4

**WARNING**

Skin and serious eye irritation; Acute oral toxicity

☐ Hold for EHS review before disposal

Date:

Sign:

R0099

Refolding buffer, pH 7.4, 2 ×

Amount	Ingredient	Stock	Final
20 mL	Tris hydrochloride (Tris-Cl), pH 7.4 ⚡ R0055	1 M	20 mM
234 g	NaCl [7647-14-5]	58.44 g/mol	4 M
2 mL	Dithiothreitol (DTT) □ ⚡ R0015 Skin irritation (H315); Serious eye irritation (H319)	1 M	2 mM
2 mL	Ethylenediaminetetraacetate (EDTA), pH 8.0 ⚡ R0017	0.5 M	1 mM
To 1 L	Water, reagent-grade		

Filter sterilize. Add reducing agent right before use. Store at 4 °C.

2 × Refolding buffer

10 mM Tris-Cl, 2 M NaCl, □ 1 mM DTT,
500 μM EDTA, pH 7.4



No GHS hazards at the indicated concentrations

□ Flush to drain with excess water

Date: Sign: R0100

Troubleshooting**Sample preparation for size exclusion chromatography (SEC)**

In Step 4:


- Sample loss when using cyanine-labeled (Cy3, Cy5, etc.) histones.
 - Instead of spin filtration, clarify the sample by centrifugation at 17 000 × g for 10 min and transfer the supernatant to a fresh tube. Repeat once if needed.

Resources**Nucleosome core particle reconstitution**

Histone	Mol. Abs.		Difference	Species	Reference
	Calculated	Empirical			
H2A	4 470 cm ⁻¹ M ⁻¹	4 050 cm ⁻¹ M ⁻¹	+10.4%	<i>B. taurus</i>	D'Anna and Isenberg (1974)
H2B	7 450 cm ⁻¹ M ⁻¹	6 700 cm ⁻¹ M ⁻¹	+11.2%	<i>B. taurus</i>	D'Anna and Isenberg (1974)
H3	4 470 cm ⁻¹ M ⁻¹	4 040 cm ⁻¹ M ⁻¹	+10.6%	<i>B. taurus</i>	D'Anna and Isenberg (1974)
H4	5 960 cm ⁻¹ M ⁻¹	5 400 cm ⁻¹ M ⁻¹	+10.4%	<i>B. taurus</i>	D'Anna and Isenberg (1974)
Dimer, (H2A-H2B)	11 920 cm ⁻¹ M ⁻¹	10 750 cm ⁻¹ M ⁻¹	+10.9%	–	Summed from D'Anna and Isenberg (1974)
Tetramer, (H3-H4) ₂	20 860 cm ⁻¹ M ⁻¹	18 880 cm ⁻¹ M ⁻¹	+10.5%	–	Summed from D'Anna and Isenberg (1974)
Octamer	44 700 cm ⁻¹ M ⁻¹	39 630 cm ⁻¹ M ⁻¹	+12.8%	–	Summed from D'Anna and Isenberg (1974)
Octamer	44 700 cm ⁻¹ M ⁻¹	45 000 cm ⁻¹ M ⁻¹	+6.7%	<i>H. sapiens</i>	Prevelige and Fasman (1987)

In Step 1: Comparison of calculated and empirical molar extinction coefficients at 280 nm for reduced denatured histones or their assembled complexes. *Bos taurus* calf thymus extracts are consistent with the aromatic content of histones from *Xenopus laevis* and *Homo sapiens*. Calculated values use standard absorbance coefficients for tyrosine and phenylalanine. No known histone contains tryptophan. The calculated molar absorptivity is the common practical value in the chromatin field.

Size exclusion chromatography of histone complexes



In Step 3: Elution profile of histone subcomplexes from a Superdex™ 200 column. The histone octamer (108.5 kDa) elutes first, separate from a small H2A·H2B dimer excess (27.0 kDa). In contrast, the (H3·H4)₂ tetramer (53.0 kDa) elutes close to the octamer peak. Note the small shoulder indicative of some octamer-like assemblies formed by the tetramer. Modified from Dyer et al. (2004).

List of references

- J. D'Anna and I. Isenberg, *Biochemistry* **13**(24), 4992—4997 (1974).
P. Prevelige and G. Fasman, *Biochemistry* **26**(10), 2944—2955 (1987).
P.N. Dyer, R.S. Edayathumangalam, C.L. White, Y. Bao, S. Chakravarthy, U.M. Muthurajan, and K. Luger, *Methods Enzymol.* **375** 23—44 (2004).
R. Sperling and E. Wachtel, *Adv. Protein Chem.* **34** 1—60 (1981).
K. Luger, T. Rechsteiner, and T. Richmond, *Methods Mol. Biol.* **119** 1—16 (1999).

Change log

- | | | |
|------------|------------------------|---|
| 2020-06-12 | Bradley J. Lukasak | Initial version. |
| 2020-07-12 | Hai T. Dao | Updated sample preparation for FPLC. |
| 2022-09-11 | Benjamin C. Buchmuller | Adaptation as SOP; convert to molar concentrations. |
| 2025-03-27 | Benjamin C. Buchmuller | Revised size-exclusion protocol. |

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see <https://creativecommons.org/licenses/by-sa/4.0/> for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies. Where a brand name, manufacturer, or specific source is cited, it is given as an example and does not imply that other suitable equivalents are excluded.

Current when printed. Visit <https://benjbuch.github.io/check/> or scan the QR code to check for updates.



a68be23

