

Immunoblotting

Immunoblotting identifies proteins using antibodies that recognize unique epitopes in a linear amino acid sequence or folded structure. Typically, proteins are first separated by electrophoresis [SOP0007], then immobilized (“blotted”) onto a membrane [SOP0008]. This facilitates antibody recognition and allows for thorough washing to remove nonspecific interactions.

Detection is either direct or indirect as dye- or enzyme-conjugated secondary antibodies bind to the primary antibody and generate a signal proportional to the amount of epitope on the membrane. Antibodies from different host species (mouse, rabbit, goat, chicken) or subclasses (IgG1, IgG2a, etc.) can be combined for multiplexed probing. To detect multiple targets with same-species antibodies, membranes can be cut or stripped and reprobed several times. Fluorescent detection (*Alternative A*) is particularly suited for multiplexed quantitative imaging across a wide linear range. Chemiluminescent detection (*Alternative B*) offers higher sensitivity for low-abundance targets and can be performed with inexpensive homemade reagents, but the signal decays over minutes and is less suited to quantification.

Risk assessment

- ▷ Wear gloves, safety glasses, lab coat
- Dispose milk in accordance with local regulations
- Some jurisdictions prohibit draining dairy products due to their biological oxygen demand



Reviewed: May 22, 2026

Procedures

» Immunoprobings

- | | |
|--|--|
| □ Incubation tray <ul style="list-style-type: none">– or: Resealable plastic bag | □ Bovine serum albumin <ul style="list-style-type: none">– or: Nonfat dry milk powder (“Blotto”) |
| □ 0.1% TBS with Tween™ 20 ⚙ R0169 | □ Antibodies |

- (1.) Prepare blocking buffer by diluting Blotto to 5% or BSA to 1–5% in 20 mL TBST. ⚠

Hint: Use Blotto for abundant targets or with high-affinity antibodies, as it is inexpensive and effective under most conditions. Use BSA otherwise or when low background is essential.

Critical: Avoid milk-based blocking buffers such as Blotto for phospho-proteins or streptavidin-based detection. Milk contains phosphatases and biotin, respectively. Phosphatase activity can be quenched with 50 mM sodium fluoride. ←

- (2.) Block the membrane face-up in 5 mL blocking buffer for 30 min at room temperature with agitation. ⌚ 30 min

- (3.) Dilute the primary antibody in blocking buffer (typically 1:1 000 to 1:10 000).

Hint: Dilutions vary by antibody type. 1:10 to 1:100 for hybridoma supernatant, 1:100–1:1 000 for polyclonal, and up to 1:10 000 or higher for monoclonal antibodies.

Hint: Primary and secondary antibodies can often be reused once or twice within two days to a week.

Critical: For multiplexed experiments, avoid combining mouse and rat primary antibodies as they are difficult to cross-adsorb. Chicken antibodies tend to non-specifically bind to PVDF and nylon-based membranes, leading to high background. ←

- (4.) Discard blocking buffer. Incubate with diluted primary antibody for 30–60 min at room temperature or overnight at 4 °C with agitation. ⌚ 30–60 min ⚠

- (5.) Transfer membrane to a clean tray.

- (6.) Wash four times with 10–20 mL TBST for 3 min each with gentle shaking. ⌚ 15 min

- (7.) Dilute secondary antibody in TBST (typically 1:2 000 to 1:10 000) according to vendor instructions.

Critical: Beware, secondary anti-mouse antibodies may be subclass-specific for IgG1, IgG2a, or IgG2b. Use highly cross-adsorbed secondaries when multiplexing with antibodies from different species. ←

- (8.) Incubate with secondary antibody at room temperature for 30–60 min with agitation.
- (9.) Wash again as before.

⌚ 30–60 min
✂

A > *Fluorescent detection*

- (1.) Specifications for infrared dyes used, for example, with LI-COR Odyssey® Imagers:
 - Use IRDye® 800CW or IRDye® 680RD secondary antibodies, at 1:20 000 dilution.
 - Use IRDye® 800CW for the lowest-abundance target; it gives the lowest background.
- (2.) Rinse the membrane with 1 × TBS to remove residual Tween™ 20.
- (3.) Clean the imager surface with distilled water, then 70% ethanol. Dry with a lint-free wipe.
- (4.) Scan the membrane using a suitable fluorescence imager.

✂

B > *Luminescent detection*

□ ECL reagent ◇ R0121

- (1.) Prepare 10 mL ECL reagent per membrane just before use.
- (2.) Drip-dry the membrane and distribute ECL reagent dropwise to cover the surface evenly. Incubate for 1 min at room temperature.
- (3.) Briefly slip the membrane through deionized water and gently drag over the edge of a container to remove excess liquid.
- (4.) Expose to film or camera. The signal will last for at least 10 min.

⌚ 1 min

✂

Stripping and reusing membranes

□ SDS stripping buffer, pH 6.8 ◇ R0122
– or: Glycine stripping buffer, pH 2.0 ◇ R0211

- (1.) Wash membrane in distilled water for 5 min.
- (2.) Strip antibodies using one of the following:

Stripping buffer	NC	PVDF
0.2 M NaOH	✓	✓
SDS stripping buffer, pH 6.8 ◇ R0122	Not recommended	✓
Glycine stripping buffer, pH 2.0 ◇ R0211	✓	✓

Critical: Do not use high concentration of SDS with nitrocellulose membranes as this can destroy membrane integrity.

←

- (3.) Wash membrane twice in distilled water for 5 min.
- (4.) Reblock and reprobe as needed.

Hint: Although it may not be strictly necessary to reblock the membrane after stripping with a mild stripping buffer, it is highly recommended to prevent high background noise. Use Blotto or casein to block when reprobing membranes. PVDF membranes can be reprobed up to five times with care.

Materials

Immunoprobng

Incubation vessel

- Incubation tray
- or Resealable plastic bag

TBS with Tween™ 20 ⚡ R0169, 0.1%, Stable for 1 month at 4 °C.

Blocking agent

- Bovine serum albumin
- or Nonfat dry milk powder ("Blotto")

Antibodies, primary and secondary

Luminescent detection

ECL reagent ⚡ R0121

Stripping and reusing membranes

Stripping buffer

- SDS stripping buffer ⚡ R0122, pH 6.8
- or Glycine stripping buffer ⚡ R0211, pH 2.0

Recipes

TBS with Tween™ 20 (TBST), 0.1%

Amount	Ingredient		Stock	Final
100 mL	Tris-buffered saline (TBS), pH 7.4	⚡ R0053	10 ×	1 ×
5 mL	Tween™	⚡ R0058	20%	0.1%
To 1 L	Water, reagent-grade			

Add sodium azide to 0.05% to increase shelflife. Stable for 1 month at 4 °C. **Hint:** Compatible with PVDF or nitrocellulose (NC) membranes; do not use Tween™ 20 with nylon membranes.

0.1% TBS with Tween™ 20 (TBST)

1 × TBS, 0.1% Tween™



No GHS hazards at the indicated concentrations

☐ Hold for EHS review before disposal

Date: Sign: R0169

Tris hydrochloride (Tris-Cl), pH 8.5, 1 M

Amount	Ingredient		Stock	Final
12.1 g	Tris base	[77-86-1]	121.14 g/mol	1 M
20 mL	HCl, 37%	[7647-01-0]	11.65 M	
	Skin corrosion (H314)			
To 100 mL	Water, reagent-grade			

Adjust pH to 8.5–8.8. Stable for 2 years at room temperature.

1 M Tris hydrochloride (Tris-Cl)

pH 8.5



WARNING

Respiratory irritation

- ☐ Dilute below the discharge limit
- ☐ Flush to drain with excess water

Date: Sign: R0118

Luminol, 100 ×

Amount	Ingredient	Stock	Final
443 mg	Luminol [521-31-3]	177.16 g/mol	250 mM
To 10 mL	Dimethyl sulfoxide (DMSO), reagent-grade		

Dispense into 50 µL aliquots. Store at –20 °C.

100 × Luminol

No GHS hazards at the indicated concentrations

- ☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0119

p-Coumaric acid, 200 ×

Amount	Ingredient	Stock	Final
148 mg	p-Coumaric acid [501-98-4]	164.16 g/mol	90 mM
To 10 mL	Dimethyl sulfoxide (DMSO), reagent-grade		

Dispense into 25 µL aliquots. Store at –20 °C.

200 × p-Coumaric acid

No GHS hazards at the indicated concentrations

- ☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0120

ECL reagent

Amount	Ingredient	Stock	Final
1 mL	Tris hydrochloride (Tris-Cl), pH 8.5 R0118	1 M	100 mM
50 µL	Luminol, in DMSO R0119	200 ×	1 ×
25 µL	p-Coumaric acid, in DMSO R0120	400 ×	1 ×
3 µL	Hydrogen peroxide [7722-84-1]	30%	5 mM
	Serious eye damage (H318)		
To 10 mL	Water, reagent-grade		

Prepare freshly; use immediately.

ECL reagent

100 mM Tris-Cl, 1 × Luminol, 1 × p-Coumaric acid, 5 mM Hydrogen peroxide

No GHS hazards at the indicated concentrations

- ☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0121

SDS stripping buffer, pH 6.8

Amount	Ingredient	Stock	Final
2 mL	Tris hydrochloride (Tris-Cl), pH 6.8 R0102	0.5 M	100 mM
500 µL	SDS R0047	20%	1%
	Skin irritation (H315); Serious eye damage (H318)		
98 µL	2-Mercaptoethanol [60-24-2]	78.14 g/mol	140 mM
	Acute dermal toxicity (H310); Serious eye damage (H318); Reproductive toxicity (H361); Organ toxicity, repeated exposure (H373)		
To 10 mL	Water, reagent-grade		

Do not adjust the pH.

SDS stripping buffer

100 mM Tris-Cl, 1% SDS, 140 mM 2-Mercaptoethanol, pH 6.8



DANGER

Skin sensitization; Acute dermal toxicity; Organ damage through prolonged or repeated exposure; Reproductive toxicity

- ☐ Collect as HAZARDOUS WASTE
☐ DO NOT wash into sewer

Date: Sign: R0122

Glycine stripping buffer, pH 2.0

Amount	Ingredient		Stock	Final
188 mg	Glycine	[56-40-6]	75.07 g/mol	25 mM
To 100 mL	Water, reagent-grade			

Adjust pH to 2.0 with hydrochloric acid.

Glycine stripping buffer
pH 2.0



DANGER

Serious eye damage; Skin corrosion

- ☐ Neutralise to pH 6–9
- ☐ Flush to drain with excess water

Date: Sign: R0211

Troubleshooting

Immunoprobng

In Step 1:

- No or little signal
 - Lower the percentages of Blotto or BSA as they might block antibody epitopes.

In Step 4:

- Weak or no signal
 - Ensure detergent concentrations are not too high; excess Tween or SDS may wash off loosely bound antibodies.

In Step 8:

- High background signal
 - Reduce antibody concentration.
 - Switch blocking buffer, or use more stringent washes.
 - Avoid milk-based blocking buffers such as Blotto for phospho-proteins or streptavidin-based detection.
- Non-specific bands appear in multiple lanes
 - Switch to highly cross-adsorbed secondary antibodies to eliminate cross-reactivity in multiplexed experiments.

Fluorescent detection

In Step 4:

- Fluorescent bleed-through between channels
 - Use spectrally distinct dyes and check for proper filter/channel selection.
 - IRDye® 800CW and 680RD are optimized for dual-channel detection with minimal overlap.
- Image noise or haze across the blot
 - Ensure platen glass is clean and dry before scanning. Dust can scatter excitation light, causing diffuse background.

Luminescent detection

In Step 4:

- Signal saturation or blown-out bands
 - Shorten exposure time or dilute antibody further.
 - Highly abundant proteins can overwhelm detection systems and obscure nearby bands.
- Blurry or smeared bands
 - Ensure even contact between blot and imaging surface.

Change log

2020-03-22	Michael Haugbro	Initial protocol; Tom Muir lab.
2023-01-15	Benjamin C. Buchmuller	Adaptation as SOP. Added fluorescent and chemiluminescent detection sections and membrane stripping; fluorescent detection informed by LI-COR Biosciences "Good West-erns Gone Bad" application guide (2008, rev. 2022); chemiluminescent ECL reagent recipe from the Michael Knop lab "Self-made ECL".
2026-05-22	Benjamin C. Buchmuller	Fix transcription errors in ECL reagent stock concentrations: Luminol 100× → 200×, p-Coumaric acid 200× → 400× (50 µL and 25 µL in 10 mL total dilute 200× and 400× respec-tively; previous values delivered half the declared 1× final concentration and would have halved chemiluminescent signal output).

From the *Lab Protocols* collection by B. C. Buchmuller and contributors (2026). Licensed under Creative Commons Attribution Share Alike 4.0 International; see <https://creativecommons.org/licenses/by-sa/4.0/> for the terms.

For research use. Provided in good faith, without warranty or liability for any use or results. Users are responsible for compliance with local regulations and institutional policies.

Current when printed. Visit <https://benjbuch.github.io/check/> or scan the QR code to check for updates.



1521f19