

Computational classification of SOD2 protein in coral and symbiotic algae from public genomic and transcriptomic data

Benjie Chen
benjie@alum.mit.edu

ABSTRACT

SOD2, or Manganese/Iron Superoxide Dismutase, plays a crucial role in maintaining cellular health across the tree of life. Existing literature is vague in identifying types of SOD2 proteins expressed in both corals and their symbiotic algae; most agree on the presence of a) mitochondrial targeting MnSOD in corals, and b) likely Mn dependent SOD2 in algae. This paper presents evidence for four classes of SOD2 proteins between coral hosts and their symbiotic algae. Public genomic and transcriptomic data reveal mitochondrial targeting and secreted versions of SOD2 among a wide range of cnidaria species, and chloroplast targeting (with a bipartite leader architecture) and secreted versions of SOD2 among several symbiodinium species. The mature/catalytic region of SOD2 is well conserved within each class, but highly differentiated between classes. This paper presents the mature region HMM profiles for each class, with threshold scores high enough such that they are predictive of leader type.

Introduction

SOD2 proteins protect cells against oxidative damage by converting toxic superoxide radicals into hydrogen peroxide and diatomic oxygen. SOD2 proteins can bind either manganese (MnSOD) or iron (FeSOD) as co-factors; their structure consists of conserved residues His26, His74, Asp159, and His163 (human sequence coordinates) forming the anchors of a trigonal bipyramidal metal pocket and variations in the outer shell, based on bound metal, to ensure superoxide dismutase activity [1]. SOD2 is present throughout tree of life, with bacterial SOD2 often preferring Fe as cofactor, mammalian SOD2 almost always requires Mn as a cofactor to function in mitochondria, and metal preference not always phylogenetically constrained [2].

Existing literature is vague on the types of SOD2 proteins in coral hosts and their symbiotic algae. While there is consensus on the presence of mitochondrial MnSOD in corals, other forms of SOD2 in coral and symbiotic algae are rarely discussed/analyzed separately. [3] reports experimental evidence for secreted superoxide dismutase activity in *Stylophora pistillata* without identifying the source of such activity. [4] experimentally identifies two forms of MnSOD in *Hydra vulgaris*, a non-coral species in the cnidaria phylum. [5] computationally compares various annotated MnSOD and FeSOD sequences from various symbiodinium species, with grouping of sequences, based on putative annotation, inconsistent with findings of this paper. Indeed, most transcriptomic analyses in the field depend on blasting against UniProt or NCBI, or matching against the HMM profile K04564, a pan-metagenomic KEGG ortholog ([6]) for SOD2. Neither approaches provide resolution beyond putative superoxide dismutase activity.

This paper presents results from an unbiased computational search and grouping of SOD2 protein sequences, using NCBI deposited genome sequences from 36 coral and 10 symbiodinium species, as well as RNAseq datasets from 5 papers. The goal of this work is to present specific and unambiguous sequence-level profiles for SOD2 proteins in *Anthozoa* and *Dinophyceae*. Although these protein groups have features that are suggestive of biological and biochemical mechanism, the scope of this paper is limited to computational;

there is no experimental work, beyond referenced RNAseq experiments, to functionally validate the different classes of proteins, determine their metal preference, or likely cellular location.

Specifically, a protein sequence database was constructed from a) protein sequences submitted to and associated with NCBI genome accessions, b) protein sequences exhaustively detected using 25k+ KO HMM profiles ([6]) on 6-frame translations of genomic sequences submitted to and associated with NCBI genome accessions, and c) protein sequences generated from transcriptomes from 5 published RNAseq studies. From this database, sequences matching the SOD2 K04564 profile, having PF00081 and PF02777 Pfam ([7]) FeSOD N and C-terminal domains, and having four key metal binding pocket residues ([1]) were retained.

A leader discovery algorithm was employed to look for, in predicted exons if protein sequence was obtained through a gene model (above (a)) or in upstream 3-frame translations (above (b)) or in the protein sequence (above (c)), possible start codons upstream of the PF00081 match. For each sequence retained from the original protein sequence database, zero (if a start codon cannot be found) or more candidate K04564 sequences may be generated from this leader discovery algorithm. This algorithm is described in more detail in the methods section below.

DeepLoc 2.1 ([8]) classification of candidate K04564 sequences and Muscle multi-sequence alignments ([9]) were used to group protein sequences by leader type, localization, and sequence similarity, for *Anthozoa* and *Dinophyceae* separately.

References

1. Perry, J., Shin, D., Getzoff, E. & Tainer, J. The structural biochemistry of the superoxide dismutases. *Biochimica et Biophysica Acta (BBA) - Proteins and Proteomics* (2010).
2. Sendra, K. *et al.* An ancient metalloenzyme evolves through metal preference modulation. *Nature Ecology and Evolution* (2023).
3. Saragonsti, E., Tchernov, D., Katsir, A. & Shaked, Y. Extracellular production and degradation of superoxide in the coral *Stylophora pistillata* and cultured *symbiodinium*. *PLOS One* (2010).
4. Dash, B., Metz, R., Huebner, H., Porter, W. & Phillips, T. Molecular characterization of two superoxide dismutases from *Hydra vulgaris*. *Gene* (2007).
5. Krueger, T. *et al.* Transcriptomic characterization of the enzymatic antioxidants FeSOD, MnSOD, APX and KatG in the dinoflagellate genus *Symbiodinium*. *BMC Evolutionary Biology* (2015).
6. Kanehisa, M. *et al.* From genomics to chemical genomics: new developments in KEGG. *Nucleic Acids Research* (2006).
7. Sonnhammer, E., Eddy, S., Birney, E., Bateman, A. & Durbin, R. Pfam: multiple sequence alignments and HMM-profiles of protein domains. *Nucleic Acids Research* (1998).
8. Nielsen, H. Practical Applications of Language Models in Protein Sorting Prediction: SignalP 6.0, DeepLoc 2.1, and DeepLocPro 1.0. *Large Language Models (LLMs) in Protein Bioinformatics* (2025).
9. Edgar, R. Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology and phylogeny. *Nature Communications* (2022).