

Results

01 • DISTRIBUTION & NORMALITY

is a variable normally distributed?

Table. Normality tests

Variable	Test	Statistic	N	p	Skew	Kurtosis (excess)
score	Shapiro-Wilk	W 0.96	14	.679	0.30	-1.33
score	K-S (Lilliefors)	D 0.13	14	.732	0.30	-1.33
anxiety	Shapiro-Wilk	W 0.98	13	.990	-0.11	-1.28
anxiety	K-S (Lilliefors)	D 0.09	13	.997	-0.11	-1.28

Kurtosis is excess (normal = 0); both via psych::describe (type 3). Shapiro-Wilk applies for 3-5000 cases.

Figure. Distribution shape — score




Figure. Distribution shape — score




Figure. Distribution shape — anxiety





Figure. Distribution shape — anxiety



How to read this test

A small p (below .05) suggests the data depart from normality. With large samples these tests over-flag trivial deviations; with small samples they have low power, so a $p > .05$ does not prove normality — it only means no departure was detected. Either way, judge normality mainly from the Q-Q plot — points hugging the diagonal indicate normality.

APA template: score: Normality was assessed with the Shapiro-Wilk test, $W = .96$, $p = .679$. anxiety: Normality was assessed with the Shapiro-Wilk test, $W = .98$, $p = .990$.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)