

Results

01 • REPEATED-MEASURES ANOVA

3+ conditions on the same subjects

Table 1. Condition descriptives

Condition	N	M	SD
score_t1	60	59.55	5.45
score_t2	60	62.42	5.89
score_t3	60	64.83	6.04

Table 2. Repeated-measures ANOVA

Source	SS	df	MS	F	p	partial η^2 [95% CI]
Condition	836.30	1.78	471.04	78.51	<.001	0.57 [0.48, 1.00]

Table 3. Sphericity (Mauchly's test)


Effect	W	p	GG ϵ	HF ϵ
condition	0.87	.020	0.89	0.91

when sphericity is violated the F-test uses the Greenhouse–Geisser / Huynh–Feldt correction; post-hoc table follows. Mauchly's test & the GG/HF corrections apply only when the repeated factor has 3+ levels (with 2 levels sphericity is automatically met and this table is omitted).

Table 4. Post-hoc comparisons

Pair	M _{diff}	SE	P _{adj}	95% CI
score_t1 - score_t2	−2.86	0.42	<.001	[−3.90, −1.83]
score_t1 - score_t3	−5.27	0.35	<.001	[−6.14, −4.41]
score_t2 - score_t3	−2.41	0.48	<.001	[−3.60, −1.22]

Figure. Means across conditions



How to read this test

Tests whether the average differs across conditions measured on the same people. Check sphericity first — if violated, read the corrected F/p. A significant result means conditions differ; post-hoc tests show which. Your significance threshold (α) is 0.05.

APA template: A repeated-measures ANOVA (GG-corrected) gave $F(1.78, 104.75) = 78.51$, $p < .001$, partial $\eta^2 = .57$ [.48, 1.00].

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)