


1 ANCOVA

Group	Adj. M	SE	95% CI			
control	31.42	1.11	[29.20, 33.65]			
drug_a	35.02	1.11	[32.80, 37.25]			
drug_b	37.69	1.10	[35.48, 39.90]			

Source	SS	df	MS	F	p	partial η^2 [95% CI]
baseline	1757.40	1	1757.40	72.23	<.001	0.56 [0.42, 1.00]
group	392.60	2	196.30	8.07	<.001	0.22 [0.07, 1.00]

Pair	Mdiff (adj.)	SE	padj	95% CI
control - drug_a	-3.60	1.58	.068	[-7.41, 0.21]
control - drug_b	-6.26	1.56	<.001	[-10.03, -2.50]
drug_a - drug_b	-2.66	1.57	.214	[-6.44, 1.11]



Compares group means after statistically removing the influence of a numeric covariate. First confirm the factor $\times$ covariate interaction is non-significant (homogeneity of slopes) — if it is significant, the adjusted-means interpretation is not valid and a different model is needed. Then read the Factor row's F/p for the adjusted group effect, and the adjusted means for the covariate-controlled

group values. Your significance threshold (α) is 0.05.

Controlling for the covariate, an ANCOVA gave $F(2,56)=8.07$, $p < .001$, partial $\eta^2=.22$ [0.07, 1.00].