

Results

01 • MANCOVA

MANOVA with covariate control

Table 1. Multivariate tests (covariate-adjusted)


Effect	Pillai / Wilks	approx F	df1	df2	p
baseline	0.55	33.87	2	55	<.001
group	0.37	6.30	4	112	<.001

Table 2. Adjusted univariate follow-ups

DV	F	df1	df2	p	partial η^2 [95% CI]
outcome	8.07	2	56	<.001	0.22 [0.07, 1.00]
outcome2	7.47	2	56	.001	0.21 [0.06, 1.00]

assumption checks include homogeneity of covariance matrices (Box's M) and homogeneity of regression slopes for each covariate. (Box's M $\chi^2(6)=11.44$, $p=.076$ · slopes $p(\text{baseline} \times \text{group})=.421$)

Figure. Adjusted means per outcome



How to read this test

Like MANOVA, but group differences on the set of outcomes are assessed after controlling for one or more covariates. Interpret the univariate follow-ups only if the multivariate p is significant, and adjust

them for the number of DVs (e.g. Bonferroni) to control familywise error. Your significance threshold (α) is 0.05.

APA template: A MANCOVA gave a covariate-adjusted group effect, Pillai's $V=.37$, $F(4,112)=6.30$, $p < .001$.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)