

Results

01 • KRUSKAL-WALLIS

nonparametric 3+ group comparison

Table 1. Rank summary

Group	N	Median	IQR	Mean rank
control	20	33.55	9.63	24.90
drug_a	20	36.50	11.97	28.15
drug_b	20	38.70	6.60	38.45


Table 2. Kruskal-Wallis test

H	df	p	ϵ^2 [95% CI]
6.56	2	.038	0.11 [0.03, 1.00]

Table 3. Dunn post-hoc

Pair	Z	p _{adj}
control - drug_a	0.59	.556
control - drug_b	2.45	.042
drug_a - drug_b	1.87	.124

Figure. Distribution by group



How to read this test

The nonparametric counterpart to one-way ANOVA. The H/p tests whether any group tends to have systematically higher or lower values (stochastic dominance) — read it as a "median difference" only when the groups have similar distribution shapes. If significant, Dunn's post-hoc shows which pairs differ, with adjusted p-values. ϵ^2 is the effect size. Your significance threshold (α) is 0.05.

APA template: A Kruskal-Wallis test gave $H(2)=6.56$, $p = .038$, $\epsilon^2=.11$ [.03, 1.00].

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)