

Results

01 • MULTIPLE LINEAR REGRESSION

one numeric outcome, several predictors

Table 1. Regression coefficients

	(1)	β	VIF
(Intercept)	20.36 (6.03) [8.11, 32.62]		
pre_score	0.61 (0.08) [0.46, 0.77]	—	1.25
age	0.02 (0.07) [−0.12, 0.16]	—	1.41
group: b	5.35 (1.64) [2.01, 8.69]	—	1.04
method: online	−2.05 (2.02) [−6.16, 2.06]	—	1.25
method: workshop	−3.25 (2.27) [−7.87, 1.37]	—	1.25
Num.Obs.	40		
R ²	0.75		
R ² Adj.	0.71		
F	20.42		
RMSE	4.68		
AIC	250.96		
BIC	262.78		
Log.Lik.	−118.48		

assumption checks: linearity, normality, homoscedasticity, and multicollinearity (VIF).

Figure. Residual diagnostics





Figure. Coefficient plot



How to read this test

Each predictor's B is its effect holding the others constant; p and CI assess it. R^2 is the joint variance explained; VIF flags predictors that overlap too much (multicollinearity). B is in each predictor's own units, so to compare which predictors matter most use the standardized coefficient (β), not B. Your significance threshold (α) is 0.05.

APA template: The model explained $R^2=.75$ of the variance, $F(5,34)=20.42$, $p < .001$; predictor pre_score gave $B=0.61$, $p < .001$.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)

