

Results

01 • POISSON / NEGATIVE BINOMIAL


predict a count outcome

Table 1. Coefficients

	Log-count (B)	IRR
(Intercept)	-1.46 (0.23)	0.23 [0.15, 0.36]
age	0.01 (0.005)	1.01 [1.00, 1.02]
group: b	0.17 (0.14)	1.19 [0.90, 1.56]
Num.Obs.	40	
Dispersion	1.69	
Residual deviance	67.57	
df	37	
Log.Lik.	-98.18	
AIC	202.37	
BIC	207.44	

dispersion check (Poisson): if variance >> mean (over-dispersion), switch to negative binomial, which instead reports theta (its overdispersion parameter) rather than this ratio.

Figure. Fit / residuals



How to read this test

Models event counts. Each predictor's incidence-rate ratio (IRR) gives the multiplicative change in the expected count per unit (>1 more, <1 fewer); p tests significance. Check dispersion to pick Poisson vs. negative binomial. If cases have unequal exposure (different observation time/area/population), add an offset of $\log(\text{exposure})$ so the model predicts rates, not raw counts — omitting it biases the IRRs. Your significance threshold (α) is 0.05.

APA template: Predictor age was associated with the count, $\text{IRR}=1.01$, 95% CI [1.00, 1.02], $p = .007$.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)