

Results

01 • STATIONARITY TESTS (ADF, KPSS)

is the series stationary?

Table. Stationarity tests

Test	Statistic	Lag	p	Conclusion
ADF	-8.38	4	.010	stationary
KPSS	1.81	3	< .010	non-stationary
PP	-26.93	3	< .010	stationary

ADF and KPSS have opposite null hypotheses — the conclusion column reconciles them. The Statistic column holds two different metrics (ADF: Dickey-Fuller τ ; KPSS: LM statistic); KPSS p-values are interpolated and shown bounded (e.g. “> .10” / “< .01”) rather than exact.

Figure. Series & structure





Figure. Series & structure



How to read this test

For ADF, a small p means stationary; for KPSS, a small p means non-stationary. If a series is non-stationary, difference it before modeling. The conclusion column states the verdict. Failing to reject is not proof of the null for either test (both have low power), so strengthen the verdict by also reading the ACF rather than relying on a single p -value. Method: R tseries package ADF/KPSS/PP (Dickey & Fuller; Kwiatkowski, Phillips, Schmidt & Shin, 1992; Phillips & Perron, 1988). Your significance threshold (α) is 0.05.

APA template: ADF gave $\tau = -8.38$, $p = .010$; KPSS gave $LM = 1.81$, $p < .010$; Phillips–Perron gave $Z = -26.93$, $p < .010$ ($\alpha = 0.05$).

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)