

Results

01 • REGRESSION DISCONTINUITY (RDD)


effect at a cutoff

Table. RD estimate

	(1)
RD treatment effect	9.90 (0.21) [9.48, 10.28]
Bandwidth	8.66
N (left)	18
N (right)	16

inference is robust (bias-corrected); the SE / CI use rdrobust robust standard errors. Bandwidth selector: MSE-optimal (mserd); kernel: triangular. Cutoff = 50.00. McCrary density manipulation test: $t = 0.08$, $p = .938$ (McCrary, 2008) — a small p flags sorting/manipulation at the cutoff.

Figure. Discontinuity



How to read this test

Estimates the jump in the outcome right at the cutoff — the estimate is the treatment effect for cases near the threshold. p/CI assess it; the RD plot shows the discontinuity visually. The jump is causal only if subjects cannot control which side of the cutoff they land on and nothing else changes at the same threshold — the footer reports the McCrary density test (a small p flags sorting/manipulation at the

cutoff); also run covariate-balance / placebo checks. Method: R rdrobust package (Calonico, Cattaneo & Titiunik, 2014, 2015).

APA template: At the cutoff, the treatment effect was 9.90, 95% CI [9.48, 10.28], $p < .001$.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)