

RESULTS

Results

01 · SUMMARY STATISTICS

central tendency & spread of numeric variables

Table. Descriptive statistics

Group	Variable	N	M	SD	95% CI	Min	Max	Median	Skew	Kurtosis (excess)
Female	exam_score	120	65.69	8.31	[64.19, 67.20]	44.84	84.12	66.27	−0.17	−0.55
Male	exam_score	120	65.44	8.03	[63.99, 66.89]	44.91	81.23	65.83	−0.19	−0.56
Female	study_hours_per_week	120	6.80	2.37	[6.37, 7.22]	1.84	11.44	6.67	−0.07	−1.11
Male	study_hours_per_week	120	6.59	2.48	[6.14, 7.04]	2.14	11.62	6.45	0.11	−1.17

one row per chosen variable; a Group column is added when "Group by" is used (stats repeat per group). 95% CI is the t-based confidence interval for the mean; kurtosis is excess (normal = 0).

Figure. Distribution — exam_score





Figure. Distribution — study_hours_per_week



Understanding the numbers

Variable. Which column each row of statistics describes - one row per variable you selected (plus a Group column when grouped). This run reports descriptives for 2 variable(s), split out by group.

N. How many valid (non-missing) values went into that row's statistics. Here, N = 120 across the reported row(s).

Mean (M). The arithmetic average - one of the two center measures reported here. Here, M = 6.59–65.69 across the reported row(s) - compare it to the Median to see whether the shape is skewed.

SD. The standard deviation - how far values typically sit from the mean, showing the spread. Here, SD = 2.37–8.31 across the reported row(s).

95% CI. The t-based confidence interval for the population mean - the range likely to contain the true mean. Here, each row's 95% CI brackets its Mean, showing the margin of uncertainty around that estimate.

Min. The smallest observed value - one end of the range. Here, Min = 1.84–44.91 across the reported row(s).

Max. The largest observed value - the other end of the range. Here, Max = 11.44–84.12 across the reported row(s).

Median. The middle value when the data are sorted - a center measure less affected by extreme scores than the mean. Here, Median = 6.45–66.27 across the reported row(s) - read it alongside the Mean above.

Skew. How far the shape departs from a normal bell curve in asymmetry. Here, Skew = –0.19–0.11 across the reported row(s).

Kurtosis (excess). How far the shape departs from a normal bell curve in tail weight/peakedness (normal = 0). Here, Kurtosis = –1.17––0.55 across the reported row(s).

How to read this test

Descriptives summarize your data: the mean and median show the center, the SD and range show the spread, and skew/kurtosis show how far the shape departs from a normal bell curve. There is no significance test here.

APA template: Table X reports descriptive statistics for the study variables.

Statistical basis: Descriptive statistics table (mean, SD, median, min/max, N, skew, kurtosis) (Revelle W (2024). psych: Procedures for Psychological, Psychometric, and Personality Research. Northwestern University, Evanston.). Full references in the exported CITATIONS.txt.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)