



Tip: the journey bar is clickable - jump back to any finished stage, or between your tests, anytime.



RESULTS

Results


01 · FREQUENCIES & CROSS-TABS

counts for categorical data

Table 1. Frequency distribution (one variable)

Category	n	Valid %	Total %	Cumulative %
Blended	80	33.3	33.3	33.3
Flipped	80	33.3	33.3	66.7
Lecture	80	33.3	33.3	100.0

Figure. Category counts



Understanding the numbers

Category. Each distinct value the variable takes, with its count and share of the total. Here, the table lists 3 categories.

n. How many cases fall in that category. Across the categories here, n sums to 240 valid case(s).

Valid %. That category's share of the valid (non-missing) cases - Valid % uses the valid n as its denominator. Here, Valid % is computed over the 240 valid case(s), excluding any Missing row.

Total %. That category's share of the grand total - Total % uses the grand total, including any Missing row, as its denominator. Here, Total % is computed over all 240 case(s) in the dataset.

Cumulative %. The running total of Valid % as you move down the category list, reaching 100% at the last category. Reading down the 3-category table here, Cumulative % climbs to 100% by the final row.

How to read this test

Frequencies show how many cases fall in each category and what share of the total that is. A cross-tab shows counts for combinations of two categories — to test whether the two are related, run a chi-square test.

APA template: Frequencies (and cross-tabulations) are reported in Table X.

Statistical basis: Frequency counts and cross-tabulations (Firke S (2023). janitor: Simple Tools for Examining and Cleaning Dirty Data. R package.). Full references in the exported CITATIONS.txt.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)