


1 Kruskal-Wallis

Group	N	Median	IQR	Mean rank
Blended	80	5.00	3.00	125.96
Flipped	80	2.00	4.00	72.51
Lecture	80	6.00	3.00	163.03

H	df	p	ϵ^2 [95% CI]
69.55	2	<.001	0.29 [0.22, 1.00]

Pair	Z	padj
Blended - Flipped	-4.90	<.001
Blended - Lecture	3.40	<.001
Flipped - Lecture	8.29	<.001



The nonparametric counterpart to one-way ANOVA. The H/p tests whether any group tends to have systematically higher or lower values (stochastic dominance) — read it as a "median difference" only when the groups have similar distribution shapes. If significant, Dunn's post-hoc shows which pairs differ, with adjusted p-values. ϵ^2 is the effect size. Your significance threshold (α) is 0.05.

A Kruskal-Wallis test gave $H(2)=69.55$, $p < .001$, $\varepsilon^2=.29$ [.22, 1.00].

Statistical basis: Kruskal-Wallis H test (nonparametric counterpart to one-way ANOVA) (Kruskal, W. H., Wallis, W. A. (1952). "Use of ranks in one-criterion variance analysis." Journal of the American Statistical Association, 47(260), 583-621.); Dunn's post-hoc pairwise comparisons (Kassambara A (2023). rstatix: Pipe-Friendly Framework for Basic Statistical Tests. R package.); ε^2 effect size (Ben-Shachar MS, Lüdtke D, Makowski D (2020). "effectsize: Estimation of Effect Size Indices and Standardized Parameters." Journal of Open Source Software, 5(56), 2815.).