

RESULTS

Results

01 · SUMMARY STATISTICS

central tendency & spread of numeric variables

Table. Descriptive statistics

Group	Variable	N	M	SD	95% CI	Min	Max	Median	Skew	Kurtosis (excess)
female	score	7	75.29	6.78	[69.02, 81.55]	66.00	85.00	75.00	−0.03	−1.63
male	score	7	76.00	7.39	[69.16, 82.84]	69.00	88.00	72.00	0.48	−1.67
female	anxiety	6	34.17	8.57	[25.18, 43.16]	22.00	45.00	33.50	−0.06	−1.75
male	anxiety	7	31.00	7.59	[23.98, 38.02]	19.00	40.00	33.00	−0.28	−1.65

one row per chosen variable; a Group column is added when "Group by" is used (stats repeat per group). 95% CI is the t-based confidence interval for the mean; kurtosis is excess (normal = 0).

Figure. Distribution — score





Figure. Distribution — anxiety



Understanding the numbers

Variable. Which column each row of statistics describes - one row per variable you selected (plus a Group column when grouped). This run reports descriptives for 2 variable(s), split out by group.

N. How many valid (non-missing) values went into that row's statistics. Here, N = 6–7 across the reported row(s).

Mean (M). The arithmetic average - one of the two center measures reported here. Here, M = 31.00–76.00 across the reported row(s) - compare it to the Median to see whether the shape is skewed.

SD. The standard deviation - how far values typically sit from the mean, showing the spread. Here, SD = 6.78–8.57 across the reported row(s).

95% CI. The t-based confidence interval for the population mean - the range likely to contain the true mean. Here, each row's 95% CI brackets its Mean, showing the margin of uncertainty around that estimate.

Min. The smallest observed value - one end of the range. Here, Min = 19.00–69.00 across the reported row(s).

Max. The largest observed value - the other end of the range. Here, Max = 40.00–88.00 across the reported row(s).

Median. The middle value when the data are sorted - a center measure less affected by extreme scores than the mean. Here, Median = 33.00–75.00 across the reported row(s) - read it alongside the Mean above.

Skew. How far the shape departs from a normal bell curve in asymmetry. Here, Skew = –0.28–0.48 across the reported row(s).

Kurtosis (excess). How far the shape departs from a normal bell curve in tail weight/peakedness (normal = 0). Here, Kurtosis = –1.75––1.63 across the reported row(s).

How to read this test

Descriptives summarize your data: the mean and median show the center, the SD and range show the spread, and skew/kurtosis show how far the shape departs from a normal bell curve. There is no significance test here.

APA template: Table X reports descriptive statistics for the study variables.

Statistical basis: Descriptive statistics table (mean, SD, median, min/max, N, skew, kurtosis) (Revelle W (2024). psych: Procedures for Psychological, Psychometric, and Personality Research. Northwestern University, Evanston.). Full references in the exported CITATIONS.txt.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)