



Tip: the journey bar is clickable - jump back to any finished stage, or between your tests, anytime.



RESULTS

Results


01 · FREQUENCIES & CROSS-TABS

counts for categorical data

Table 1. Frequency distribution (one variable)

Category	n	Valid %	Total %	Cumulative %
control	6	50.0	50.0	50.0
treatment	6	50.0	50.0	100.0

Figure. Category counts



Understanding the numbers

Category. Each distinct value the variable takes, with its count and share of the total. Here, the table lists 2 categories.

n. How many cases fall in that category. Across the categories here, n sums to 12 valid case(s).

Valid %. That category's share of the valid (non-missing) cases - Valid % uses the valid n as its denominator. Here, Valid % is computed over the 12 valid case(s), excluding any Missing row.

Total %. That category's share of the grand total - Total % uses the grand total, including any Missing row, as its denominator. Here, Total % is computed over all 12 case(s) in the dataset.

Cumulative %. The running total of Valid % as you move down the category list, reaching 100% at the last category. Reading down the 2-category table here, Cumulative % climbs to 100% by the final row.

How to read this test

Frequencies show how many cases fall in each category and what share of the total that is. A cross-tab shows counts for combinations of two categories — to test whether the two are related, run a chi-square test.

APA template: Frequencies (and cross-tabulations) are reported in Table X.

Statistical basis: Frequency counts and cross-tabulations (Firke S (2023). janitor: Simple Tools for Examining and Cleaning Dirty Data. R package.). Full references in the exported CITATIONS.txt.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)