


1 Friedman

Condition	Mean rank
score_t1	1.23
score_t2	2.04
score_t3	2.73

χ^2	df	p	Kendall's W [95% CI]
67.19	2	<.001	0.56 [0.48, 1.00]

Pair	padj
score_t1 - score_t2	<.001
score_t1 - score_t3	<.001
score_t2 - score_t3	<.001



The nonparametric counterpart to repeated-measures ANOVA. The χ^2/p tests whether ranks differ across conditions; Kendall's W is the effect size (0–1 agreement). Post-hoc shows which conditions differ. Your significance threshold (α) is 0.05. N = 60 subjects.

A Friedman test gave $\chi^2(2)=67.19$, $p < .001$, $W=.56$ [.48, 1.00].

Statistical basis: Friedman rank test (nonparametric counterpart to repeated-

measures ANOVA) (Friedman, M. (1937). "The use of ranks to avoid the assumption of normality implicit in the analysis of variance." *Journal of the American Statistical Association*, 32(200), 675-701.); Kendall's W effect size (agreement among rankings) (Kendall, M. G. (1948). *Rank Correlation Methods*. Charles Griffin & Company.); Nemenyi post-hoc pairwise comparisons (Nemenyi, P. B. (1963). *Distribution-free multiple comparisons* [Doctoral dissertation, Princeton University].).