




1 Multiple linear regression

	(1)	β	VIF
(Intercept)	20.36 (6.03) [8.11, 32.62]		
pre_score	0.61 (0.08) [0.46, 0.77]	—	1.25
age	0.02 (0.07) [−0.12, 0.16]	—	1.41
group: b	5.35 (1.64) [2.01, 8.69]	—	1.04
method: online	−2.05 (2.02) [−6.16, 2.06]	—	1.25
method: workshop	−3.25 (2.27) [−7.87, 1.37]	—	1.25
Num.Obs.	40		
R ²	0.75		
R ² Adj.	0.71		
F	20.42		
RMSE	4.68		
AIC	250.96		
BIC	262.78		
Log.Lik.	−118.48		





Each predictor's B is its effect holding the others constant; p and CI assess it. R^2 is the joint variance explained; VIF flags predictors that overlap too much (multicollinearity). B is in each predictor's own units, so to compare which predictors matter most use the standardized coefficient (β), not B. Your significance threshold (α) is 0.05.

The model explained $R^2=.75$ of the variance, $F(5,34)=20.42$, $p < .001$; predictor pre_score gave $B=0.61$, $p < .001$.

Statistical basis: Ordinary least squares regression coefficients (B), SE, CI (R Core Team (2026). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.); Standardized coefficients (β) (Lüdtke D, Ben-Shachar MS, Patil I, Makowski D (2020). "Extracting, Computing and Exploring the Parameters of Statistical Models using R." Journal of Open Source Software, 5(53), 2445.); Variance inflation factor (VIF) cutoff guidance (O'Brien, R. M. (2007). "A caution regarding rules of thumb for variance inflation factors." Quality & Quantity, 41(5), 673-690.).