

RESULTS

Results

01 · REGRESSION DISCONTINUITY (RDD)


effect at a cutoff

Table. RD estimate

	(1)
RD treatment effect	9.90 (0.21) [9.48, 10.28]
Bandwidth	8.66
N (left)	18
N (right)	16
Bandwidth sensitivity — half h (4.33): estimate = 9.86, 95% CI [8.74, 10.64]	
Bandwidth sensitivity — double h (17.32): estimate = 9.94, 95% CI [9.55, 10.24]	

inference is robust (bias-corrected): the SE / CI use rdrobust robust standard errors. Bandwidth selector: MSE-optimal (mserd); kernel: triangular. Cutoff = 50.00. McCrary density manipulation test: t = 0.08, p = .938 (McCrary, 2008) — a small p flags sorting/manipulation at the cutoff.

Figure. Discontinuity



Understanding the numbers

RD treatment effect. The estimated jump in the outcome right at the cutoff - the treatment effect for cases near the threshold. Here, the estimated jump is 9.90.

Bandwidth. The MSE-optimal bandwidth: how far from the cutoff on each side observations are used to estimate the jump. Here, bandwidth = 8.66.

N (left). The effective number of observations within the bandwidth on the side below the cutoff. Here, N (left) = 18.

N (right). The effective number of observations within the bandwidth on the side above the cutoff. Here, N (right) = 16.

Bandwidth sensitivity. Re-estimating the RD effect at half and double the MSE-optimal bandwidth - a stable estimate across these should not swing wildly. Here, halving the bandwidth ($h = 4.33$) gives 9.86 [8.74, 10.64], and doubling it ($h = 17.32$) gives 9.94 [9.55, 10.24].

How to read this test

Estimates the jump in the outcome right at the cutoff — the estimate is the treatment effect for cases near the threshold. p/CI assess it; the RD plot shows the discontinuity visually. The jump is causal only if subjects cannot control which side of the cutoff they land on and nothing else changes at the same threshold — the footer reports the McCrary density test (a small p flags sorting/manipulation at the cutoff); also run covariate-balance / placebo checks. Method: R rdrobust package (Calonico, Cattaneo & Titiunik, 2014, 2015).

APA template: At the cutoff, the treatment effect was 9.90, 95% CI [9.48, 10.28], $p < .001$.

Statistical basis: Robust bias-corrected local-polynomial regression discontinuity inference (Calonico, S., Cattaneo, M. D., Titiunik, R. (2014). "Robust Nonparametric Confidence Intervals for Regression-Discontinuity Designs." *Econometrica*, 82(6), 2295-2326.); Implementation (rdrobust package) (Calonico S, Cattaneo MD, Farrell MH, Titiunik R (2017). "rdrobust: Software for Regression-Discontinuity Designs." *The Stata Journal*, 17(2), 372-404.). Full references in the exported CITATIONS.txt.

After export · optional feedback — an anonymous, optional satisfaction survey — opens a short Google Form in a new tab

[How did it go? — Share feedback →](#)