



Tip: the journey bar is clickable - jump back to any finished stage, or between your tests, anytime.



RESULTS

Results

01 · FRIEDMAN

nonparametric repeated measures

Table 1. Rank summary

Condition	Mean rank
rt_baseline	3.00
rt_low	2.00
rt_high	1.00


Table 2. Friedman test

χ^2	df	p	Kendall's W [95% CI]
80.00	2	<.001	1.00 [1.00, 1.00]

Table 3. Post-hoc (Nemenyi)

Pair	p_{adj}
rt_baseline - rt_low	<.001
rt_baseline - rt_high	<.001
rt_low - rt_high	<.001

Figure. Across conditions



Understanding the numbers

χ^2 . The Friedman test statistic, testing whether ranks differ systematically across conditions. Here, $\chi^2(2) = 80.00$.

Condition. Which repeated-measures condition this row describes. Here, the conditions compared are rt_baseline, rt_low, rt_high.

df. The degrees of freedom for the Friedman test, based on the number of conditions. Here, $df = 2$.

Mean rank. The average rank of each condition's scores across subjects - the quantity the test compares. Here, the mean rank is highest for rt_baseline (3.00) and lowest for rt_high (1.00).

p. The probability of rank differences this large (or larger) across conditions if there were truly no systematic difference. Here, $p < .001$ - below the conventional .05 threshold, ranks differ across conditions.

p adj. The Nemenyi post-hoc adjusted p-value for a specific pair of conditions. Here, the strongest pairwise difference is rt_baseline - rt_high, $p \text{ adj} < .001$.

Kendall's W. The effect size for the Friedman test - the degree of agreement in ranking across conditions, from 0 to 1. Here, $W = 1.00 [1.00, 1.00]$.

N. The number of subjects with complete data across every condition. Here, $N = 40$ subjects.

How to read this test

The nonparametric counterpart to repeated-measures ANOVA. The χ^2/p tests whether ranks differ across conditions; Kendall's W is the effect size (0–1 agreement). Post-hoc shows which conditions differ. Your significance threshold (α) is 0.05. $N = 40$ subjects.

APA template: A Friedman test gave $\chi^2(2)=80.00$, $p < .001$, $W=1.00 [1.00, 1.00]$.

Statistical basis: Friedman rank test (nonparametric counterpart to repeated-measures ANOVA) (Friedman, M. (1937). "The use of ranks to avoid the assumption of normality implicit in the analysis of variance." Journal of the American Statistical Association, 32(200), 675-701.); Kendall's W effect size (agreement among rankings) (Kendall, M. G. (1948). Rank Correlation Methods. Charles Griffin & Company.); Nemenyi post-hoc pairwise comparisons (Nemenyi, P. B. (1963). Distribution-free multiple comparisons [Doctoral dissertation, Princeton University].). Full references in the exported CITATIONS.txt.

After export · optional feedback - an anonymous, optional satisfaction survey - opens a short Google Form in a new tab

[How did it go? - Share feedback →](#)