


1 Friedman

Condition	Mean rank
rt_baseline	3.00
rt_low	2.00
rt_high	1.00

χ^2	df	p	Kendall's W [95% CI]
80.00	2	<.001	1.00 [1.00, 1.00]

Pair	padj
rt_baseline - rt_low	<.001
rt_baseline - rt_high	<.001
rt_low - rt_high	<.001



The nonparametric counterpart to repeated-measures ANOVA. The χ^2/p tests whether ranks differ across conditions; Kendall's W is the effect size (0–1 agreement). Post-hoc shows which conditions differ. Your significance threshold (α) is 0.05. N = 40 subjects.

A Friedman test gave $\chi^2(2)=80.00$, $p < .001$, $W=1.00$ [1.00, 1.00].

Statistical basis: Friedman rank test (nonparametric counterpart to repeated-

measures ANOVA) (Friedman, M. (1937). "The use of ranks to avoid the assumption of normality implicit in the analysis of variance." *Journal of the American Statistical Association*, 32(200), 675-701.); Kendall's W effect size (agreement among rankings) (Kendall, M. G. (1948). *Rank Correlation Methods*. Charles Griffin & Company.); Nemenyi post-hoc pairwise comparisons (Nemenyi, P. B. (1963). *Distribution-free multiple comparisons* [Doctoral dissertation, Princeton University].).