


1 Chi-square independence

Row \ Column	No	Yes	Total
Blended	30 [29.00] (37.5% / 34.5%) r=0.28	50 [51.00] (62.5% / 32.7%) r=-0.28	80
Flipped	17 [29.00] (21.3% / 19.5%) r=-3.42	63 [51.00] (78.8% / 41.2%) r=3.42	80
Lecture	40 [29.00] (50.0% / 46.0%) r=3.13	40 [51.00] (50.0% / 26.1%) r=-3.13	80
Total	87	153	240

χ^2	df	p	Cramér's V [95% CI]
14.39	2	<.001	0.24 [0.13, 1.00]



Tests whether two categorical variables are associated. A p below alpha means they are related; Cramér's V gives the strength of association (0–1); each cell's standardized residual (r) beyond about ± 1.96 flags a category combination that significantly drives the association. Valid only when expected counts are mostly ≥ 5 (the app flags this and suggests Fisher's exact) and each case appears once (raw counts, not percentages). Your significance threshold (α) is 0.05.

A chi-square test of independence gave $\chi^2(2, N=240)=14.39$, $p < .001$, $V=.24$ [.13, 1.00].

Statistical basis: Chi-square test of independence (Pearson, K. (1900). "On the criterion that a given system of deviations from the probable in the case of

a correlated system of variables is such that it can be reasonably supposed to have arisen from random sampling." *Philosophical Magazine*, 50(302), 157-175.); Cramér's V effect size (w-family; Cohen, 1988) (Cohen, J. (1988). *Statistical Power Analysis for the Behavioral Sciences* (2nd ed.). Routledge.).