



# 1 Simple linear regression

|                      | (1)                               | $\beta$ |
|----------------------|-----------------------------------|---------|
| (Intercept)          | 56.29<br>(1.42)<br>[53.50, 59.08] |         |
| study_hours_per_week | 1.39<br>(0.20)<br>[0.99, 1.78]    | 0.41    |
| Num.Obs.             | 240                               |         |
| R <sup>2</sup>       | 0.17                              |         |
| R <sup>2</sup> Adj.  | 0.17                              |         |
| F                    | 48.59                             |         |
| RMSE                 | 7.41                              |         |
| AIC                  | 1648.70                           |         |
| BIC                  | 1659.14                           |         |
| Log.Lik.             | -821.35                           |         |





$R^2$  is the share of outcome variance explained. The predictor's  $B$  is the slope (change in outcome per unit), with  $p$  testing whether it differs from zero and the CI showing precision.  $\beta$  is the standardized slope (change in outcome in SDs per 1 SD change in the predictor), useful for comparing effect sizes on a common scale. Your significance threshold ( $\alpha$ ) is 0.05.

A simple linear regression gave  $B=1.39$ ,  $t(238)=6.97$ ,  $p < .001$ ,  $R^2=.17$ .

Statistical basis: Ordinary least squares regression coefficient ( $B$ ), SE, CI (R Core Team (2026). R: A Language and Environment for Statistical Computing. R Foundation for Statistical Computing, Vienna.); Standardized coefficient ( $\beta$ ) (Lüdtke D, Ben-Shachar MS, Patil I, Makowski D (2020). "Extracting, Computing and Exploring the Parameters of Statistical Models using R." Journal of Open Source Software, 5(53), 2445.).