

Quantifying the global burden of parental loss for children: a cause of death approach

Benjamin-Samuel Schlüter^{1,*}, Antonino Polizzi², Diego Alburez-Gutierrez³, Monica Alexander⁴

¹Department of Statistical Sciences, University of Toronto, Canada

²Leverhulme Centre for Demographic Science, University of Oxford, England

³Kinship Inequalities Research Group, Max Planck Institute for Demographic Research, Germany

⁴Departments of Statistical Sciences and Sociology, University of Toronto, Canada

Abstract

Parental loss is an important life course transition that has the potential to impair physical and mental well-being. The timing and cause of parental death are important mediators of the bereavement process. Previous research on parental loss has focused on a subset of causes of death in few countries. We aim to provide a global assessment of the role of different causes of death in the experience of parental loss. Using estimates from the United Nations World Population Prospects and the Global Burden of Disease study, we apply a formal kinship matrix approach to estimate the prevalence and causes of parental loss worldwide. Although the prevalence of experiencing parental loss has generally declined over time, children in the Global South are still more likely to experience bereavement. In countries such as Afghanistan, Mexico, and Zimbabwe, a large proportion of parental deaths are conflict- or epidemic-related. In the United States, the top cause of father bereavement for children is violent death.

1 Introduction

Shared lifetime between children and their parents is longer today than ever before. Declining fertility has led to shrinking kinship networks in many regions of the world during the 20th century. At the same time, widespread improvements in mortality conditions have increased the time shared between younger and older generations (Bengtson 2001), as, on average, children lose their parents later in life.

While mortality conditions have improved, the presence of premature adult death still means many children lose parents at a young age. Parental bereavement has the potential to impair physical and mental well-being (Amato and Anthony 2014; Li et al. 2014; Luecken 2008; Pham et al. 2018; Saarela and Rostila 2019), while the timing and cause of parental death are important mediators of the bereavement process (Li et al. 2014). The early and sudden loss of a parent has been shown to have a particularly negative impact on the well-being and development of affected children (Chen, Chen, and Liu 2009; Leopold and Lechner 2015; Patterson, Verdery, and Daw 2020). In contrast,

*benjamin.schluter@utoronto.ca.

degenerative diseases, such as dementia and Alzheimer’s disease, are associated with increased family caregiver burden before the parental death, with often detrimental effects on the caregivers’ well-being and the quality of family relationships ([Alzheimer’s Association Report 2022](#)).

Despite the importance of assessing the causes of parental death to fully understand the bereavement process of children, our knowledge of the cause-of-death patterns in parental loss is limited. Previous research on parental loss has focused on a small number of countries to study the effects of Covid-19 ([Susan D. Hillis et al. 2021b](#); [Susan D. Hillis et al. 2021a](#); [Snyder et al. 2022](#); [Verdery et al. 2020](#)), all-cause mortality (REFERENCE OR DROP ALL CAUSES?), and the HIV/AIDS epidemic ([Zagheni 2011](#)).

Kin loss is not just the product of present mortality rates. The probability of losing a parent is affected by the fertility rates of the parental generation (which determine the number of children ‘at risk’ of orphanhood), the past mortality conditions affecting the parental and child generation (i.e., if both parents have already died, a child is no longer at risk of orphanhood), and the present mortality conditions (which determine the risk that a parent will die in a given calendar year). Consequently, as countries differ in their mortality and fertility conditions as well as their epidemiological profiles ([Dattani et al. 2023](#); [United Nations 2022](#)), the short- and long-term burden associated with parental loss can be expected to vary across the globe.

In this project, we aim to provide a global assessment of the role of different causes of death in the experience of parental loss. We use a formal kinship matrix approach ([Caswell 2019](#)) in combination with mortality and fertility data from the United Nations World Population Prospects (UNWPP), and to cause-of-death information from the Global Burden of Disease (GBD) study to estimate causes of parental bereavement. In this extended abstract, we present preliminary results for four countries: Afghanistan, Mexico, United States of America (USA), and Zimbabwe.

2 Data and Method

We combined two sources of data to estimate the worldwide parental loss by cause of death: cause-specific mortality, and life table data with its associated population counts, and fertility rates.

We used the 2022 Revision of World Population Prospects ([United Nations 2022](#)) to obtain period life table by sex; period fertility rates for females aged 15 to 49 years old, and yearly population counts by sex, all by 1-year age class and grouped into individual calendar year over the period 1950-2019. Cause of death information was obtained from modeled data from the Institute for Health Metrics and Evaluation ([2019](#)) available over the period 1990-2019. Cause of death information was collected from this source for ages 0, 1-4, and then in five-year age groups until the open age-group 95 and more. We converted these counts into 1-year age group until age 100+, using a penalized composite link model ([Rizzi, Gampe, and Eilers 2015](#)). Twenty-two causes of death were selected corresponding to the level two of the GBD hierarchy of causes.

In order to estimate parental loss by cause, we used the kinship matrix projection model developed by Caswell ([2019](#)) focusing only on the projected kin that are parent (mother and father).

The model combined several extensions to use time-varying and sex-differentiated vital rates while accounting for death from multiple causes ([Caswell 2022](#); [Caswell, Margolis, and Verdery 2023](#); [Caswell and Song 2021](#)). The projection model for parents can be expressed as follows

$$\begin{pmatrix} \mathbf{d}_L^f \\ \mathbf{d}_L^m \\ \mathbf{d}_D^f \\ \mathbf{d}_D^m \end{pmatrix} (x+1, t+1) = \left(\begin{array}{cc|c} \mathbf{U}_t^f & \mathbf{0} & \mathbf{0} \\ \mathbf{0} & \mathbf{U}_t^m & \\ \hline \mathbf{M}_t^f & \mathbf{0} & \\ \mathbf{0} & \mathbf{M}_t^m & \mathbf{0} \end{array} \right) \begin{pmatrix} \mathbf{d}_L^f \\ \mathbf{d}_L^m \\ \mathbf{d}_D^f \\ \mathbf{d}_D^m \end{pmatrix} (x, t)$$

where $\omega = 101$ is the number of ages considered; $\alpha = 22$ are the twenty-two different causes of death considered; the matrix \mathbf{U}_t of dimension $(\omega \times \omega)$ contains the survival probabilities on its main subdiagonal; the matrix \mathbf{M}_t of dimension $(\alpha\omega \times \omega)$ contains the probabilities of dying from the causes considered on its main diagonals; \mathbf{d}_L refers to the age distribution of the parent living and \mathbf{d}_D reflects the age distribution of the parent dying by cause, in year t when a child is aged x ; upper scripts f and m corresponds to female and male, respectively; subscript t refers to the year. The block matrix on the right-hand side allows to project the parents' age distribution (alive or dead) over time, as their child ages. The model is fit to each country separately.

The model requires as input the age distribution of parents of offspring (see Caswell and Song (2021) for more details). We assumed that in a given year t , both parents were alive at the time of birth and the age distribution of parents at the birth of their child (when $x = 0$) is expressed as $\boldsymbol{\pi}_t = \frac{\mathbf{f}_t \circ \mathbf{n}_t}{\|\mathbf{f}_t \circ \mathbf{n}_t\|}$, where \mathbf{f}_t is a vector of dimension $(\omega \times 1)$ containing age-specific fertility rates and \mathbf{n}_t is a vector of dimension $(\omega \times 1)$ being the age distribution of the overall population. Hence, at the birth of a child in year t ,

$$\begin{pmatrix} \mathbf{d}_L^f \\ \mathbf{d}_L^m \\ \mathbf{d}_D^f \\ \mathbf{d}_D^m \end{pmatrix} (0, t) = \begin{pmatrix} \boldsymbol{\pi}_t^f \\ \boldsymbol{\pi}_t^m \\ \mathbf{0} \\ \mathbf{0} \end{pmatrix}$$

We assumed that female and male age-specific fertility rates were equal as the UN does not provide male fertility (Alburez-Gutierrez, Williams, and Caswell 2023). Cause-specific mortality was not available for the period 1950–1990. During this period, we simplified the block-matrix to only contain survival probabilities, meaning that we only projected the age distribution of parents alive before 1990 ($\mathbf{M}_t = \mathbf{0}$ for $t < 1990$). We additionally recorded the age distribution of parents dying by cause from 1990 until 2019. As it is commonly done in these models (Snyder et al. 2022; Verdery et al. 2020), before 1950, we assumed that the earliest available rates have been operating for a long time (stable population assumption).

3 Preliminary Results

Figure 1 shows the prevalence of underage (below age 18) children who ever lost a father (left panel) or mother (right panel) over the period 1970–2019 for the four selected countries, Afghanistan, Mexico, the United States, and Zimbabwe. There are clear differences in country-specific trends over time. In the U.S., the prevalence of ever having lost a father remained relatively constant over the five decades of observation, at about 5%, while this prevalence declined sharply between the 1970s and the mid-2000s in Mexico. In Afghanistan, the prevalence of paternal death also showed a declining trend, with the exception of the periods 1979–1992 and 2017–2019, which were marked by large-scale conflict. Zimbabwe represents an exception to the general trend of low or declining paternal bereavement, with the likelihood of ever having lost a father increasing sharply during the 1980s and 1990s, which coincided with the HIV/AIDS crisis. During the 2010s, the likelihood of

paternal bereavement in Zimbabwe started to decline again.

While the prevalence of maternal loss was generally lower than the one of paternal loss, trends in maternal and paternal bereavement followed similar trends in each country. Prevalences of maternal bereavement were usually affected little by periods of conflict (e.g., Afghanistan in the 1980s, 1977–1979 in Zimbabwe).




Figure 1: Prevalence of children losing a parent, by parent sex and country over period 1970-2019

Figure 2 breaks down the prevalence of paternal and maternal bereavement among underage children by causes of death for the period 1990–2019. For each country-sex combination, the prevalence of paternal and maternal loss is shown for the top three causes of death over the entire thirty-year period.

In line with trends displayed in Figure 1, paternal and maternal bereavement from “self-harm and interpersonal violence”, “cardiovascular diseases”, and “neoplasms” in the United States have displayed relatively stagnant trends in the 21st century. Similarly, following the all-cause trends in Figure 1, maternal bereavement from “cardiovascular diseases”, “maternal and neonatal disorders”, and “neoplasms” in Afghanistan and maternal bereavement from “neoplasms”, “cardiovascular diseases”, and “diabetes and kidney diseases” in Mexico showed declining trends, albeit to varying degrees. In Zimbabwe, parental bereavement due to “HIV/AIDS and sexually transmitted infections” clearly stands out as the driver of the all-cause trends shown in Figure 1. At the peak of the HIV/AIDS crisis in Zimbabwe, 16% of children had previously lost their father to “HIV/AIDS and sexually

transmitted infections”, while 20% of children had previously lost their mother. Finally, offsetting trends among different causes of death were seen for fathers in Afghanistan and Mexico, where, at different time points, high and rising bereavement from “self-harm and interpersonal violence” counteracted improvements in the remaining two causes of death (Afghanistan: “cardiovascular diseases” and “transport injuries”; Mexico: “digestive diseases” and “transport injuries”).




Figure 2: Prevalence of children losing a parent, by parent sex and country top three causes of parental death over period 1990-2019

Finally, Figure 3 shows the prevalence (y-axis) of having lost at least one parent (father or mother) by age (x-axis) and cause of death (area color) in Afghanistan (left) and Mexico (right) in the year 2019. The top panels show the probability of parental loss from “cardiovascular diseases” and “neoplasms”, while the bottom panels focus on the remaining top ten causes of death in each country in 2019 (excluding “cardiovascular diseases” and “neoplasms”).

In Afghanistan (20%), the prevalence of having lost a parent to any cause was higher in childhood (i.e., below age 18) than in Mexico (9%). Looking at the bottom panel of Figure 3, it becomes clear that “self-harm and interpersonal violence”, “transport injuries”, and “unintentional injuries” were the main contributors to this pattern.

The adult peak of parental loss was reached at around age 50 in Afghanistan and at around age 60 in Mexico. Parental loss from “self-harm and interpersonal violence” was more important for parental loss at all ages in Afghanistan. In contrast, “diabetes and kidney diseases”, “digestive diseases”, and

“neurological disorders” played a more important role in Mexico.




Figure 3: Age- and cause-specific bereavement prevalences of losing at least one parent, by country top twelve causes in 2019

4 Summary, Discussion, and Next Steps

In our project, we aim to provide a global assessment of the role of different causes of death in the experience of parental loss. Using a formal matrix approach, our study highlights the stark global differences in the experience of parental loss. Although the prevalence of experiencing parental loss has generally declined over time, children in the Global South are still more likely to lose a parent. In the second half of the 20th century, many countries in the Global South, such as Afghanistan, Mexico, and Zimbabwe, have experienced short- or longer-term mortality crises resulting from conflicts or epidemics. These mortality crises have temporarily exacerbated experiences of parental loss (as in Afghanistan and Zimbabwe) or slowed down improvements in parental longevity (as in Mexico). Our findings show that in countries such as Afghanistan, Mexico, and Zimbabwe, a large proportion of parental deaths are conflict- or epidemic-related and can occur suddenly in a context already marked by social upheaval. Moreover, in Afghanistan, the high prevalence of mortality from maternal and neonatal disorders suggests that many children may grow up without a mother. While parental loss is less common in the United States, a large proportion of bereaved children in the U.S. have lost their father to a violent death. Future work will focus on presenting and interpreting results for all countries and deriving region-base estimates.

The preliminary results presented in this extended abstract represent only a small subset of countries and are subject to two main limitations. First, in many data-scarce contexts, the age-specific mortality and fertility rates and the cause-of-death information provided by UNWPP and GBD represent estimated quantities themselves. Consequently, biases in the mortality and fertility information are carried over to the estimates of parental loss derived from our matrix kinship models. GBD, and in some instances UNWPP, provide uncertainty bounds around their mortality and fertility estimates. For the PAA 2024 annual meeting, we aim to explicitly incorporate this uncertainty in our estimates of parental loss. Moreover, for selected countries, such as the United States and Mexico, we plan to conduct robustness checks with cause-of-death information derived directly from vital registration systems.

Second, in the absence of male fertility data, we approximate male fertility rates using an “androgynous model” to derive kin estimates, which assumes that female and male fertility patterns in a given country are identical. Existing information on female and male fertility suggests that the reproductive period of males is generally longer and the mean age of childbirth higher ([Dudel and Klüsener 2021](#); [Schoumaker 2019](#)). Estimates of kin relationships derived from population registers vs. matrix models for Sweden suggest that the “androgynous model” approximates register-based estimates of different kin relationships reasonably well ([Alburez-Gutierrez, Williams, and Caswell 2023](#)). For the PAA 2024 annual meeting, we aim to test the sensitivity of our results to different assumptions about male fertility patterns.

Despite these limitations, our study provides important insights into the cause-of-death patterns in parental loss in different world regions. Against general increases in shared lifetime between children and their parents, our results suggest important variations in the short- and long-term burdens associated with parental loss across the globe. Thus, our findings will prove useful in the design of adequate support systems that help underage and adult children in their grieving process and protect children from potential social and economic repercussions arising from the death of their parents.

References

- Alburez-Gutierrez, D., Williams, I., and Caswell, H. (2023). Projections of human kinship for all countries. doi:[10.31235/osf.io/hn3zm](https://doi.org/10.31235/osf.io/hn3zm).
- Alzheimer’s Association Report (2022). 2022 Alzheimer’s disease facts and figures. *Alzheimer’s & Dementia* 18(4):700–789. doi:[10.1002/alz.12638](https://doi.org/10.1002/alz.12638).
- Amato, P.R. and Anthony, C.J. (2014). Estimating the effects of parental divorce and death with fixed effects models. *Journal of Marriage and Family* 76(2):370–386. doi:[10.1111/jomf.12100](https://doi.org/10.1111/jomf.12100).
- Bengtson, V.L. (2001). Beyond the nuclear family: The increasing importance of multigenerational bonds. *Journal of Marriage and Family* 63(1):1–16. doi:[10.1111/j.1741-3737.2001.00001.x](https://doi.org/10.1111/j.1741-3737.2001.00001.x).
- Caswell, H. (2019). The formal demography of kinship. *Demographic Research* 41:679–712.
- Caswell, H. (2022). The formal demography of kinship IV. *Demographic Research* 47:359–396.
- Caswell, H., Margolis, R., and Verdery, A.M. (2023). The formal demography of kinship v: Kin loss, bereavement, and causes of death. *SocArXiv*. doi:[10.31235/osf.io/mk64p](https://doi.org/10.31235/osf.io/mk64p).
- Caswell, H. and Song, X. (2021). The formal demography of kinship III. *Demographic Research* 45:517–546.
- Chen, S.H., Chen, Y.-C., and Liu, J.-T. (2009). The impact of unexpected maternal death on education: First evidence from three national administrative data links. *The American Economic Review* 99(2):149–153. doi:[10.1257/aer.99.2.149](https://doi.org/10.1257/aer.99.2.149).
- Dattani, S., Spooner, F., Ritchie, H., and Roser, M. (2023). *Causes of Death*. Our World in Data. <https://ourworldindata.org/causes-of-death>.
- Dudel, C. and Klüsener, S. (2021). Male–female fertility differentials across 17 high-income countries: Insights from a new data resource. *European Journal of Population* 37(2):417–441. doi:[10.1007/s10680-020-09575-9](https://doi.org/10.1007/s10680-020-09575-9).
- Hillis, Susan D., Blenkinsop, A., Villaveces, A., Annor, F.B., Liburd, L., Massetti, G.M., Demissie, Z., Mercy, J.A., Nelson III, C.A., Cluver, L., Flaxman, S., Sherr, L., Donnelly, C.A., Ratmann, O., and Unwin, H.J.T. (2021a). COVID-19-associated orphanhood and caregiver death in the United States. *Pediatrics* 148(6):e2021053760. doi:[10.1542/peds.2021-053760](https://doi.org/10.1542/peds.2021-053760).
- Hillis, Susan D., Unwin, H.J.T., Chen, Y., Cluver, L., Sherr, L., Goldman, P.S., Ratmann, O., Donnelly, C.A., Bhatt, S., Villaveces, A., Butchart, A., Bachman, G., Rawlings, L., Green, P., Nelson III, C.A., and Flaxman, S. (2021b). Global minimum estimates of children affected by COVID-19-associated orphanhood and deaths of caregivers: A modelling study. *The Lancet* 398(10298):391–402. doi:[10.1016/S0140-6736\(21\)01253-8](https://doi.org/10.1016/S0140-6736(21)01253-8).
- Institute for Health Metrics and Evaluation (2019). *GBD Results*. University of Washington. <https://vizhub.healthdata.org/gbd-results/>.
- Leopold, T. and Lechner, C.M. (2015). Parents’ death and adult well-being: Gender, age, and adaptation to filial bereavement. *Journal of Marriage and Family* 77(3):747–760. doi:[10.1111/jomf.12186](https://doi.org/10.1111/jomf.12186).
- Li, J., Vestergaard, M., Cnattingius, S., Gissler, M., Bech, B.H., Obel, C., and Olsen, J. (2014). Mortality after parental death in childhood: A nationwide cohort study from three nordic countries. *PLOS Medicine* 11(7):e1001679. doi:[10.1371/journal.pmed.1001679](https://doi.org/10.1371/journal.pmed.1001679).
- Luecken, L.J. (2008). Long-term consequences of parental death in childhood: Psychological and physiological manifestations. In: Stroebe, M. S., Hansson, R. O., Schut, H. and Stroebe, W. (eds.). *Handbook of Bereavement Research and Practice: Advances in Theory and Intervention*. Washington, D.C.: American Psychological Association: 397–416. doi:[10.1037/14498-019](https://doi.org/10.1037/14498-019).
- Patterson, S.E., Verdery, A.M., and Daw, J. (2020). Linked lives and childhood experience of family death on educational attainment. *Socius* 6:2378023120975594.
- Pham, S., Porta, G., Biernesser, C., Walker Payne, M., Iyengar, S., Melhem, N., and Brent,

- D.A. (2018). The burden of bereavement: Early-onset depression and impairment in youths bereaved by sudden parental death in a 7-year prospective study. *American Journal of Psychiatry* 175(9):887–896.
- Rizzi, S., Gampe, J., and Eilers, P.H. (2015). Efficient estimation of smooth distributions from coarsely grouped data. *American Journal of Epidemiology* 182(2):138–147.
- Saarela, J. and Rostila, M. (2019). Mortality after the death of a parent in adulthood: A register-based comparison of two ethno-linguistic groups. *European Journal of Public Health* 29(3):582–587. doi:[10.1093/eurpub/cky189](https://doi.org/10.1093/eurpub/cky189).
- Schoumaker, B. (2019). Male fertility around the world and over time: How different is it from female fertility? *Population and Development Review* 45:459–487. <https://about.jstor.org/terms>.
- Snyder, M., Alburez-Gutierrez, D., Williams, I., and Zagheni, E. (2022). Estimates from 31 countries show the significant impact of COVID-19 excess mortality on the incidence of family bereavement. *Proceedings of the National Academy of Sciences* 119(26):e2202686119. doi:[10.1073/pnas.2202686119](https://doi.org/10.1073/pnas.2202686119).
- United Nations (2022). *World Population Prospects: The 2022 Revision*.
- Verdery, A.M., Smith-Greenaway, E., Margolis, R., and Daw, J. (2020). Tracking the reach of COVID-19 kin loss with a bereavement multiplier applied to the United States. *Proceedings of the National Academy of Sciences* 117(30):17695–17701. doi:[10.1073/pnas.2007476117](https://doi.org/10.1073/pnas.2007476117).
- Zagheni, E. (2011). The impact of the HIV/AIDS epidemic on kinship resources for orphans in zimbabwe. *Population and Development Review* 37(4):761–783.