




Year; events / valid responses	Prevalence, % [95% CI]
All adults aged 18+	
2022 1716 / 26,490	6.11 [5.75, 6.48]
2025 1463 / 23,441	5.18 [4.84, 5.54]
With a history of stroke	
2022 140 / 959	12.49 [7.88, 19.23]
2025 157 / 862	21.15 [14.30, 30.12]
Without a history of stroke	
2022 1569 / 25,496	5.91 [5.56, 6.28]
2025 1305 / 22,551	4.88 [4.55, 5.23]




Population / outcome
Difference [95% CI]
All adults aged 18+
Swallowing difficulties
−0.93 [−1.43, −0.42]
Substantial difficulty
−0.40 [−0.62, −0.18]
Symptoms for ≥180 days
+0.10 [−0.12, 0.32]
With a history of stroke
Swallowing difficulties
+8.66 [−1.03, 18.35]
Substantial difficulty
+4.47 [−1.05, 9.98]
Symptoms for ≥180 days
+7.12 [−0.63, 14.87]
Without a history of stroke
Swallowing difficulties
−1.03 [−1.53, −0.53]
Substantial difficulty
−0.44 [−0.66, −0.22]
Symptoms for ≥180 days
+0.05 [−0.16, 0.26]
−5
0
5
10
15
20
Prevalence difference (percentage points)


Population / analysis method
Difference [95% CI]
All adults aged 18+
Crude prevalence
−0.75 [−1.24, −0.25]
Direct age standardization
−0.93 [−1.43, −0.42]
Age and sex adjustment
−0.73 [−1.21, −0.25]
Self-responses only: direct age standardization
−1.06 [−1.56, −0.56]
Self-responses only: age and sex adjustment
−0.88 [−1.35, −0.40]
With a history of stroke
Crude prevalence
+5.20 [0.89, 9.52]
Direct age standardization
+8.66 [−1.03, 18.35]
Age and sex adjustment
+5.32 [0.76, 9.88]
Self-responses only: direct age standardization
+4.03 [−5.59, 13.64]
Self-responses only: age and sex adjustment
+3.46 [−0.64, 7.56]
−10
−5
0
5
10
15
20
Prevalence difference (percentage points)


Age; events / valid responses
Difference [95% CI]
All adults aged 18+
18–44 years
2022: 538/9,547 2025: 329/8,238
−1.95 [−2.67, −1.24]
45–64 years
2022: 525/8,458 2025: 445/6,980
+0.04 [−0.79, 0.88]
65–74 years
2022: 339/4,864 2025: 317/4,443
+0.03 [−1.14, 1.19]
75–84 years
2022: 208/2,674 2025: 266/2,856
+1.55 [−0.12, 3.22]
85+ years
2022: 106/947 2025: 106/924
+0.08 [−3.49, 3.65]
With a history of stroke
18–44 years
2022: 5/52 2025: 11/55
+13.82 [−4.28, 31.93]
45–64 years
2022: 37/236 2025: 38/217
+0.87 [−6.64, 8.37]
65–74 years
2022: 42/316 2025: 47/241
+9.18 [1.55, 16.81]
75–84 years
2022: 38/239 2025: 34/243
+0.27 [−7.99, 8.53]
85+ years
2022: 18/116 2025: 27/106
+11.26 [−2.24, 24.77]
−10
0
10
20
30
40
Prevalence difference (percentage points)


Population / analysis assumption
Difference [95% CI]
All adults aged 18+
Main analysis: valid responses
−0.93 [−1.43, −0.42]
Quarters 1–3 only
−0.61 [−1.17, −0.05]
Missing responses: 2022 yes / 2025 no
−5.22 [−5.81, −4.63]
Missing responses: 2022 no / 2025 yes
+2.15 [1.56, 2.74]
With a history of stroke
Main analysis: valid responses
+8.66 [−1.03, 18.35]
Quarters 1–3 only
+10.46 [0.30, 20.63]
Missing responses: 2022 yes / 2025 no
+2.28 [−7.30, 11.85]
Missing responses: 2022 no / 2025 yes
+14.51 [4.99, 24.04]
−10
0
10
20
30
Prevalence difference (percentage points)