
Swallowing difficulties at age 65+

12-month prevalence by reported history of stroke




US 2000 age standard; survey weights, strata and clusters accounted for.

Source: NCHS, NHIS 2022 and 2025; authors' calculations.

Absolute differences between survey years

Age-standardized prevalence in 2025 minus 2022




Open markers: sparse age cells. Secondary outcomes and subgroups are exploratory.

Source: NCHS, NHIS 2022 and 2025; authors' calculations.

Sensitivity of the year comparison

Swallowing difficulties: 2025 minus 2022; two population groups




CIs are conditional on each assumption; extreme scenarios serve as bounds.

Source: NCHS, NHIS 2022 and 2025; authors' calculations.

Stroke-group comparison across analysis methods

Swallowing difficulties at age 65+: prevalence in 2025 minus 2022

Analysis / included responses	Difference [95% CI]
Primary: direct age standardization All valid responses	+6.27 [1.08; 11.47]
Model-based age and sex adjustment All valid responses	+5.84 [0.80; 10.88]
Direct age standardization Confirmed self-reports only	+4.53 [−0.42; 9.48]
Model-based age and sex adjustment Confirmed self-reports only	+4.05 [−0.78; 8.88]



Age: US 2000 standard. Adjusted margins: fixed 50:50 sex distribution within age groups.

Extensions are post-hoc; confirmed self-report restriction changes the target population.

Source: NCHS, NHIS 2022 and 2025; authors' calculations. Survey design accounted for.