NHIS 2025 / SUPPLEMENTARY FIGURE S2 / ADULTS AGED 65+

Model-standardized prevalence of loneliness

M3 logistic standardization | common analytic sample: N = 8,094; 429 events



Absolute prevalence difference
3.67 percentage points [95% CI 1.75, 5.60]

Marginal prevalence ratio
1.86 [95% CI 1.43, 2.42]

Frequent loneliness: always or usually lonely. Predictions standardized over the same weighted covariate distribution.
M3 adjusts for demographic, social and health characteristics plus in-person and phone/video contact.
Survey-logistic models; ten income imputations; predictive-margin uncertainty. Associations, not causal effects.

Source: NHIS 2025 public-use data; supplementary survey analysis

NHIS 2025 / SUPPLEMENTARY FIGURE S3 / ADULTS AGED 65+

The full distribution of loneliness responses

Available exposure and outcome responses: N = 8,231 | survey-weighted percentages



Points: survey-weighted percentages. Lines: Korn-Graubard 95% confidence intervals; sample sizes are unweighted.
All five categories are shown without covariate adjustment. Each exposure group sums to 100% before rounding.
All ten proportions pass the implemented NCHS precision criteria. Cross-sectional, self-reported responses.

Source: NHIS 2025 public-use data; supplementary survey analysis