

Supporting Information:

Velocity Map Imaging Spectroscopy of C_2H^- and C_2D^- : a benchmark study of vibronic coupling interactions

Benjamin A. Laws,^{*,†,‡} Zachariah D. Levey,[†] Andrei M. Sanov,[¶] John F.
Stanton,[§] Timothy W. Schmidt,[†] and Stephen T. Gibson[‡]

[†]*School of Chemistry, University of New South Wales, Sydney NSW 2052, Australia*

[‡]*Research School of Physics, The Australian National University, Canberra ACT 2601, Australia*

[¶]*Department of Chemistry and Biochemistry, The University of Arizona, Tucson, Arizona 85721,
United States*

[§]*Department of Chemistry, University of Florida, Gainesville, Florida 32611, United States*

E-mail: b.laws@unsw.edu.au

Contents

1	C_2H^- Spectral Assignments	S-2
2	C_2D^- Spectral Assignments	S-2
	References	S-5

1 C₂H[−] Spectral Assignments

Spectral assignments for all peaks resolved in the photoelectron spectra of C₂H[−] from this work are presented in Table S1. Peaks are labelled with respect to the photoelectron spectrum of C₂H[−] at 300 nm in Figure S1. The experimental binding energy of each transition is given, alongside the anisotropy (+/−), the corresponding vibronic symmetry, and the calculated energy from Ref. S1. Assignments are given as $\tilde{X}(v_1, v_2, v_3)\tilde{A}(v_1, v_2, v_3)$ where superscripts represent hot-band transitions.




Figure S1: Photoelectron spectrum of C₂H[−] at 300 nm, showing the labelling of peaks used in this work.

2 C₂D[−] Spectral Assignments

Spectral assignments for all peaks resolved in C₂D[−] from this work are presented in Table S2. Peaks are labelled with respect to the photoelectron spectrum of C₂D[−] at 355 nm in Figure S2. The experimental binding energy of each transition is given, alongside the anisotropy (+/−), the corresponding vibronic symmetry, and the calculated energy from Ref. S1.

Table S1: Peak positions (cm^{-1}), and assignments for the C_2H^- photoelectron spectra from this work. The sign of the anisotropy parameter is shown for each transition, along with it's vibronic symmetry, and the calculated position from Ref.^{S1}.

Peak	eBE (cm^{-1})	v (cm^{-1})	β	Symmetry	v_{calc}^a	Assignment
	23 591	-231	+	Σ^+		$\tilde{X}(0, 2^2, 0)$
	23 685	-137	+	Σ^+		$\tilde{X}(0, 1^1, 0)$
	23 823	0	+	Σ^+	0	$\tilde{X}(0, 0, 0)$
	24 184	361	−	Π	371	$\tilde{X}(0, 1, 0)$
	24 630	807	+	Σ^+	794	$\tilde{X}(0, 2, 0)$
	25 663	1840	+	Σ^+	1838	$\tilde{X}(0, 0, 1)$
	25 916	2093	−	Π	2096	$\tilde{X}(0, 1, 1)$
	25 993	2170	−	Π	2166	$\tilde{X}(0, 5, 0)$
	26 362	2539	+	Σ^+	2536	$\tilde{X}(0, 2, 1)$
	26 757	2934	−	Π	2933	$\tilde{X}(0, 3, 1)$
	26 929	3106	−	Π	3104	$\tilde{X}(0, 7, 0)$
	27 175	3352	+	Σ^+	3371	$\tilde{X}(0, 4, 1)$
a	27 430	3607	−	Π	3604	$\tilde{X}(0, 1, 2)\tilde{X}(1, 1, 0)\tilde{A}(0, 0, 0)$
b	27 515	3692	−	Π	3690	$\tilde{X}(1, 1, 0)\tilde{X}(0, 1, 2)\tilde{A}(0, 0, 0)$
c	27 612	3788	−	Π	3790	$\tilde{X}(0, 5, 1)\tilde{A}(0, 0, 0)$
d	27 851	4028	−	Π	4011	$\tilde{X}(0, 9, 0)\tilde{X}(0, 5, 1)\tilde{A}(0, 0, 0)$
e	27 941	4118	−	Π	4093	$\tilde{X}(0, 9, 0)\tilde{A}(0, 0, 0)$
f	28 200	4377	+	Σ^+	4375	$\tilde{X}(0, 6, 1)$
g	28 349	4526	+	Σ^+	4524	$\tilde{X}(0, 10, 0)$
h	28 423	4600	−	Π	4593	$\tilde{X}(0, 3, 2)\tilde{A}(0, 0, 0)$
i	28 562	4739	−	Π	4702	$\tilde{X}(0, 7, 1)\tilde{A}(0, 1, 0)$
j	28 714	4891	−	Π	4879	$\tilde{X}(0, 7, 1)$
k	28 834	5011	−	Π	5004	$\tilde{X}(0, 11, 0)$
l	29 049	5226	−	Π	5222	$\tilde{X}(0, 1, 3)\tilde{A}(0, 0, 1)$
m	29 227	5404	+	Σ^+	5406	$\tilde{X}(0, 12, 0)\tilde{A}(0, 1, 0)$
n	29 283	5460	−	Π	5445	$\tilde{X}(0, 5, 2)\tilde{A}(0, 0, 1)$
o	29 465	5642	−	Π	5630	$\tilde{X}(0, 9, 1)\tilde{A}(0, 0, 1)$
p	29 740	5917	−	Π	5914	$\tilde{X}(0, 13, 0)$
q	29 844	6021	+	Σ^+	6054	$\tilde{X}(0, 6, 2)\tilde{X}(0, 2, 3)$
r	30 021	6198	−	Π	6200	$\tilde{X}(0, 3, 3)\tilde{A}(0, 2, 0)$
s	30 086	6263	−	Π	6266	$\tilde{X}(1, 7, 0)$

^a from calculations of Ref. S1

Table S2: Peak positions (cm^{-1}) and assignments for the C_2D^- photoelectron spectra from this work. The sign of the anisotropy parameter is shown for each transition, along with it's vibronic symmerty, and calculated position from Ref.^{S1}.

Peak	eBE (cm^{-1})	v (cm^{-1})	β	Symmetry	v_{calc}^a	Assignment
	23 751	-197	+	Σ^+		$\tilde{X}(0, 2^2, 0)$
	23829	-120	+	Σ^+		$\tilde{X}(0, 2^1, 0)$
	23 949	0	+	Σ^+	0	$\tilde{X}(0, 0, 0)$
	24 227	278	—	Π	287	$\tilde{X}(0, 1, 0)$
	24 577	629	+	Σ^+	615	$\tilde{X}(0, 2, 0)$
	24 906	957	—	Π	953	$\tilde{X}(0, 3, 0)$
	25 506	1557	+			
	25 688	1740	+	Σ^+	1744	$\tilde{X}(0, 0, 1)$
	25 901	1952	—	Π	1962	$\tilde{X}(0, 1, 1)$
	26 245	2297	+	Σ^+	2302	$\tilde{X}(0, 2, 1)$
	26 556	2607	—	Π	2612	$\tilde{X}(0, 3, 1)$
	26 747	2798	—	Π	2796	$\tilde{X}(1, 1, 0)$
a	27 253	3304	—	Π	3309	$\tilde{X}(0, 5, 1)$
b	27 369	3420	—	Π	3426	$\tilde{X}(1, 3, 0)$
c	27 450	3501	—	Π	3511	$\tilde{X}(0, 1, 2)$
d	27 552	3603	—			
e	27 793	3844	—	Π	3838	$\tilde{A}(0, 0, 0)$
f	27 906	3957	+	Σ^+	3967	$\tilde{X}(0, 2, 2)$

^a from calculations of Ref. S1




Figure S2: Photoelectron spectrum of C_2D^- at 266 nm, showing the labelling of peaks used in this work.

References

- (S1) Tarroni, R.; Carter, S. Theoretical calculation of vibronic levels of C_2H and C_2D to 10,000 cm^{-1} . *The Journal of Chemical Physics* **2003**, *119*, 12878–12889.