

Supporting Information:

PAH growth in flames and space: phenalenyl radical from acenaphthylene

Zachariah D. Levey,[†] Benjamin A. Laws,[†] Srivathsan P. Sundar,[‡] Klaas Nauta,[†] Scott H. Kable,[†] Gabriel da Silva,[‡] John F. Stanton,[¶] and Timothy W. Schmidt^{*,§}

[†]*School of Chemistry, University of New South Wales, Sydney NSW 2052, Australia*

[‡]*Department of Chemical Engineering, The University of Melbourne, Parkville 3010, Australia*

[¶]*Department of Chemistry, University of Florida, Gainesville, Florida 32611, USA*

[§]*Centre of Excellence in Exciton Science, University of New South Wales, Sydney NSW 2052, Australia*

E-mail: timothy.schmidt@unsw.edu.au

Reaction schemes for CH insertion

As there are three unique H atoms on the six-membered rings of ACYN, C-H bond insertion may proceed via three distinct pathways, each leading to a unique tropzyl-like RSR intermediate. One of these pathways, for CH insertion at the 1-position, was presented in Figure 7. The pathways (and corresponding tropyl-like intermediates) for CH insertion at the 2-position and 3-position are shown below in Figures S1 and S2.




Figure S1: Reaction scheme for CH insertion at the 2-position, forming the second tropylium-like RSR. Calculations were performed at MO6-2X/6-31G(2df,p) level of theory. The energies are in kcal/mol.




Figure S2: Reaction scheme for CH insertion at the 3-position, forming the third tropylium-like RSR. Calculations were performed at MO6-2X/6-31G(2df,p) level of theory. The energies are in kcal/mol.