

Formation of PhenRad via CH₂ ring insertion...

Z. Levey, B. A. Laws,* K. Nauta, and T. Schmidt

School of Chemistry, University of New South Wales, Sydney NSW 2052, Australia

E-mail: B.Laws@unsw.edu.au

Abstract

This is the manuscripts abstract...

Introduction

Intro....^{1,2}

Results

Here is a figure. You can then refer to the figure using Fig. 1.




Figure 1: A figure showing the depletion rate for different laser delays.

Discussion

Equations may be written using general LaTeX format, and the same labelling method.

$$I(\theta, \epsilon) = \frac{\sigma_{\text{total}}(\epsilon)}{4\pi} [1 + \beta(\epsilon)P_2(\cos \theta)], \quad (1)$$

where β in Eq. (1) is a really cool parameter.

We can also make tables, like this one, Table 1.

Table 1: This is my table caption.

Peak	eBE (cm ⁻¹)	ν (cm ⁻¹)	β	Symmetry
	23 591	-231	+	Σ^+

Conclusion

Experimental Details

Acknowledgement This research was supported by the Australian Research Council Discovery Project Grant DP160102585. The author's thank Andrei Sanov for discussion on his modified Cooper-Zare anisotropy model.

References

- (1) Porfiriev, D. P.; Azyazov, V. N.; Mebel, A. M. Conversion of acenaphthalene to phenalene via methylation: A theoretical study. *Combustion and Flame* **2020**, *213*, 302 – 313.
- (2) Laws, B. A.; Cavanagh, S. J.; Lewis, B. R.; Gibson, S. T. NOO peroxy isomer exposed with velocity-map imaging. *J. Phys. Chem. Lett.* **2017**, *8*, 4397–4401.