

Formation of PhenRad via CH ring insertion...

Z. Levey, B. A. Laws,* K. Nauta, and T. Schmidt

School of Chemistry, University of New South Wales, Sydney NSW 2052, Australia

E-mail: B.Laws@unsw.edu.au

Abstract

This is the manuscripts abstract...

Introduction

The formation mechanism of polycyclic aromatic hydrocarbons (PAHs) is an increasingly important area of research due to their role in combustion, atmospheric, circumstellar, and interstellar chemistry. Incomplete combustion of hydrocarbon fuels results in the formation of soot particles which, when released into the atmosphere, contribute to global warming and environmental damage.¹⁻⁴ Inhalation and ingestion of these particles is also of concern for human health due to their toxic and carcinogenic properties.⁵⁻⁸ PAHs are suspected to play the role of early intermediates in the soot nucleation process.^{9,10} Comparably with terrestrial soot, PAHs are viewed as the initial precursors of stellar dust grains being formed in the circumstellar shells of carbon rich evolved stars.¹¹⁻¹⁴ The most convincing evidence for the presence of PAHs in the interstellar medium (ISM) comes from infrared emission features termed the aromatic infrared bands (AIBs). These strong features between 3-20um are ubiquitous in the spectra of interstellar objects and are characteristic of large PAH molecules.^{15,16} Modelling based on these optical observations allows the inference of chemical abundances and size distributions with an estimated 20% of the total cosmic carbon abundance locked

up in PAHs of sizes 50—100 C atoms.^{17,18} PAHs are also strongly considered as carriers of the diffuse interstellar bands (DIBs) – a series of unidentified absorption features in the visible to near-infrared region.¹⁹ Aromatic species such as benzene and benzonitrile, as well as fullerenes, C₆₀ (in its neutral and cationic state) and C₇₀, have been identified in the ISM through spectroscopic techniques,^{20–23} however, there has not yet been an identification of PAHs in the ISM.

Due to the diversity of physical conditions within the ISM, the formation mechanism for interstellar PAHs is ambiguous. Current astrochemical models of PAH formation are predominately derived from combustion chemistry models. Bottom-up molecular growth processes in hot and dense circumstellar envelopes of carbon-rich asymptotic giant branch stars are largely based on the hydrogen abstraction-acetylene addition (HACA) mechanism.¹⁵ HACA was first described as an activation of a radical site by H-abstraction/addition followed by sequential addition of acetylene, finally resulting in ring closure²⁴ and has been shown to occur experimentally for the formation of naphthalene from benzene via the phenylacetylene intermediate.^{25,26} However, there are some inherent shortcomings of this mechanism. Notably, modelling studies have reported that HACA under-predicts the concentration of PAHs in flames.²⁷ Also, the injection of PAHs from stellar sources occurs on a time scale that is significantly longer than the destruction of PAHs caused by interstellar shockwaves and bombardment by energetic cosmic rays suggesting that crucial PAH growth routes are missing within the astrochemical model.^{28–30} Alternate PAH formation mechanisms have been proposed in conjunction with HACA to account for the deficiencies in PAH concentrations. These include the recombination of resonance stabilized radicals (RSRs),^{31–33} condensation of small PAHs,³⁴ phenyl addition/cyclization (PAC)³⁵ and methyl addition/cyclization (MAC).³⁶

Acenaphthylene (ACYN) is considered the first significant island of stability “pulling” the HACA sequence forward.³⁷ Theoretical investigations of the reaction between 1-naphthyl radical and acetylene at elevated temperatures above 1000 K show predominant formation of ACYN over phenanthrene and anthracene. Rapid cyclization after the addition of a single

acetylene molecule adjacent to the bay region of a PAH to form a five-membered ring occurs much faster than the addition of a second acetylene, followed by cyclization to form a third aromatic ring.³⁸ It was shown that PAHs with exclusively six-membered rings accounted for $\leq 6\%$ of the total yield from the reaction of naphthalene and acetylene, with most products containing a five-membered ring ($\sim 75\%$). Hence, a mechanism converting a five-member ring on the edge of a PAH molecule to a six-member ring is necessary to explain reaction pathways to tricyclic and larger PAHs.

Ring expansion has been thoroughly investigated for an isolated five-member ring, the cyclopentadienyl radical,^{31,39,40} and a five-member ring attached to benzene, the 1-indenyl radical.⁴¹⁻⁴³ This reaction can readily occur by methylation (addition of a CH_3 radical) to form benzene and naphthalene, respectively. Additionally, expansion of a five-member ring to a six-member ring has been demonstrated through the reaction of pyrrole with methyldi- dyne radical (CH) to form pyridine.⁴⁴ The CH radical in its ground state has been detected in combustion environments,⁴⁵⁻⁴⁷ the interstellar medium⁴⁸ and under plasma conditions.⁴⁹ Reactions between the CH radical and small unsaturated hydrocarbons and carbonyls can result in CH insertion to the $\text{C}-\text{C}$ π -bond through a cyclic intermediate (cycloaddition) followed by ring-opening isomerization.⁵⁰⁻⁵³ Isotopomer distribution experiments provided additional evidence for this cycloaddition mechanism through the reactions of deuterated methyldi- dyne radical (CD) with ethylene and pyrrole.^{44,50} Recently, ab initio calculations in combination with RRKM-ME calculations for the reaction of 1-acenaphthyl and methyl radicals exhibit ring expansion of the five-member ring by methylation to produce the phenalenyl radical.⁵⁴ The phenalenyl radical is an open-shell, neutral, RSR consisting of three benzene rings fused together by a central carbon atom. It is considered a prototypical open-shell graphene fragment⁵⁵ and is posited to have a role in soot inception and growth.³³ The gas phase, resonance enhanced multi-photon ionisation (REMPI) spectrum for the $\text{D}_0 \rightarrow \text{D}_1$ state of the phenalenyl radical has been previously recorded.⁵⁶

In this work, we experimentally investigated the ring-expansion PAH growth mechanism

for ACYN to phenalenyl radical. ACYN and methane (CH_4) molecules interact and undergo subsequent radical gas-phase reactions induced by means of an electrical discharge. This allowed for the simultaneous examination of the methylation and cycloaddition mechanisms previously described. The reaction products and intermediates were identified using the mass selective REMPI technique in conjunction with isotope labelling experiments. We determined that ACYN formed the phenalenyl radical through a cycloaddition CH insertion reaction.

Results

Here is a figure. You can then refer to the figure using Fig. 2.




Figure 1: A figure showing the depletion rate for different laser delays.

Discussion

Equations may be written using general LaTeX format, and the same labelling method.

$$I(\theta, \epsilon) = \frac{\sigma_{\text{total}}(\epsilon)}{4\pi} [1 + \beta(\epsilon) P_2(\cos \theta)], \quad (1)$$

where β in Eq. (1) is a really cool parameter.




Figure 2: A figure showing the depletion rate for different laser delays.

We can also make tables, like this one, Table 1.

Table 1: This is my table caption.

Peak	eBE (cm^{-1})	ν (cm^{-1})	β	Symmetry
	23 591	-231	+	Σ^+

Conclusion

Experimental Details

Acknowledgement

This research was supported by the Australian Research Council Discovery Project Grant DP160102585. The author's thank Andrei Sanov for discussion on his modified Cooper-Zare anisotropy model.

References

- (1) Bond, T. C.; Doherty, S. J.; Fahey, D. W.; Forster, P. M.; Berntsen, T.; Deangelo, B. J.; Flanner, M. G.; Ghan, S.; Kärcher, B.; Koch, D.; et al., Bounding the role of black carbon in the climate system: A scientific assessment. *Journal of Geophysical Research Atmospheres* **2013**, *118*, 5380–5552.
- (2) Ramanathan, V.; Carmichael, G. Global and regional climate changes due to black carbon. *Nature Geoscience* **2008**, *1*, 221–227.
- (3) Shrestha, G.; Traina, S.; Swanston, C. Black Carbon’s Properties and Role in the Environment: A Comprehensive Review. *Sustainability* **2010**, *2*, 294–320.
- (4) Finlayson-Pitts, B. J. Tropospheric Air Pollution: Ozone, Airborne Toxics, Polycyclic Aromatic Hydrocarbons, and Particles. *Science* **1997**, *276*, 1045–1051.
- (5) Tiwari, M.; Sahu, S. K.; Pandit, G. G. Inhalation Risk Assessment of PAH Exposure Due to Combustion Aerosols Generated from Household Fuels. *Aerosol and Air Quality Research* **2015**, *15*, 582–590.
- (6) Tiwari, M.; Sahu, S.; Pandit, G. Distribution of PAHs in different compartment of creek ecosystem: Ecotoxicological concern and human health risk. *Environmental Toxicology and Pharmacology* **2017**, *50*, 58 – 66.
- (7) Lahaye, J. *Soot in Combustion Systems and Its Toxic Properties*; Nato Conference Series; Springer US, 2013.
- (8) Amegah, A. K.; Quansah, R.; Jaakkola, J. J. K. Household Air Pollution from Solid Fuel Use and Risk of Adverse Pregnancy Outcomes: A Systematic Review and Meta-Analysis of the Empirical Evidence. *PLOS ONE* **2014**, *9*, 1–23.
- (9) Frenklach, M. Reaction mechanism of soot formation in flames. *Phys. Chem. Chem. Phys.* **2002**, *4*, 2028–2037.

- (10) Wang, H. Formation of nascent soot and other condensed-phase materials in flames. *Proceedings of the Combustion Institute* **2011**, *33*, 41 – 67.
- (11) Henning, T.; Salama, F. Carbon in the Universe. *Science* **1998**, *282*, 2204–2210.
- (12) Jäger, C.; Huisken, F.; Mutschke, H.; Jansa, I. L.; Henning, T. FORMATION OF POLYCYCLIC AROMATIC HYDROCARBONS AND CARBONACEOUS SOLIDS IN GAS-PHASE CONDENSATION EXPERIMENTS. *The Astrophysical Journal* **2009**, *696*, 706–712.
- (13) Draine, B. T.; Li, A. Infrared Emission from Interstellar Dust. I. Stochastic Heating of Small Grains. *The Astrophysical Journal* **2001**, *551*, 807–824.
- (14) Draine, B. T.; Li, A. Infrared Emission from Interstellar Dust. IV. The Silicate-Graphite-PAH Model in the Post-SpitzerEra. *The Astrophysical Journal* **2007**, *657*, 810–837.
- (15) Tielens, A. G. G. M. The molecular universe. *Reviews of Modern Physics* **2013**, *85*, 1021–1081.
- (16) Pech, C.; Joblin, C.; Boissel, P., The profiles of the aromatic infrared bands explained with molecular carriers. *A&A* **2002**, *388*, 639–651.
- (17) Tielens, A. Interstellar Polycyclic Aromatic Hydrocarbon Molecules. *Annual Review of Astronomy and Astrophysics* **2008**, *46*, 289–337.
- (18) Dwek, E.; Arendt, R. G.; Fixsen, D. J.; Sodroski, T. J.; Odegard, N.; Weiland, J. L.; Reach, W. T.; Hauser, M. G.; Kelsall, T.; Moseley, S. H.; et al., Detection and Characterization of Cold Interstellar Dust and Polycyclic Aromatic Hydrocarbon Emission, from COBE Observations. *The Astrophysical Journal* **1997**, *475*, 565–579.
- (19) Cox, N. L.J., The PAH-DIB Hypothesis. *EAS Publications Series* **2011**, *46*, 349–354.

- (20) Cernicharo, J.; Heras, A. M.; Tielens, A. G. G. M.; Pardo, J. R.; Herpin, F.; Guélin, M.; Waters, L. B. F. M. [ITAL]Infrared Space Observatory's[/ITAL] Discovery of C[TINF]4[/TINF]H[TINF]2[/TINF], C[TINF]6[/TINF]H[TINF]2[/TINF], and Benzene in CRL 618. *The Astrophysical Journal* **2001**, *546*, L123–L126.
- (21) McGuire, B. A.; Burkhardt, A. M.; Kalenskii, S.; Shingledecker, C. N.; Remijan, A. J.; Herbst, E.; McCarthy, M. C. Detection of the aromatic molecule benzonitrile (c-C₆H₅CN) in the interstellar medium. *Science* **2018**, *359*, 202–205.
- (22) Cami, J.; Bernard-Salas, J.; Peeters, E.; Malek, S. E. Detection of C₆₀ and C₇₀ in a Young Planetary Nebula. *Science* **2010**, *329*, 1180–1182.
- (23) Campbell, E. K.; Holz, M.; Gerlich, D.; Maier, J. P. Laboratory confirmation of C₆₀+ as the carrier of two diffuse interstellar bands. *Nature* **2015**, *523*, 322–323.
- (24) Frenklach, M.; Clary, D. W.; Gardiner, W. C.; Stein, S. E. Detailed kinetic modeling of soot formation in shock-tube pyrolysis of acetylene. *Symposium (International) on Combustion* **1985**, *20*, 887 – 901, Twentieth Symposium (International) on Combustion.
- (25) Parker, D. S. N.; Kaiser, R. I.; Troy, T. P.; Ahmed, M. Hydrogen Abstraction/Acetylene Addition Revealed. *Angewandte Chemie International Edition* **2014**, *53*, 7740–7744.
- (26) Yang, T.; Troy, T. P.; Xu, B.; Kostko, O.; Ahmed, M.; Mebel, A. M.; Kaiser, R. I. Hydrogen-Abstraction/Acetylene-Addition Exposed. *Angewandte Chemie International Edition* **2016**, *55*, 14983–14987.
- (27) Raj, A.; Prada, I. D. C.; Amer, A. A.; Chung, S. H. A reaction mechanism for gasoline surrogate fuels for large polycyclic aromatic hydrocarbons. *Combustion and Flame* **2012**, *159*, 500 – 515.
- (28) Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M., Polycyclic aromatic hydrocarbon processing in a hot gas. *A&A* **2010**, *510*, A37.

- (29) Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M., Polycyclic aromatic hydrocarbon processing in interstellar shocks. *A&A* **2010**, *510*, A36.
- (30) Micelotta, E. R.; Jones, A. P.; Tielens, A. G. G. M., Polycyclic aromatic hydrocarbon processing by cosmic rays. *A&A* **2011**, *526*, A52.
- (31) Melius, C. F.; Colvin, M. E.; Marinov, N. M.; Pit, W. J.; Senkan, S. M. Reaction mechanisms in aromatic hydrocarbon formation involving the C₅H₅ cyclopentadienyl moiety. *Symposium (International) on Combustion* **1996**, *26*, 685 – 692.
- (32) Miller, J. A.; Melius, C. F. Kinetic and thermodynamic issues in the formation of aromatic compounds in flames of aliphatic fuels. *Combustion and Flame* **1992**, *91*, 21 – 39.
- (33) Johansson, K. O.; Head-Gordon, M. P.; Schrader, P. E.; Wilson, K. R.; Michelsen, H. A. Resonance-stabilized hydrocarbon-radical chain reactions may explain soot inception and growth. *Science* **2018**, *361*, 997–1000.
- (34) Siegmann, K.; Sattler, K. Formation mechanism for polycyclic aromatic hydrocarbons in methane flames. *The Journal of Chemical Physics* **2000**, *112*, 698–709.
- (35) Shukla, B.; Susa, A.; Miyoshi, A.; Koshi, M. Role of Phenyl Radicals in the Growth of Polycyclic Aromatic Hydrocarbons. *The Journal of Physical Chemistry A* **2008**, *112*, 2362–2369, PMID: 18298104.
- (36) Shukla, B.; Miyoshi, A.; Koshi, M. Role of Methyl Radicals in the Growth of PAHs. *Journal of the American Society for Mass Spectrometry* **2010**, *21*, 534 – 544.
- (37) Frenklach, M.; Mebel, A. M. On the mechanism of soot nucleation. *Phys. Chem. Chem. Phys.* **2020**, *22*, 5314–5331.
- (38) Kislov, V. V.; Sadovnikov, A. I.; Mebel, A. M. Formation Mechanism of Polycyclic

- Aromatic Hydrocarbons beyond the Second Aromatic Ring. *The Journal of Physical Chemistry A* **2013**, *117*, 4794–4816, PMID: 23672431.
- (39) Moskaleva, L.; Mebel, A.; Lin, M. The $\text{CH}_3 + \text{C}_5\text{H}_5$ reaction: A potential source of benene at high temperatures. *Symposium (International) on Combustion* **1996**, *26*, 521 – 526.
- (40) Sharma, S.; Green, W. H. Computed Rate Coefficients and Product Yields for $\text{c-C}_5\text{H}_5 + \text{CH}_3 \rightarrow \text{Products}$. *The Journal of Physical Chemistry A* **2009**, *113*, 8871–8882, PMID: 19591475.
- (41) Mebel, A. M.; Georgievskii, Y.; Jasper, A. W.; Klippenstein, S. J. Pressure-dependent rate constants for PAH growth: formation of indene and its conversion to naphthalene. *Faraday Discussions* **2016**, *195*, 637–670.
- (42) Jasper, A. W.; Hansen, N. Hydrogen-assisted isomerizations of fulvene to benzene and of larger cyclic aromatic hydrocarbons. *Proceedings of the Combustion Institute* **2013**, *34*, 279 – 287.
- (43) Zhao, L.; Kaiser, R. I.; Lu, W.; Xu, B.; Ahmed, M.; Morozov, A. N.; Mebel, A. M.; Howlader, A. H.; Wnuk, S. F. Molecular mass growth through ring expansion in polycyclic aromatic hydrocarbons via radical–radical reactions. *Nature Communications* **2019**, *10*.
- (44) Soorkia, S.; Taatjes, C. A.; Osborn, D. L.; Selby, T. M.; Trevitt, A. J.; Wilson, K. R.; Leone, S. R. Direct detection of pyridine formation by the reaction of CH (CD) with pyrrole: a ring expansion reaction. *Physical Chemistry Chemical Physics* **2010**, *12*, 8750.
- (45) Love, N.; Parthasarathy, R. N.; Gollahalli, S. R. Concentration Measurements of CH and OH Radicals in Laminar Biofuel Flames. *International Journal of Green Energy* **2011**, *8*, 113–120.

- (46) Tinaut, F. V.; Reyes, M.; Giménez, B.; Pastor, J. V. Measurements of OH* and CH* Chemiluminescence in Premixed Flames in a Constant Volume Combustion Bomb under Autoignition Conditions. *Energy & Fuels* **2011**, *25*, 119–129.
- (47) Zhang, T.; Guo, Q.; Liang, Q.; Dai, Z.; Yu, G. Distribution Characteristics of OH*, CH*, and C2* Luminescence in CH₄/O₂ Co-flow Diffusion Flames. *Energy & Fuels* **2012**, *26*, 5503–5508.
- (48) Gerin, M., et al. Interstellar CH absorption in the diffuse interstellar medium along the sight-lines to G10.6-0.4 (W31C), W49N, and W51. *A&A* **2010**, *521*, L16.
- (49) Zhou, J.; Fisher, E. R. Surface Reactivity and Energetics of CH Radicals during Plasma Deposition of Hydrogenated Diamondlike Carbon Films. *The Journal of Physical Chemistry B* **2006**, *110*, 21911–21919, PMID: 17064158.
- (50) Goulay, F.; Trevitt, A. J.; Meloni, G.; Selby, T. M.; Osborn, D. L.; Taatjes, C. A.; Vereecken, L.; Leone, S. R. Cyclic Versus Linear Isomers Produced by Reaction of the Methylidyne Radical (CH) with Small Unsaturated Hydrocarbons. *Journal of the American Chemical Society* **2009**, *131*, 993–1005, PMID: 19123915.
- (51) Trevitt, A. J.; Prendergast, M. B.; Goulay, F.; Savee, J. D.; Osborn, D. L.; Taatjes, C. A.; Leone, S. R. Product Branching Fractions of the CH + Propene Reaction from Synchrotron Photoionization Mass Spectrometry. *The Journal of Physical Chemistry A* **2013**, *117*, 6450–6457.
- (52) Goulay, F.; Trevitt, A. J.; Savee, J. D.; Bouwman, J.; Osborn, D. L.; Taatjes, C. A.; Wilson, K. R.; Leone, S. R. Product Detection of the CH Radical Reaction with Acetaldehyde. *The Journal of Physical Chemistry A* **2012**, *116*, 6091–6106.
- (53) Trevitt, A. J.; Goulay, F. Insights into gas-phase reaction mechanisms of small carbon radicals using isomer-resolved product detection. *Phys. Chem. Chem. Phys.* **2016**, *18*, 5867–5882.

- (54) Porfiriev, D. P.; Azyazov, V. N.; Mebel, A. M. Conversion of acenaphthalene to phenylene via methylation: A theoretical study. *Combustion and Flame* **2020**, *213*, 302 – 313.
- (55) Morita, Y.; Suzuki, S.; Sato, K.; Takui, T. Synthetic organic spin chemistry for structurally well-defined open-shell graphene fragments. *Nature Chemistry* **2011**, *3*, 197–204.
- (56) O'Connor, G. D.; Troy, T. P.; Roberts, D. A.; Chalyavi, N.; Fückel, B.; Crossley, M. J.; Nauta, K.; Stanton, J. F.; Schmidt, T. W. Spectroscopy of the Free Phenalenyl Radical. *Journal of the American Chemical Society* **2011**, *133*, 14554–14557, PMID: 21854045.