

# Formation of PhenRad via CH<sub>2</sub> ring insertion...

Z. Levey, B. A. Laws,\* K. Nauta, and T. Schmidt

*School of Chemistry, University of New South Wales, Sydney NSW 2052, Australia*

E-mail: B.Laws@unsw.edu.au

## Abstract

This is the manuscripts abstract...

## Introduction

Intro....<sup>1,2</sup> oconnor paper says...<sup>3</sup>

## Results

Here is a figure. You can then refer to the figure using Fig. 1.




Figure 1: A figure showing the depletion rate for different laser delays.

## Discussion

Equations may be written using general LaTeX format, and the same labelling method.

$$I(\theta, \epsilon) = \frac{\sigma_{\text{total}}(\epsilon)}{4\pi} [1 + \beta(\epsilon)P_2(\cos \theta)], \quad (1)$$

where  $\beta$  in Eq. (1) is a really cool parameter.

We can also make tables, like this one, Table 1.

Table 1: This is my table caption.

Peak	eBE (cm <sup>-1</sup> )	$\nu$ (cm <sup>-1</sup> )	$\beta$	Symmetry
23	591	-231	+	$\Sigma^+$

## Conclusion

## Experimental Details

**Acknowledgement** This research was supported by the Australian Research Council Discovery Project Grant DP160102585. The author's thank Andrei Sanov for discussion on his modified Cooper-Zare anisotropy model.

## References

- (1) Porfiriev, D. P.; Azyazov, V. N.; Mebel, A. M. Conversion of acenaphthalene to phenalene via methylation: A theoretical study. *Combustion and Flame* **2020**, *213*, 302 – 313.
- (2) Laws, B. A.; Cavanagh, S. J.; Lewis, B. R.; Gibson, S. T. NOO peroxy isomer exposed with velocity-map imaging. *J. Phys. Chem. Lett.* **2017**, *8*, 4397–4401.
- (3) O'Connor, G. D.; Troy, T. P.; Roberts, D. A.; Chalyavi, N.; Fückel, B.; Crossley, M. J.; Nauta, K.; Stanton, J. F.; Schmidt, T. W. Spectroscopy of the Free Phenalenyl Radical. *Journal of the*

*American Chemical Society* **2011**, *133*, 14554–14557, PMID: 21854045.