


Notes on Trigonometry

Paul F. Roysdon, Ph.D.

I. DEFINITIONS

Right triangle definition

For this definition we assume that $0 < \theta < \frac{\pi}{2}$ or $0^\circ < \theta < 90^\circ$.




$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}} \quad \csc \theta = \frac{\text{hypotenuse}}{\text{opposite}}$$

$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} \quad \sec \theta = \frac{\text{hypotenuse}}{\text{adjacent}}$$

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} \quad \cot \theta = \frac{\text{adjacent}}{\text{opposite}}$$

Unit circle definition

For this definition θ is any angle.



$$\sin \theta = \frac{y}{1} = y \quad \csc \theta = \frac{1}{y}$$

$$\cos \theta = \frac{x}{1} = x \quad \sec \theta = \frac{1}{x}$$

$$\tan \theta = \frac{y}{x} \quad \cot \theta = \frac{x}{y}$$

II. FACTS AND PROPERTIES

Domain

The domain is all the values of θ that can be plugged into the function.

$\sin \theta$, θ can be any angle.

$\cos \theta$, θ can be any angle.

$\tan \theta$, $\theta \neq (n + \frac{1}{2})\pi, n = 0, \pm 1, \pm 2, \dots$

$\csc \theta$, $\theta \neq n\pi, n = 0, \pm 1, \pm 2, \dots$

$\sec \theta$, $\theta \neq (n + \frac{1}{2})\pi, n = 0, \pm 1, \pm 2, \dots$

$\cot \theta$, $\theta \neq n\pi, n = 0, \pm 1, \pm 2, \dots$

Range

The range is all possible values to get out of the function.

$$-1 \leq \sin \leq 1$$

$$-1 \leq \cos \leq 1$$

$$-\infty \leq \tan \leq \infty$$

$$\csc \theta \geq 1 \text{ and } \csc \theta \leq -1$$

$$\sec \theta \geq 1 \text{ and } \sec \theta \leq -1$$

$$-\infty \leq \cot \leq \infty$$

Period

The period of a function is the number T such that $f(\theta + T) = f(\theta)$. If ω is a fixed number and θ is any angle, the following periods exist.

$$\sin(\omega\theta) \implies T = \frac{2\pi}{\omega} \quad \csc(\omega\theta) \implies T = \frac{2\pi}{\omega}$$

$$\cos(\omega\theta) \implies T = \frac{2\pi}{\omega} \quad \sec(\omega\theta) \implies T = \frac{2\pi}{\omega}$$

$$\tan(\omega\theta) \implies T = \frac{\pi}{\omega} \quad \cot(\omega\theta) \implies T = \frac{\pi}{\omega}$$

III. FORMULAS AND IDENTITIES

Tangent and Cotangent Identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\cot \theta = \frac{\cos \theta}{\sin \theta}$$

Reciprocal Identities

$$\sin \theta = \frac{1}{\csc \theta} \quad \csc \theta = \frac{1}{\sin \theta}$$

$$\cos \theta = \frac{1}{\sec \theta} \quad \sec \theta = \frac{1}{\cos \theta}$$

$$\tan \theta = \frac{1}{\cot \theta} \quad \cot \theta = \frac{1}{\tan \theta}$$

Pythagorean Identities

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$\tan^2 \theta + 1 = \sec^2 \theta$$

$$1 + \cot^2 \theta = \csc^2 \theta$$

Even/Odd Formulas

$$\sin(-\theta) = -\sin \theta \quad \csc(-\theta) = -\csc \theta$$

$$\cos(-\theta) = \cos \theta \quad \sec(-\theta) = \sec \theta$$

$$\tan(-\theta) = -\tan \theta \quad \cot(-\theta) = -\cot \theta$$

Periodic Formulas

For $n \in \mathbb{Z}$

$$\sin(\theta + 2\pi n) = \sin \theta \quad \csc(\theta + 2\pi n) = \csc \theta$$

$$\cos(\theta + 2\pi n) = \cos \theta \quad \sec(\theta + 2\pi n) = \sec \theta$$

$$\tan(\theta + \pi n) = \tan \theta \quad \cot(\theta + \pi n) = \cot \theta$$

Double Angle Formulas

$$\sin(2\theta) = 2 \sin \theta \cos \theta$$

$$\cos(2\theta) = \cos^2 \theta - \sin^2 \theta$$

$$= 2 \cos^2 \theta - 1$$

$$= 1 - 2 \sin^2 \theta$$

$$\tan(2\theta) = \frac{2 \tan \theta}{1 - \tan^2 \theta}$$

Degrees to Radians Formulas

If x is an angle in degrees and t is an angle in radians then

$$\frac{\pi}{180} = \frac{t}{x} \implies t = \frac{\pi x}{180}, \text{ and } x = \frac{180t}{\pi}$$

Half Angle Formulas

$$\sin \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{2}}$$

$$\cos \frac{\theta}{2} = \pm \sqrt{\frac{1 + \cos \theta}{2}}$$

$$\tan \frac{\theta}{2} = \pm \sqrt{\frac{1 - \cos \theta}{1 + \cos \theta}}$$

$$\sin^2 \theta = \frac{1}{2}(1 - \cos(2\theta))$$

$$\cos^2 \theta = \frac{1}{2}(1 + \cos(2\theta))$$

$$\tan^2 \theta = \frac{1 - \cos(2\theta)}{1 + \cos(2\theta)}$$

Sum and Difference Formulas

$$\sin(\alpha \pm \beta) = \sin \alpha \cos \beta \pm \cos \alpha \sin \beta$$

$$\cos(\alpha \pm \beta) = \cos \alpha \cos \beta \mp \sin \alpha \sin \beta$$

$$\tan(\alpha \pm \beta) = \frac{\tan \alpha \pm \tan \beta}{1 \mp \tan \alpha \tan \beta}$$

Product to Sum Formulas

$$\sin \alpha \sin \beta = \frac{1}{2}[\cos(\alpha - \beta) - \cos(\alpha + \beta)]$$

$$\cos \alpha \cos \beta = \frac{1}{2}[\cos(\alpha - \beta) + \cos(\alpha + \beta)]$$

$$\sin \alpha \cos \beta = \frac{1}{2}[\sin(\alpha + \beta) + \sin(\alpha - \beta)]$$

$$\cos \alpha \sin \beta = \frac{1}{2}[\sin(\alpha + \beta) - \sin(\alpha - \beta)]$$

Sum to Product Formulas

$$\sin \alpha + \sin \beta = 2 \sin \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right)$$

$$\sin \alpha - \sin \beta = 2 \cos \left(\frac{\alpha + \beta}{2} \right) \sin \left(\frac{\alpha - \beta}{2} \right)$$

$$\cos \alpha + \cos \beta = 2 \cos \left(\frac{\alpha + \beta}{2} \right) \cos \left(\frac{\alpha - \beta}{2} \right)$$

$$\cos \alpha - \cos \beta = -2 \sin \left(\frac{\alpha + \beta}{2} \right) \sin \left(\frac{\alpha - \beta}{2} \right)$$

Cofunction Formulas

$$\sin \left(\frac{\pi}{2} - \theta \right) = \cos \theta \quad \csc \left(\frac{\pi}{2} - \theta \right) = \sec \theta$$

$$\cos \left(\frac{\pi}{2} - \theta \right) = \sin \theta \quad \sec \left(\frac{\pi}{2} - \theta \right) = \csc \theta$$

$$\tan \left(\frac{\pi}{2} - \theta \right) = \cot \theta \quad \cot \left(\frac{\pi}{2} - \theta \right) = \tan \theta$$

IV. INVERSE TRIG FUNCTIONS

Definition

$$y = \sin^{-1} x \implies x = \sin y$$

$$y = \cos^{-1} x \implies x = \cos y$$

$$y = \tan^{-1} x \implies x = \tan y$$

Domain and Range

Function	Domain	Range
$y = \sin^{-1} x$	$-1 \leq x \leq 1$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$
$y = \cos^{-1} x$	$-1 \leq x \leq 1$	$0 \leq y \leq \pi$
$y = \tan^{-1} x$	$-\infty \leq x \leq \infty$	$-\frac{\pi}{2} \leq y \leq \frac{\pi}{2}$

Inverse Properties

$$\begin{aligned}\sin(\sin^{-1}(x)) &= x & \sin^{-1}(\sin(\theta)) &= \theta \\ \cos(\cos^{-1}(x)) &= x & \cos^{-1}(\cos(\theta)) &= \theta \\ \tan(\tan^{-1}(x)) &= x & \tan^{-1}(\tan(\theta)) &= \theta\end{aligned}$$


Alternate Notation

$$\begin{aligned}\sin^{-1} x &= \arcsin x \\ \cos^{-1} x &= \arccos x \\ \tan^{-1} x &= \arctan x\end{aligned}$$

V. LAW OF SINES, COSINES AND TANGENTS

Law of Sines

$$\frac{\sin \alpha}{a} = \frac{\sin \beta}{b} = \frac{\sin \gamma}{c}$$



Law of Cosines

$$\begin{aligned}a^2 &= b^2 + c^2 - 2bc \cos \alpha \\ b^2 &= a^2 + c^2 - 2ac \cos \beta \\ c^2 &= a^2 + b^2 - 2ab \cos \gamma\end{aligned}$$

Mollweides Formula

$$\frac{a+b}{c} = \frac{\cos \frac{1}{2}(\alpha - \beta)}{\sin \frac{1}{2}\gamma}$$

Law of Tangents

$$\begin{aligned}\frac{a-b}{a+b} &= \frac{\tan \frac{1}{2}(\alpha - \beta)}{\tan \frac{1}{2}(\alpha + \beta)} \\ \frac{b-c}{b+c} &= \frac{\tan \frac{1}{2}(\beta - \gamma)}{\tan \frac{1}{2}(\beta + \gamma)} \\ \frac{a-c}{a+c} &= \frac{\tan \frac{1}{2}(\alpha - \gamma)}{\tan \frac{1}{2}(\alpha + \gamma)}\end{aligned}$$

VI. UNIT CIRCLE

For any ordered pair on the unit circle (x, y) : $\cos \theta = x$ and $\sin \theta = y$. Example: $\cos \left(\frac{5\pi}{3} \right) = \frac{1}{2}$, and $\sin \left(\frac{5\pi}{3} \right) = -\frac{\sqrt{3}}{2}$.

