

WESTERN FRONT

Assessment Question: How has medicine changed from c1250 to the modern day?

Name: _____

Class: _____

Progress & Assessment Tracker	Do Now	RAG	Effort	Level	WWW (What Went Well)	EBI (Even Better If)
KT5.1: What was the historical and medical context of the Western Front?	/ 6	○				
KT5.2: How did the trench environment create new medical challenges?	/ 6	○				
KT5.3: What new illnesses and wounds did soldiers face?	/ 6	○				
KT5.4: How did the RAMC and FANY operate the chain of evacuation?	/ 6	○				
KT5.5: What incredible medical advances were forged on the Western Front?	/ 6	○				
Final Unit Grade:						



Section A: The Historic Environment

This section assesses your knowledge of the historic environment of the Western Front and counts for 10% of your total GCSE.

Q1(a) & 1(b): Describe One Feature...

4
Marks

5
mins

5:00

TARGET OBJECTIVE

AO1 (Demonstrate knowledge and understanding of key features)

THE 2-MARK TRIAGE FORMULA

[Sentence 1: State feature clearly]

[Sentence 2: Add specific context]

Examiner Red Flags

The Single-Sentence List:

Naming a feature without elaboration (e.g., "One feature

Grade 9 Checklist

Structural Separation: Have I written exactly two distinct

was gas attacks"). This only secures 1 mark.

Vague Generalizations:

Broad statements without precise historical vocabulary.

"Too Much Detail": Writing a whole paragraph. Examiners award the 2 marks as soon as the feature and one detail are met. Excess writing wastes precious time.

sentences for 1(a) and two for 1(b)?

First-Sentence Punch: Does my first sentence explicitly state one physical, technological, or administrative feature?

Precise Contextual Anchor: Does my second sentence deploy named, specific historical data?


Zero Overlap: Are my answers for 1(a) and 1(b) completely distinct?

Question 2(a): Source Utility

8 Marks

12-15 mins

15:00



TARGET OBJECTIVE

AO3 (Analyse, evaluate, and use sources to make substantiated judgements)

THE GRADE 9 UTILITY STRUCTURE

1 Enquiry-Focused Thesis

2 Source Paragraph (Content, Provenance, Context)

3 Explicit Utility Verdict (Do NOT compare)

Examiner Red Flags

Generic Provenance

Mnemonics: Using rote-learned checklists to make sweepingly dismissive statements (e.g., "Source A is a diary so it is biased/unreliable").

The Comparison Trap:

Wasting time comparing Source A and Source B. There are zero marks available for comparing.

Simple Comprehension:

Simply listing what the source "shows" or "says" without drawing historical inferences.

Grade 9 Checklist

Enquiry Alignment: Does the first sentence of each paragraph state how useful that specific source is for the precise enquiry?

Double-Source Balance:

Have I given equal analytical weight to both sources?

Inference from Content:

Have I pulled a specific quote or visual detail and explicitly explained what it reveals?

Provenance Deconstruction:

Have I evaluated why the source was created, who created it, and when?

Contextual Verification:


Have I integrated independent historical knowledge?

Question 2(b): Source Follow-Up

4 Marks

5 mins

5:00



TARGET OBJECTIVE

A03 (Formulate historical questions and plan a historical enquiry)

THE ENQUIRIES CONNECTION MAP

1. Precise Source Detail

2. Broadening Question

3. Contemporary Source

4. How it Helps

Examiner Red Flags

The "Unlinked" Chain:

Proposing a follow-up question that has no logical connection to the physical quote selected in Box 1.

Anachronistic Sources:

Suggesting "interviews with soldiers," "the internet," or "textbooks." You must select a primary source.

Circular Explanations:

Writing "This would help answer my question because it would tell me what I want to know" in Box 4 receives 0 marks.

Grade 9 Checklist

Box 1 (Detail): Have I copied a single, direct, and highly specific quote?

Box 2 (Question): Is my question tightly focused on the detail in Box 1?

Box 3 (Source Type): Have I suggested a highly specific, contemporary primary source? (e.g. RAMC medical diaries)

Box 4 (How it Helps): Have I explained exactly what information my suggested source would contain?

L1: KT5.1: What was the historical and medical context of the Western Front?

LEARNING OBJECTIVES

- ☐ Explain the state of surgical asepsis, radiology, and transfusion science on the eve of World War I.

In the decades leading up to the First World War, medicine was experiencing a period of rapid and exciting transformation. The devastating surgical problems of pain and infection were finally being conquered, allowing doctors to carry out more complex operations than ever before. However, while incredible breakthroughs like aseptic surgery, x-rays, and blood typing had laid a strong foundation for modern medicine, there were still major limitations—particularly regarding blood storage and the mobility of equipment—that would pose severe challenges when the war broke out in 1914.

Did you know?

- 1895: The year **Wilhelm Roentgen** discovered x-rays, allowing doctors to diagnose internal injuries without surgery.
- 1901: The year **Karl Landsteiner** discovered different blood groups, which successfully stopped patients from rejecting transfused blood.
- Aseptic Surgery: The practice of ensuring no germs enter the operating theatre in the first place, achieved through steam-sterilisation (autoclaves), rubber gloves, and surgical gowns.

Do Now Activity

Score: / 6

1. What lifestyle choice was definitively linked to lung cancer in the 1950s?

2. Name one modern technology used to diagnose lung cancer.

3. How does the government try to prevent people from smoking?

4. Explain why lung cancer is so difficult to treat effectively.

5. Compare the government's response to lung cancer with its response to cholera in the 19th century.

6. Recall: Who was Edward Jenner and what was their key contribution to medicine?

Vocabulary Check

Fill in the blanks using the vocabulary words below.

Aseptic surgery | Mobile X-ray Unit | Blood Depot

Before the outbreak of the First World War, major breakthroughs in Britain laid the foundation for wartime medical advances. In operating theatres, surgeons transitioned to _____ to prevent microbes from entering wounds during procedures. Meanwhile, the discovery of radiation allowed for the use of _____ designs to find deeply embedded shrapnel in wounded soldiers closer to the battlefield. Finally, to treat severe blood loss and shock, doctors developed methods of preserving and storing blood in a _____ so that transfusions could be carried out quickly without the donor needing to be present.

Source A : The Western Front



The Western Front

The Context of the Western Front When World War I broke out in 1914, the British Expeditionary Force (BEF) was sent to Northern France. The terrain of the Western Front was primarily flat agricultural land, particularly in the Ypres Salient and the Somme. Constant artillery bombardment destroyed the drainage systems, turning the battlefield into a horrific, muddy swamp. The war quickly devolved into a stalemate, with both sides digging hundreds of miles of trenches. This static war of attrition meant that millions of men lived in extremely unhygienic conditions, exposed to the weather, rats, and human waste. The unique environment of the Western Front created unprecedented medical challenges that the Royal Army Medical Corps (RAMC) was initially completely unprepared for.

The Move Towards Aseptic Surgery By the late 1890s, surgeons had moved past Joseph Lister's antiseptic methods (killing germs already in the wound) and towards aseptic surgery, which focused on preventing germs from getting into the operating theatre in the first place. All medical staff were required to meticulously wash their hands, faces, and arms before entering the theatre. Surgeons began wearing sterilised rubber gloves and surgical gowns, which dramatically decreased the rate of infection. Surgical instruments were steam-sterilised using a machine called an autoclave, invented in 1881 by Charles Chamberland. Some theatres even pumped air over a heating system to sterilise it before it entered the room. By 1900, aseptic surgery was the standard, meaning that complex, intrusive operations could be carried out safely in civilian hospitals without the high risk of fatal postoperative infections.

Q1. Identify the main geographical feature of the Western Front.

The Development of X-Rays X-rays were discovered accidentally in 1895 by the German physicist Wilhelm Roentgen, who noticed that the rays passed through flesh but not through bone. They were used for diagnostic purposes almost immediately; by 1896, radiology departments were opened in British hospitals. For example, Dr John Hall-Edwards at Birmingham General Hospital became one of the first doctors to use an x-ray to locate a needle in a woman's hand before operating. However, early x-rays had severe problems: the radiation doses released were 1,500 times stronger than today, causing patients to suffer burns and hair loss. Furthermore, the glass tubes were incredibly fragile, the machines were far too heavy to be moved easily, and taking a single image could take up to 90 minutes, requiring the patient to stay completely still. While x-rays successfully allowed doctors to diagnose internal problems without cutting the patient open, the machines were heavy, slow, and dangerous, meaning they were not yet suited for the fast-paced, mobile environment of a battlefield.

Q2. Describe the conditions of the trenches during the war of attrition.

Wilhelm Roentgen

Blood Transfusions and Storage Problems James Blundell carried out the first human-to-human blood transfusions in 1818 to save women suffering from blood loss during childbirth. However, early transfusions were highly dangerous because blood types were not understood, often resulting in rejection

and death. In 1901, the Austrian doctor Karl Landsteiner revolutionised the process by discovering three blood groups (A, B, and O), with a fourth (AB) found a year later. In 1907, blood group O was identified as the universal blood group. Despite this, a massive problem remained: blood coagulated (clotted) as soon as it left the body, meaning it could not be stored. Because blood could not be stored in a bank, transfusions had to be carried out with the donor and the patient connected directly to each other by a tube in the exact same room. This made dealing with mass casualties on a battlefield virtually impossible.

KNOWLEDGE CHECK (Q5)

What was the main limitation of performing blood transfusions in hospital settings before the discovery of sodium citrate in 1915?

- ☐ A. Blood types had not yet been discovered, so transfusions were always fatal.
 - ☐ B. Blood could not be stored because it quickly coagulated (clotted), meaning a donor had to be physically present.
 - ☐ C. Syringes had not yet been invented.
 - ☐ D. The Church banned the transfer of blood between two people.
-

Q3. Explain how artillery bombardment worsened the medical situation.

Q4. To what extent was the RAMC prepared for the medical challenges of the Western Front?

Karl Landsteiner

Pair & Share Activity

Prompt: Which pre-war medical development was the most critical for the survival of soldiers on the Western Front: the rise of aseptic surgery, the invention of mobile X-rays, or the progress in storing and transfusing blood?

Think: Jot down your choice and one piece of evidence from your knowledge (such as the high rate of infection from muddy trench soil making aseptic methods vital, or blood transfusions preventing fatal shock before surgery).

Historian's Corner: The Debate over the Source of Medical Innovation in World War I

Historians of military medicine debate whether the rapid medical advances of the First World War represented brand new, revolutionary breakthroughs or merely the accelerated application of existing peacetime technologies. Traditionalist historians argue that the Western Front acted as a vast clinical laboratory, forcing rapid, unprecedented experimentation that yielded genuine breakthroughs in surgery and logistics. However, revisionist historians argue that almost all key technologies—including aseptic surgery, sodium citrate blood preservation, and portable X-ray tubes—had already been discovered in civilian laboratories before 1914. They suggest that the war's real contribution was not scientific creation, but rather the massive state funding and logistical mobilization that allowed these pre-existing civilian ideas to be scaled up and standardized across thousands of casualties.

GCSE Exam Practice

Source A

From the personal pocket diary of Private Thomas Atkins, serving with the British Infantry in the Ypres Salient, November 1916.

"The communication trenches are flooded with three feet of icy water, and the clay walls are collapsing into thick slime. Our boots are permanently waterlogged, and we have to dig out our firesteps constantly. The enemy shelled our support lines yesterday, throwing up colossal geysers of wet mud that choked our drainage channels."

Source B

An official archival photograph taken on the Somme in 1916, showing British infantrymen in a muddy communication trench.



Provenance Scaffolding:

Source A is a private diary from a soldier on the front line; consider how this affects its honesty compared to an official account. Source B is an official photograph, which might be censored, but what does the undeniable presence of thigh-deep mud tell you regardless?

Q6. 1 (a). How useful are Sources A and B for an enquiry into the difficulties faced by soldiers in the trench system of the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

[illegible]

GCSE Exam Practice

Q Describe one feature of the First Battle of Ypres. (2 marks)

Q Describe one feature of the Battle of the Somme. (2 marks)

Exam Practice

1. Describe one feature of the support trench system on the Western Front. (2 marks)

2. Describe one feature of the problems involved in transporting wounded soldiers away from the battleground. (2 marks)

3. How useful are Sources A and B for an enquiry into the difficulties faced by soldiers in the trench system of the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

Source A

Source A: From the personal pocket diary of Private Thomas Atkins, serving with the British Infantry in the Ypres Salient, November 1916. 'The communication trenches are flooded with three feet of icy water, and the clay walls are collapsing into thick slime. Our boots are permanently waterlogged, and we have to dig out our firesteps constantly. The enemy shelled our support lines yesterday, throwing up colossal geysers of wet mud that choked our drainage channels.'

Source B

Source B: An official photograph taken on the Somme in October 1916. It shows British soldiers attempting to maintain a communication trench.



4. How could you follow up Source A to find out more about the work of the stretcher bearers on the Western Front? (4 marks)

Source A

Source A: A diary entry from an RAMC officer at the Somme, 1916: 'The terrain is devastated. We cannot use the motor ambulances as the heavy rains and artillery have turned the roads into deep quagmires of mud. We are forced to use six horses to pull a single wagon.'

Documentary / General Notes

Title / Topic: _____

☐ Emerging (1-2) ☐ Emerging+ (3) ☐ Expected (4-5) ☐ Expected+ (6-7) ☐ Greater Depth (8-9)

Teacher Comment:

L2: KT5.2: How did the trench environment create new medical challenges?

LEARNING OBJECTIVES

- ☐ Explain the causes, symptoms, and preventive strategies developed to combat trench foot and trench fever.
- ☐ Analyze the impact of weaponry on injuries, specifically focusing on shrapnel, head injuries, and gas gangrene from soil.

The First World War on the Western Front forced the British army into a static, defensive conflict across Flanders and northern France. Instead of a fast-moving war of movement, soldiers lived and fought in a complex network of muddy trenches. This brutal environment not only produced staggering casualties in major battles like the Somme and Ypres, but the devastated terrain and narrow trenches created immense problems for the medical services trying to transport and treat the wounded.

Did you know?

- Ypres Salient: A vulnerable 'bulge' in the Allied trench line in Belgium that was surrounded by the enemy on three sides.
- Zig-zag: The pattern in which trenches were deliberately dug to prevent enemy fire and explosive blasts from travelling straight down the line.
- 2.5 miles: The length of the underground tunnel network dug into the chalk at Arras to safely shelter troops and a hospital in 1916.

Do Now Activity

Score: / 6

1. When did the First World War begin and end?

2. What was the 'Western Front'?

3. Name the three major battles fought by the British on the Western Front.

4. Explain the impact of the terrain at Ypres on the soldiers.

5. Why was the casualty rate so high on the Western Front?

6. Recall: Who was Joseph Lister and what was their key contribution to medicine?


Vocabulary Check

Write a historically accurate sentence connecting two terms from the glossary box below.

Trench Foot | Trench Fever | Shell Shock

Your Sentence:

Source A : Map of the Somme Offensive



Map of the Somme Offensive

Trench Environment and Illnesses The trenches were breeding grounds for disease. The most notorious environmental illness was Trench Foot, caused by standing in cold, waterlogged mud for days without changing socks. It caused feet to swell, blister, and turn gangrenous, often requiring amputation. The army eventually ordered soldiers to change socks twice a day and rub whale oil on their feet. Another major issue was Trench Fever, which caused severe joint pain and high fevers. It took years for doctors to realize it was transmitted by the millions of body lice living in the soldiers' unwashed uniforms. To combat this, the army set up massive delousing stations and bathhouses to disinfect clothing with hot steam.

The Theatre of War (Key Battles) The Ypres Salient: A vulnerable 'bulge' in the Allied line in Belgium, surrounded by the enemy on three sides, allowing the Germans to fire down from the higher ground. The First Battle of Ypres (Autumn 1914) was critical; the British held onto the salient to stop the Germans advancing to the sea, successfully retaining control of the vital English Channel ports. Major battles later included the Second Battle of Ypres (1915), which saw the first use of chlorine gas by the Germans, and the Third Battle of Ypres (Passchendaele, 1917), where heavy rain turned the ground into a deadly, waterlogged sea of mud. During the fight for Hill 60 (April 1915), the British successfully used offensive mining to tunnel under and blow up the German positions. The Somme (1916): A massive British attack

aiming to relieve pressure on the French at Verdun. It became a medical nightmare, resulting in nearly 60,000 British casualties on the very first day and over 400,000 by the end of the campaign. Arras (1917): Because the ground was chalky and easy to tunnel through, British and New Zealand miners dug a vast 2.5-mile network of tunnels. This safely sheltered 25,000 troops and included a fully functioning underground hospital.

Q1. Identify the cause of Trench Foot.

The Trench System and its Organisation The frontline trench was the firing line where attacks were launched from. Soldiers stood on a firestep to shoot over the sandbag parapet. The support trench was located about 80 metres behind the frontline; troops would retreat here if the frontline came under heavy attack. The reserve trench was at least 100 metres further back, holding backup troops ready to be mobilised for a counter-attack. Communication trenches linked these rows together, allowing the safe movement of men, equipment, and wounded soldiers between the front and the rear. To prevent enemy fire or the blast wave from an exploding shell from travelling straight down the line, trenches were deliberately dug in a zig-zag pattern. The complex, narrow, and zig-zag design of the trench system made it incredibly difficult and physically exhausting for stretcher-bearers to safely manoeuvre seriously wounded men around the tight corners while under fire.

Q2. Describe the symptoms of Trench Fever.

Significance of the Terrain and Transport Problems Constant artillery shelling destroyed the region's roads and natural drainage systems. This left the landscape heavily cratered and filled with deep, waterlogged mud. Heavy motor ambulances frequently got permanently stuck in the deep mud, meaning the army had to rely heavily on horse-drawn ambulance wagons, which often required six horses to pull them through the sludge. These wagons were violently bumpy and often worsened severe injuries. Stretcher-bearers (working in teams of four to six) faced the exhausting, dangerous task of carrying the wounded through muddy shell-holes in No Man's Land, often under heavy fire or in the dark. Once inside the trenches, it was incredibly difficult to manoeuvre stretchers due to crowded troops moving in opposite directions. The devastated terrain and narrow trench system severely slowed down the evacuation of wounded soldiers; this delay caused men to die from blood loss or shock, and allowed deep infections like gas gangrene to take hold before they could reach a Casualty Clearing Station.

KNOWLEDGE CHECK (Q5)

How did the British Army attempt to prevent soldiers from developing trench foot in wet and muddy conditions?

- ☐ A. By providing them with waterproof rubber wellington boots.

- ☐ B. By issuing orders for soldiers to rub their feet with whale oil and regularly change their socks.
- ☐ C. By administering a trench foot vaccine to all soldiers.
- ☐ D. By heating the trenches with underground pipes.

Q3. Explain how the army attempted to prevent Trench Fever.

Q4. Evaluate the effectiveness of military discipline in preventing environmental illnesses.

Pair & Share Activity

Prompt: Which aspect of the trench environment presented the most difficult challenge for the Royal Army Medical Corps (RAMC): the physical environment (producing trench foot), the biological vectors (producing trench fever), or the psychological strain of industrial warfare (producing shell shock)?

Think: Jot down your choice and one piece of evidence from the sources, such as the thousands of men evacuated for trench foot, the difficulty of delousing crowded trenches to stop lice, or the fact that shell shock was a completely new condition requiring specialists.

Your Notes:

Historian's Corner: The Debate over the Treatment and Diagnosis of Shell Shock (NYD.N)

Historians of military medicine actively debate the British Army's evolving response to shell shock (NYD.N) during the First World War. Traditionalist histories often focused on the harshness of the military command, highlighting that early in the war, soldiers showing symptoms of psychological trauma were frequently accused of cowardice or malingering, with some even facing court-martial and execution. However, revisionist

historians argue that the British medical services adapted remarkably quickly to an unprecedented crisis. They point out that by 1916, the RAMC recognized shell shock as a genuine medical condition, establishing specialist treatment centers close to the front lines. They argue that the policy of treating soldiers quickly and locally (focusing on rest and reassurance) was a highly advanced clinical approach that aimed to prevent chronic mental illness, even though the primary military motive remained returning men to active fighting.

GCSE Exam Practice

Source A

From the personal memories of William Easton, an 18-year-old volunteer with the East Anglian Field Ambulance, describing the Passchendaele/Ypres campaign in 1917.

"During the third offensive at Ypres, we navigated paths choked with thick, hip-deep mire. In the devastated sector of Hooze, we struggled to relocate casualties. The physical exertion of a single carry left us utterly spent; delivering just two or three wounded men to safety daily was our absolute limit. We worked in teams of four, using shoulder straps to stabilize the stretcher and prevent agonising falls."

Source B

An official archival photograph taken on the Somme in 1916, showing a horse-drawn ambulance navigating a deeply rutted, waterlogged dirt road.



Provenance Scaffolding:

Think about who wrote or produced this source. For Source A, it's an East Anglian Field Ambulance worker at Ypres. Does this personal, on-the-ground experience make it more reliable for understanding the physical toll? For Source B, it's an official photograph from the Somme. Was it meant to document reality, or perhaps carefully framed for morale?

Q6. 2 (a). How useful are Sources A and B for an enquiry into the impact of the terrain on the transport of the wounded on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

GCSE Exam Practice

3. How useful are Sources A and B for an enquiry into the problem of ill health arising from the trench environment? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

Source A

Source A: A report from a Medical Officer in 1915: 'Altogether about 200 men had to be evacuated from our section of the trenches with swollen, numb feet due to the freezing mud. Several require amputation.'

Source B

Source B: A diary entry from an RAMC surgeon: 'The men are brought in with horrific shrapnel injuries. The metal drives the filthy, manured soil deep into the muscle, creating an environment where gas gangrene thrives. We often have to amputate to save their lives.'

4. How could you follow up Source A to find out more about the problem of trench foot? (4 marks)

Source A

Source A: A report from a Medical Officer in 1915: 'Altogether about 200 men had to be evacuated from our section of the trenches with swollen, numb feet due to the freezing mud. Several require amputation.'

Documentary / General Notes

Title / Topic: _____

☐ Emerging (1-2) ☐ Emerging+ (3) ☐ Expected (4-5) ☐ Expected+ (6-7) ☐ Greater Depth (8-9)

Teacher Comment:

L3: KT5.3: What new illnesses and wounds did soldiers face?

LEARNING OBJECTIVES

- ☐ Explain the connection between artillery weaponry, shrapnel, and the introduction of the steel Brodie helmet in 1915.
- ☐ Analyse how the muddy, manured environment of Flanders led to severe wound infections like gas gangrene and tetanus.

Living and fighting in the waterlogged trenches of the Western Front exposed soldiers to horrifying new medical dangers. Beyond the immediate, devastating trauma of being hit by shrapnel or rapid-fire machine guns, men faced the constant, invisible threat of deadly soil-borne infections like gas gangrene. Furthermore, the brutal, freezing environment itself caused severe illnesses like trench foot and shellshock, while the introduction of poison gas added a terrifying new psychological and physical dimension to casualty care.

Did you know?

- 58%: The percentage of all wounds on the Western Front caused by high-explosive shells and shrapnel.
- 1915: The year the steel Brodie helmet was introduced, drastically reducing fatal head injuries by 80%.
- 80,000: The estimated number of British troops who suffered from the severe psychological trauma of shellshock.

Do Now Activity

Score: / 6

1. What was the purpose of the duckboards in the trenches?

2. Name the pattern in which trenches were dug.

3. What was 'No Man's Land'?

4. Explain why the zig-zag pattern of trenches was crucial for survival.

5. To what extent did the trench environment itself cause casualties?

6. Recall: Who was Francis Crick and what was their key contribution to medicine?

Vocabulary Check

Write a clear definition and a historically accurate sentence for the term: **Gas Gangrene**

Source A : The devastated, muddy terrain of Passchendaele



The devastated, muddy terrain of Passchendaele

Wounds and Infections The weapons of WWI caused devastating injuries. High-explosive artillery shells caused over 60% of all wounds. When these shells exploded, hot jagged shrapnel tore through flesh, dragging the muddy, bacteria-filled uniform deep into the wound. This immediately caused deadly infections like Gas Gangrene, which could kill a man within 24 hours. Soldiers also faced the horrors of poison gas. Chlorine, Phosgene, and Mustard gas caused blinding, severe lung damage, and horrific internal and external blisters. While gas only caused about 6,000 British deaths, the psychological terror it inflicted was immense, requiring soldiers to carry heavy, uncomfortable gas masks at all times.

Illnesses from the Trench Environment Trench foot was a painful swelling caused by standing in cold mud and water for hours. It restricted blood flow, eventually causing the flesh to rot, a condition known as gangrene. Preventative orders included rubbing whale oil into the feet and changing socks regularly, but severe cases usually required amputation. Trench fever was a debilitating condition causing severe flu-like symptoms, high temperatures, and aching muscles. It was incredibly widespread, affecting an estimated half a million men. In 1918, doctors finally proved it was spread by lice living in the seams of clothing, leading to the creation of special delousing stations. Shellshock was a poorly understood psychological condition caused by the constant trauma of bombardment and the noise of the big guns.

Symptoms included nightmares, loss of speech, uncontrollable shaking, and complete mental breakdown. It affected an estimated 80,000 British troops, with some being treated at the specialist Craiglockhart Hospital in Edinburgh. The army officially recorded the condition under the code NYD.N (Not Yet Diagnosed, Nervous). (Key Detail: You must use the exact military acronym 'NYD.N.' to show precise historical knowledge of how the army categorized shellshock). Because the brutal trench environment itself caused massive casualties, the medical corps had to dedicate huge resources to preventative measures just to keep the army fit enough to fight.

Q1. Identify the weapon that caused over 60% of all wounds on the Western Front.

The Devastation of Shrapnel and Explosives High-explosive shells and shrapnel (jagged metal fragments packed inside artillery shells) were the greatest killers, responsible for 58% of all wounds. They scattered fast-moving metal over a massive area, causing horrific trauma. Machine guns and rifles accounted for roughly 39% of wounds. A machine gun could fire between 450 and 600 rounds a minute, instantly fracturing bones and piercing major internal organs. The sheer destructive power of modern industrial weapons meant soldiers suffered complex, jagged injuries and massive blood loss that traditional military surgeons had never encountered on this scale before.

Q2. Describe the physical and psychological effects of poison gas.

The Deadly Threat of Infection The farmland of France and Belgium had been heavily fertilised with manure before the war, meaning the soil was packed with lethal bacteria that caused tetanus and gas gangrene. When shrapnel or a bullet pierced a soldier, it dragged dirty uniform fabric and infected mud deep into the wound. While the danger of tetanus was successfully reduced by administering anti-tetanus injections from late 1914, there was no initial cure for gas gangrene. This infection rapidly produced gas inside the dying tissue, turning flesh black and emitting a foul smell, and could kill a man within a day. Because lethal bacteria were instantly driven deep into the body at the moment of injury, traditional surface antiseptics entirely failed, forcing doctors to find radically new surgical methods to stop the infection.

KNOWLEDGE CHECK (Q4)

Why were deep shrapnel wounds on the Western Front almost guaranteed to develop the deadly infection known as gas gangrene?

☐ A. Because the German army dipped their bullets in poison before firing them.

- ☐ B. Because the French farmland had been heavily fertilized with manure for centuries, filling the soil with deadly tetanus and gangrene bacteria.
- ☐ C. Because the British soldiers refused to wash their uniforms.
- ☐ D. Because chlorine gas from chemical attacks infected the wounds.

Q3. Explain why shrapnel wounds almost always became infected.

Head Injuries and the Brodie Helmet At the start of the war, British soldiers only wore soft cloth caps. Because men stood in trenches, their heads were highly exposed, and approximately 20% of all wounds were to the head, face, and neck. In 1915, the British army introduced the steel Brodie helmet, which included a strong strap to prevent it from being blown off in an explosion. The widespread introduction of the Brodie helmet was a monumental preventative success, rapidly reducing the number of fatal head wounds caused by shrapnel by an estimated 80%.

Q5. To what extent could doctors successfully treat deep infections during WWI?

The Terror of Poison Gas Chlorine gas was first used by the Germans at the Second Battle of Ypres (April 1915), causing victims to slowly suffocate as their lungs filled with fluid. Before gas masks were officially issued in July 1915, soldiers urinated on cotton pads and pressed them to their faces to neutralise the chemical. Phosgene was introduced later in 1915; it was faster-acting and more deadly, capable of killing an exposed person within two days. Mustard gas (1917) was an odourless gas that caused severe internal and external blistering and could burn the skin straight through clothing. Although poison gas was a terrifying psychological weapon, the rapid development of effective gas masks meant it was actually the cause of under 5% of British deaths (around 6,000 soldiers), making it far less lethal than artillery fire.

Pair & Share Activity

Prompt: Which posed a more difficult medical challenge for the Royal Army Medical Corps (RAMC) on the Western Front: the physical injuries caused by industrial weaponry (such as shrapnel wounds and head trauma) or the environmental infections caused by the Flanders terrain (such as gas gangrene, tetanus, and trench foot)?

Think: Jot down your choice and one piece of evidence from the sources, such as the introduction of the Brodie helmet reducing head wounds vs. the manured soil causing gas gangrene infections that standard antiseptics couldn't treat.

Your Notes:

Historian's Corner: The Debate over the Diagnosis and Treatment of Shell Shock (NYD.N)

Historians of military medicine are deeply divided over the British Army's response to psychological trauma, which was medically recorded as shell shock or NYD.N ('Not Yet Diagnosed, Nervous'). Traditionalist historians argue that the military high command was unsympathetic and hostile, often viewing traumatized soldiers as cowards or malingerers, which in some instances led to court-martials and executions for desertion. Conversely, revisionist historians argue that the RAMC adapted with remarkable speed to an entirely new clinical phenomenon. They highlight that by 1916, the army had established specialized psychiatric units close to the front line, utilizing the advanced concept of 'proximity, immediacy, and expectation' (treating men locally so they expected to return to their units), which laid the groundwork for modern trauma therapy.

GCSE Exam Practice

Source A

From a wartime notebook of Lance Sergeant Elmer Cotton in 1915, providing a private, firsthand account of a gas attack.

"A chlorine discharge floods the victim's respiratory tract, effectively suffocating them while standing on dry ground. The resulting symptoms are excruciating: a blinding headache, an overwhelming craving for water—though drinking it brings immediate fatality—and a severe, stabbing sensation in the lungs. Patients continuously expectorate a distinctive greenish fluid. It represents a truly horrific way to perish."

Source B

An official photograph capturing blinded soldiers forming a chain outside an Advanced Dressing Station.



Provenance Scaffolding:

For Source A, consider that it is from a wartime notebook of Lance Sergeant Elmer Cotton in 1915. This is a private, unedited firsthand account. How does this affect its usefulness? For Source B, an official photograph showing blinded soldiers forming a chain. Photographs capture reality but can also be staged. Does the clear visual evidence of the gas's impact make it highly useful regardless?

Q6. 2 (a). How useful are Sources A and B for an enquiry into the effects of poison gas attacks on soldiers on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

[illegible]

GCSE Exam Practice

Q Describe one feature of the effects of poison gas. (2 marks)

Q Describe one feature of shrapnel wounds. (2 marks)

Exam Practice

1. Describe one feature of aseptic surgery in the early 20th century. (2 marks)

2. Describe one feature of early x-ray machines. (2 marks)

3. How could you follow up Source A to find out more about the treatment and prevention of the effects of gas attacks? (4 marks)

Source A

Source A: From an official report written by a medical officer at an Advanced Dressing Station near Ypres, May 1915: 'At 4:00 AM, the enemy released a dense cloud of yellowish-green gas toward our trenches. Many of the men did not adjust their woolen flannel pads quickly enough. Over fifty casualties were brought into our Dressing Station, their eyes swollen shut and blinded by the vapor. We washed their eyes with bicarbonate solution, but several have already suffocated.'

Documentary / General Notes

Title / Topic: _____

☐ Emerging (1-2) ☐ Emerging+ (3) ☐ Expected (4-5) ☐ Expected+ (6-7) ☐ Greater Depth (8-9)

Teacher Comment:

L4: KT5.4: How did the RAMC and FANY operate the chain of evacuation?

LEARNING OBJECTIVES

- ☐ Explain the four main stages of the Western Front chain of evacuation and the mode of transport between them.
- ☐ Evaluate the roles, significance, and contributions of the RAMC and FANY on the Western Front.

The devastating environment and weaponry of the Western Front required a highly organised system to rescue and treat casualties. The Royal Army Medical Corps (RAMC) and nursing volunteers like the FANY worked tirelessly to transport men through a multi-stage 'chain of evacuation'. From exhausted stretcher-bearers navigating deep mud to the fully equipped underground hospital at Arras, the medical services continually adapted to save lives under unimaginable pressure.

Did you know?

- **Triage:** The vital system used at the Casualty Clearing Station to divide wounded soldiers into three categories to prioritise medical treatment.
- **512:** The number of motor ambulances purchased through The Times newspaper's public appeal in 1914.
- **700:** The number of stretcher beds available inside the fully electrified underground hospital at Arras.

Do Now Activity

Score: / 6

1. What is Trench Foot?

2. Which weapons caused the most wounds on the Western Front?

3. Why were wounds highly prone to infection on the Western Front?

4. Explain how gas attacks affected the soldiers.

5. How effective were the preventative measures against gas attacks?

6. Recall: Who was Florence Nightingale and what was their key contribution to medicine?

Vocabulary Check

Fill in the blanks using the vocabulary words below.

Chain of Evacuation Casualty Clearing Station (CCS) First Aid Nursing Yeomanry (FANY)
--

To save soldiers wounded in the trenches, they had to be moved quickly along the _____ before soil-borne bacteria caused fatal infections. While basic first aid was given close to the battlefield, soldiers with life-threatening wounds were sent further back to a _____ for urgent surgery. To transport these casualties safely, the army relied on volunteers from the _____, who braved shellfire to drive motorized ambulances between frontline dressing stations and the rear hospitals.

The Chain of Evacuation To deal with the massive casualties, the RAMC designed a highly organized 'Chain of Evacuation'. Stretcher bearers would rescue men from No Man's Land and take them to the Regimental Aid Post (RAP) just behind the frontline trench, which provided basic first aid. From there, horse-drawn or motorized ambulances transported the wounded to Advanced Dressing Stations (ADS). Those who needed surgery were sent further back to Casualty Clearing Stations (CCS), which were large tented hospitals located near railway lines. The CCS dealt with the most critical, life-saving surgeries, particularly amputations to stop gangrene. If a soldier survived, they were put on a hospital train to a Base Hospital near the French coast, where specialists worked, before being shipped back to 'Blighty' (Britain).

The Work of the RAMC and Nurses The RAMC was the branch of the army responsible for all medical care, from treating the wounded to maintaining sanitation to keep the men healthy. Due to the sheer scale of casualties, the RAMC expanded massively from 9,000 men in 1914 to 113,000 by 1918. Professional nurses from Queen Alexandra's Imperial Military Nursing Service also grew from 300 to 10,000. The First Aid Nursing Yeomanry (FANY) was a women's voluntary organisation; they famously drove motor ambulances, transported supplies to the frontline, set up cinemas, and even operated a mobile bath unit that could bathe up to 40 men an hour. The rapid expansion of the RAMC and the vital inclusion of female volunteers were absolutely essential to prevent the medical system from entirely collapsing under the unprecedented, staggering number of casualties.

Q1. Identify the first stage of the Chain of Evacuation.

Transport in the Chain of Evacuation Initially, the army relied on horse-drawn ambulance wagons, which were violently bumpy, could only carry a few men, and often worsened severe injuries. A public appeal by The Times newspaper raised money for 512 new motor ambulances, which arrived in October 1914. However, motor vehicles frequently got stuck in the deep mud, meaning horse-drawn wagons (sometimes needing six horses) and exhausted human stretcher-bearers (working in teams of four or six while under enemy fire) remained crucial. To move the wounded to Base Hospitals, specially designed ambulance trains and canal barges were used, with some trains capable of carrying up to 800 casualties at once. The devastated, cratered terrain severely hindered transport; this delay and the bumpy journeys often exacerbated injuries and caused fatal blood loss before soldiers could ever reach life-saving surgery.

Q2. Describe the role of Casualty Clearing Stations (CCS).

Stages of Treatment: RAP and Dressing Stations The Regimental Aid Post (RAP) was located within 200m of the frontline, often in communication trenches or dugouts. A Regimental Medical Officer provided immediate first aid to get walking wounded back to the fight, but could not perform surgery on serious injuries. The wounded were then moved to an Advanced Dressing Station (ADS) or Main Dressing Station (MDS), located about 400m to half a mile back in abandoned buildings, dug-outs, or tents. Here, Field Ambulance units (staffed by medical officers, orderlies, and eventually nurses) provided better bandages and pain relief, keeping men for a maximum of one week. These early stages were vital for preventing frontline trenches from becoming blocked with wounded men, quickly patching up minor injuries, and organising the transport of serious cases further back.

Q3. Explain how patients were transported between the different stages of the evacuation chain.

Stages of Treatment: CCS and Base Hospitals The Casualty Clearing Station (CCS) was located 7 to 12 miles back, safely out of artillery range but near railway lines. Here, patients underwent triage, being sorted into three groups: the walking wounded, those needing hospital treatment, and those with no chance of recovery. Because deadly gas gangrene set in so quickly, the CCS soon took over the role of performing critical, life-saving surgery (like amputations) from the Base Hospitals. Base Hospitals, located near the French and Belgian coastal ports, eventually focused on the longer-term recovery of patients and developed specialist wards for head wounds or gas poisoning. By shifting life-saving surgical operations forward to the CCS, the medical services successfully reduced the time wounded men had to wait for surgery, drastically decreasing deaths from infection.

KNOWLEDGE CHECK (Q5)

What was the main difference between the medical treatments performed at a Regimental Aid Post (RAP) and a Casualty Clearing Station (CCS)?

- ☐ A. The RAP performed complex surgery, while the CCS only applied bandages.
 - ☐ B. The RAP provided immediate first aid near the frontline, while the CCS was equipped with operating theatres and X-ray machines to perform life-saving surgery.
 - ☐ C. The RAP was for officers only, while the CCS was for regular soldiers.
 - ☐ D. The RAP was located in Britain, while the CCS was in France.
-

Q4. Evaluate the significance of the Chain of Evacuation.

The Underground Hospital at Arras In 1916, British and Commonwealth troops linked existing chalk tunnels and quarries beneath the town of Arras. They created a massive underground town that included a fully functioning hospital (nicknamed Thompson's Cave) within 800m of tunnels. It contained 700 spaces for stretchers to be used as beds, operating theatres, rest stations, a mortuary, and was supplied with electricity and piped water. The underground hospital at Arras was highly significant because it functioned as a fully equipped medical facility incredibly close to the frontline, while remaining completely safe from enemy artillery bombardment.

Pair & Share Activity

Prompt: Was the medical service's success on the Western Front driven more by the highly organized military logistics of the RAMC, or by the courage and adaptability of voluntary organizations like the FANY?

Think: Jot down your choice and one piece of evidence from your knowledge (such as the systematic stages of triage from RAP to Base Hospital vs. FANY volunteers driving motorized ambulances under active bombardment).

Your Notes:

Historian's Corner: The Debate over the Primary Role of Casualty Clearing Stations

Historians of military medicine actively debate how the roles of Casualty Clearing Stations (CCSs) and Base Hospitals shifted during the First World War. Traditional accounts argued that Base Hospitals, situated near the safe French coast, remained the primary hubs for surgical intervention throughout the war. However, revisionist historians have shown that the extreme threat of gas gangrene and tetanus from Flanders soil forced a rapid role reversal. Because debridement (wound excision) had to be performed within hours of injury, CCSs located near the frontline railheads became the primary hubs for life-saving surgery, while Base Hospitals were relegated to long-term convalescence. This adaptation demonstrates that wartime medical logistics was not a static plan but a dynamic, rapidly evolving system.

GCSE Exam Practice

Source A

From the 1919 published autobiography of Pat Beauchamp, a volunteer FANY ambulance driver.

"With the Somme offensive raging, dozens of transport cars would form a convoy to meet incoming hospital trains at the railway siding. Once positioned, stretcher bearers loaded our vehicles with four severely injured, non-walking cases. Traversing the ruined, potholed cobblestone roads was an agonizing ordeal. The violent jolts caused immense suffering, yet we were forced to drive slowly to protect those inside."

Source B

A 1916 archival photograph from the Somme showing a motor ambulance navigating a flooded road.



Provenance Scaffolding:

Source A is from the 1919 published autobiography of a FANY ambulance driver. It's a memoir published right after the war. How might writing for a public audience affect her description of the difficulties? Source B is a 1916 photograph from the Somme. Official photographs were subject to censorship. Does its depiction of flooded roads match your own knowledge of the Somme battlefield?

Q6. 2 (a). How useful are Sources A and B for an enquiry into the problems of transporting the wounded on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

[illegible]

GCSE Exam Practice

Q Describe one feature of the FANY. (2 marks)

Q Describe one feature of the Chain of Evacuation. (2 marks)

Exam Practice

1. Describe one feature of the underground hospital at Arras. (2 marks)

2. Describe one feature of the work of the FANY on the Western Front. (2 marks)

3. How useful are Sources A and B for an enquiry into the work of medical staff in the Casualty Clearing Stations (CCS) on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

Source A

Source A: A letter from an RAMC nurse working at a Casualty Clearing Station (CCS) in 1916: 'We worked night after night, in the thunderous noise of raging battles. The surgeons are

Source B

Source B: A diary entry from a member of the FANY in 1917: 'I drove my motor ambulance through the terrible mud for 14 hours straight today, bringing

constantly amputating limbs to stop the spread of gangrene before putting the men on the trains.'

men from the Main Dressing Station down to the railhead.'

4. How could you follow up Source A to find out more about the work of medical staff in the Casualty Clearing Stations? (4 marks)

Source A

Source A: A letter from an RAMC nurse working at a Casualty Clearing Station (CCS) in 1916: 'We worked night after night, in the thunderous noise of raging battles. The surgeons are constantly amputating limbs to stop the spread of gangrene before putting the men on the trains.'

Documentary / General Notes

Title / Topic: _____

☐ Emerging (1-2) ☐ Emerging+ (3) ☐ Expected (4-5) ☐ Expected+ (6-7) ☐ Greater Depth (8-9)

Teacher Comment:

L5: KT5.5: What incredible medical advances were forged on the Western Front?

LEARNING OBJECTIVES

- ☐ Explain how the Thomas splint and mobile X-ray units solved critical evacuation and surgical challenges.
- ☐ Analyze the development of blood transfusion and the significance of the first blood bank at the Battle of Cambrai.

The horrific new weapons of the First World War created devastating, complex injuries that traditional 19th-century surgeons simply did not know how to cure. Faced with thousands of men rapidly dying from gas gangrene, massive blood loss, and shattered limbs, the medical services were forced to turn the Western Front into a vast laboratory. Driven by sheer desperation, they successfully pioneered radical new surgical techniques, mobile diagnostic technology, and miraculous methods for storing blood directly on the battlefield.

Did you know?

- 82%: The dramatically increased survival rate for soldiers with compound leg fractures after the Thomas Splint was introduced in late 1915.
- 1917: The year **Oswald Hope Robertson** established the world's first 'blood depot' (blood bank) before the Battle of Cambrai.
- Debridement: The rapid surgical technique of cutting away dead, damaged, and infected tissue from around a wound to prevent the spread of gas gangrene.

Do Now Activity

Score: / 6

1. What does RAMC stand for?

2. What was the role of the FANY?

3. Describe the 'Chain of Evacuation'.

4. Explain the importance of the Casualty Clearing Stations (CCS).

5. Why was the work of the RAMC so dangerous?

6. Recall: Who was James Simpson and what was their key contribution to medicine?

Vocabulary Check

Write a historically accurate sentence connecting two terms from the glossary box below.

Thomas Splint Carrel-Dakin Method Blood Depot

Your Sentence:

Medical Advances on the Western Front The sheer volume of casualties on the Western Front forced doctors to innovate rapidly. One of the greatest breakthroughs was the Thomas Splint, a metal frame designed by **Hugh Owen Thomas**. Before 1915, 80% of soldiers with broken femurs died from shock or bleeding. The Thomas Splint kept the leg perfectly still during transport, increasing the survival rate to an incredible 82%. Another major advance was in blood transfusions. At the start of the war, transfusions had to be done arm-to-arm. However, in 1915, Richard Lewisohn discovered that adding sodium citrate to blood stopped it from clotting, allowing it to be stored on ice. This led to the creation of the first blood banks at the Battle of Cambrai in 1917, allowing surgeons to treat shock and perform complex surgeries.

Hugh Owen Thomas

New Techniques in the Treatment of Wounds and Infection Because traditional aseptic surgery was impossible in dirty casualty stations and carbolic acid simply did not kill gas gangrene bacteria effectively, radical new solutions had to be found. Debridement (Wound Excision): Surgeons began aggressively cutting away all dead, damaged, and infected tissue from around a wound to physically remove the gangrene. This had to be done incredibly quickly before the wound was stitched up to stop the infection from spreading into healthy flesh. The Carrel-Dakin method: If debridement wasn't enough, surgeons used a system where tubes pumped a chemical sterilised salt solution directly into deep wounds to continually wash out and kill the infection-causing bacteria. The method was effective but highly difficult, as the solution only stayed fresh for six hours and had to be made as it was needed. Amputation: If antiseptics or wound excision failed to stop the spread of infection, the limb was amputated to save the man's life. By 1918, over 240,000 men had lost limbs. Because infection spread so rapidly in the heavily manured soil of the Western Front, surgeons were forced to abandon slow peacetime methods and instead pioneer radical, invasive techniques to physically flush out or cut away deadly bacteria before it killed the patient.

Q1. Identify the device designed by Hugh Owen Thomas.

The Thomas Splint In 1914 and 1915, a soldier with a gunshot or shrapnel wound that fractured their femur (thigh bone) only had a 20% chance of survival. Traditional splints did not keep the leg straight, meaning the jagged bone caused massive internal bleeding and shock during bumpy transport. In December 1915, Robert Jones introduced the Thomas Splint (a device originally designed by his uncle, Hugh Thomas) to the Western Front. This splint rigidly pulled the leg lengthways, completely stopping the broken bones from grinding on each other. The introduction of the Thomas Splint was one of the greatest medical successes of the war. By keeping the leg perfectly still, it prevented fatal blood loss and shock, miraculously increasing the survival rate for thigh fractures from 20% to 82%.

KNOWLEDGE CHECK (Q3)

How did the implementation of the Thomas splint on the Western Front in 1916 directly reduce mortality rates among wounded soldiers?

- ☐ A. It stopped the spread of infection by killing the bacteria that caused gas gangrene.
 - ☐ B. It cured shell shock by holding the soldier firmly in place.
 - ☐ C. It kept fractured leg bones rigidly in place during transport, preventing fatal blood loss and shock.
 - ☐ D. It provided a steady drip of blood for transfusions on the battlefield.
-

Q2. Describe how Richard Lewisohn improved blood transfusions.

The Use of Mobile X-Ray Units X-rays were absolutely essential for accurately locating bullets and deeply embedded shrapnel so they could be removed before infection set in. While static machines were used in Base Hospitals, six mobile x-ray units (vans) were developed to operate in the British sector so they could be driven directly to Casualty Clearing Stations closer to the fighting. There were limitations: the glass tubes overheated very quickly (meaning machines could only be used for an hour before needing to cool down), and they could not detect fragments of dirty clothing driven into wounds. Even though the mobile images were not quite as clear as static machines and they overheated, they were incredibly significant because they allowed life-saving diagnostic technology to be transported directly to the frontline, heavily reducing deaths from infection.

Q4. Explain how the Thomas Splint improved survival rates.

Blood Transfusions and the Blood Bank at Cambrai Severe blood loss caused victims' bodies to go into fatal shock. Dr Lawrence Bruce Robertson pioneered the use of the indirect blood transfusion method (using a syringe and tube) at Boulogne in 1915. However, early transfusions were heavily limited because blood could not be stored and would quickly clot. Scientists rapidly discovered chemical solutions: In 1915, Richard Lewisohn added sodium citrate to stop blood from clotting. Richard Weil then discovered this allowed blood to be refrigerated for up to two days. In 1916, Francis Rous and James Turner added citrate glucose, a monumental breakthrough that allowed blood to be safely stored for up to four weeks. These chemical discoveries allowed the American doctor Oswald Hope Robertson to establish the world's

first 'blood depot' (blood bank) before the Battle of Cambrai in 1917. He stored 22 units of pre-donated universal Type O blood in glass bottles packed with ice and sawdust, successfully treating severely shocked soldiers and proving that stored blood could save lives on the battlefield.

Q5. To what extent did the Western Front accelerate medical progress?

Oswald Hope Robertson

Pair & Share Activity

Prompt: Which medical advance on the Western Front represented a more impressive leap forward: the mechanical simplicity of the Thomas Splint which stabilized fractures, or the complex chemical engineering of blood preservation and storage?

Think: Jot down your choice and one piece of evidence from the sources (e.g., the Thomas splint keeping the leg rigid to prevent internal bleeding during transport, or sodium citrate allowing blood to be stored in depots for the first time without donors present).

Your Notes:

Historian's Corner: The Debate over the Western Front as a Catalyst for Medical Innovation

Military and social historians actively debate the true significance of the First World War in driving medical progress. Traditional histories argue that the Western Front served as a massive, unprecedented laboratory of rapid innovation, where the extreme pressures of trench warfare forced doctors to experiment and make revolutionary breakthroughs in antiseptics, radiology, and blood storage. Conversely, revisionist historians argue that almost all of these 'wartime breakthroughs' were actually based on pre-war civilian discoveries that had already been established in laboratories before 1914, such as the basic design of the Thomas splint, Marie Curie's work with radioactivity, and pre-war transfusion experiments. From this perspective, the war did not spark scientific creation itself, but rather acted as an institutional and financial accelerator, providing the state funding and millions of casualties necessary to scale up and standardize pre-existing civilian technologies.

GCSE Exam Practice


Source A

From an official clinical report written by Captain Arthur Vaughan, a surgeon serving with the Royal Army Medical Corps (RAMC) at a Casualty Clearing Station (CCS) behind the frontline, October 1916.

"Our medical team at the Casualty Clearing Station has fully adopted the new leg traction splint. Before we received this iron frame, almost every compound femur fracture arrived at our ward in a state of terminal shock due to the jagged bones grating inside the flesh during transit. Since implementing this immobilization frame last month, we have kept the fractured limbs completely rigid. Out of forty admissions with thigh wounds, we have lost only six patients."

Source B

A studio photograph taken to train medical orderlies, showing a soldier's leg secured in a Thomas splint.



Provenance Scaffolding:

Source A is a clinical report by a surgeon; does this make its statistics reliable? Source B is a staged training photograph; while not taken on the chaotic frontline, does this actually make it better for demonstrating how the device technically worked?

Q6. 2 (a). How useful are Sources A and B for an enquiry into the effectiveness of new medical developments used to treat wounded soldiers on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

Exam Practice

Source A

'Our medical team at the Casualty Clearing Station has fully adopted the new leg traction splint. Before we received this iron frame, almost every compound femur fracture arrived at our ward in a state of terminal shock due to the jagged bones grating inside

Source B

(Visual Description: A detailed, black-and-white studio photograph showing a soldier lying flat on a wooden stretcher. His right leg is suspended and held rigid inside a metallic Thomas splint, which consists of a padded leather ring pushed up against the groin and parallel steel rods running down both sides of the limb. Bandages

the flesh during transit. Since implementing this immobilization frame last month, we have kept the fractured limbs completely rigid. Out of forty admissions with thigh wounds, we have lost only six patients.'

and canvas straps are wrapped tightly around the thigh and calf to keep the leg completely straight. A canvas sling is hooked to the end of the splint to pull the foot forward, creating traction. Two medical orderlies in clean aprons stand alongside, demonstrating how to attach the secure traction cords.)

4. How could you follow up Source A to find out more about new techniques being used on the Western Front to deal with injuries? (4 marks)

Source A

Source A: A report from a surgeon at a CCS in 1916: 'Several unsuccessful trials this morning to extract a shell fragment from a soldier's brain. Finally, we decided to try using a large wire nail as a magnet.'

Documentary / General Notes

Title / Topic: _____

☐ Emerging (1-2) ☐ Emerging+ (3) ☐ Expected (4-5) ☐ Expected+ (6-7) ☐ Greater Depth (8-9)

Teacher Comment:

Factors Overview: Western Front

Edexcel focuses heavily on the factors that drove medical progress (or held it back). For each factor below, write one specific historical example from this period that either helped or hindered medical progress.

Factor	Specific Historical Example & Impact
Individuals	
The Church & Religion	
Government & Wealth	
Science & Technology	
Attitudes in Society	
War	

Factor	Specific Historical Example & Impact