

WESTERN FRONT

Paper 1: Medicine Through Time with the Western Front



COURSE TEXTBOOK

Enquiry: How has medicine changed from c1250 to the modern day?

Contents

Lesson 1: KT5.1: What was the historical and medical context of the Western Front?

Lesson 2: KT5.2: How did the trench environment create new medical challenges?

Lesson 3: KT5.3: What new illnesses and wounds did soldiers face?

Lesson 4: KT5.4: How did the RAMC and FANY operate the chain of evacuation?

Lesson 5: KT5.5: What incredible medical advances were forged on the Western Front?

L1: KT5.1: What was the historical and medical context of the Western Front?

LEARNING OBJECTIVES

- ☐ Explain the state of surgical asepsis, radiology, and transfusion science on the eve of World War I.

In the decades leading up to the First World War, medicine was experiencing a period of rapid and exciting transformation. The devastating surgical problems of pain and infection were finally being conquered, allowing doctors to carry out more complex operations than ever before. However, while incredible breakthroughs like aseptic surgery, x-rays, and blood typing had laid a strong foundation for modern medicine, there were still major limitations-particularly regarding blood storage and the mobility of equipment-that would pose severe challenges when the war broke out in 1914.

Did you know?

- 1895: The year Wilhelm Roentgen discovered x-rays, allowing doctors to diagnose internal injuries without surgery.
- 1901: The year Karl Landsteiner discovered different blood groups, which successfully stopped patients from rejecting transfused blood.
- Aseptic Surgery: The practice of ensuring no germs enter the operating theatre in the first place, achieved through steam-sterilisation (autoclaves), rubber gloves, and surgical gowns.

Recall & Retrieval

- | | |
|---|--|
| 1. What lifestyle choice was definitively linked to lung cancer in the 1950s? | 2. Name one modern technology used to diagnose lung cancer. |
| 3. State one way the modern government tries to prevent people from smoking. | 4. When did the First World War begin and end? |
| 5. What was the 'Western Front'? | 6. What is a 'magic bullet'? |
| 7. What disease did Edward Jenner vaccinate against in 1796? | 8. What was established in Britain in 1948 to provide free medical care? |
| 9. Who was the Minister of Health responsible for establishing the NHS in 1948? | 10. Which anesthetic was discovered by James Simpson in 1847? |

Vocabulary Check

The Odd One Out: Circle *ONE* term that does not belong with the others. Explain your reasoning: What historical connection links the other terms, and why is your chosen term different?

Salient | Trench System | Dugout | Battle of the Somme (1916) | Battle of Arras (1917) | Battle of Cambrai (1917)

Teacher Model / Pupil Reasoning:

Source S1 : Wilhelm Röntgen's First Medical X-Ray (1895)



The world's first medical radiograph, revealing the bones and wedding ring of Anna Bertha Ludwig Röntgen, taken by Wilhelm Röntgen in December 1895.

Source S2 : Aerial Photographic Map of Trenches at Ypres (1916)



Aerial reconnaissance photograph of frontline, support, and communication trenches zigzagging through shell-cratered terrain near Ypres.

Source S3 : British Soldiers in Waterlogged Mud Trench (1917)



British soldiers standing knee-deep in liquid mud in a flooded communication trench on the Western Front.



Source B: *Source B: Aerial Photographic Map of Trenches at Ypres (1916)*

The Context of the Western Front When World War I broke out in 1914, the British Expeditionary Force (BEF) was sent to Northern France. The terrain of the Western Front was primarily flat agricultural land, particularly in the Ypres Salient and the Somme. Constant artillery bombardment destroyed the natural drainage systems, turning the battlefield into a horrific, waterlogged swamp of mud. The war quickly devolved into a stalemate, with both sides digging hundreds of miles of trenches. This static war of attrition meant that millions of men lived in extremely unhygienic conditions, exposed to the weather, rats, and human waste. Furthermore, high-explosive artillery fire constantly destroyed impassable roads and severed underground telephone cables, cutting off frontline medical officers. The RAMC was forced to rely on runners, runner-dogs, visual signalling flags, and homing pigeons to send urgent casualty reports or request stretcher-bearers. Runners were frequently killed or delayed traversing shell craters, creating catastrophic delays in moving wounded men away from the front line. The unique environment of the Western Front created unprecedented medical challenges that the Royal Army Medical Corps (RAMC) was initially completely unprepared for.



Source C: *Source C: British Soldiers in Waterlogged Mud Trench (1917)*

The Move Towards Aseptic Surgery By the late 1890s, surgeons had moved past Joseph Lister's antiseptic methods (killing germs already in the wound) and towards aseptic surgery, which focused on preventing germs from getting into the operating theatre in the first place. All medical staff were required to meticulously wash their hands, faces, and arms before entering the theatre. Surgeons began wearing sterilised rubber gloves and surgical gowns, which dramatically decreased the rate of infection. Surgical instruments were steam-sterilised using a machine called an autoclave, invented in 1881 by Charles Chamberland. Some theatres even pumped air over a heating system to sterilise it before it entered the room. By 1900, aseptic surgery was the standard, meaning that complex, intrusive operations could be carried out safely in civilian hospitals without the high risk of fatal postoperative infections.

Q1. Identify the main geographical feature of the Western Front.



Source A: *Source A: Wilhelm Röntgen's First Medical X-Ray (1895)*

The Development of X-Rays X-rays were discovered accidentally in 1895 by the German physicist Wilhelm Roentgen, who noticed that the rays passed through flesh but not through bone. They were used for diagnostic purposes almost immediately; by 1896, radiology departments were opened in British hospitals. For example, Dr John Hall-Edwards at Birmingham General Hospital became one of the first doctors to use an x-ray to locate a needle in a woman's hand before operating. However, early x-rays had severe problems: the radiation doses released were 1,500 times stronger than today, causing patients to suffer burns and hair loss. Furthermore, the glass tubes were incredibly fragile, the machines were far too heavy to be moved easily, and taking a single image could take up to 90 minutes, requiring the patient to

stay completely still. While x-rays successfully allowed doctors to diagnose internal problems without cutting the patient open, the machines were heavy, slow, and dangerous, meaning they were not yet suited for the fast-paced, mobile environment of a battlefield.

Q2. Describe the conditions of the trenches during the war of attrition.

Wilhelm Roentgen — German Physicist

Accidentally discovered X-rays in 1895 while experimenting with a cathode ray tube. His discovery allowed doctors to see inside the human body without cutting it open, revolutionizing diagnostics and the treatment of bone fractures.

Key actions: Discovered X-rays in 1895 by covering a cathode-ray tube with black cardboard, noticing that invisible rays could pass through solid objects and expose photographic plates to show bone structures.

Significance: Created the medical science of radiology. For the first time, doctors could diagnose internal fractures, bone diseases, and embedded foreign objects non-invasively, eliminating the need for exploratory surgery.

Limitations: Early X-ray technology was highly primitive and dangerous: exposure times were incredibly long (requiring patients to sit still for up to 90 minutes), the machines were massive and fragile (difficult to move), and they released toxic doses of radiation that caused severe skin burns and hair loss.

Blood Transfusions and Storage Problems James Blundell carried out the first human-to-human blood transfusions in 1818 to save women suffering from blood loss during childbirth. However, early transfusions were highly dangerous because blood types were not understood, often resulting in rejection and death. In 1901, the Austrian doctor Karl Landsteiner revolutionised the process by discovering three blood groups (A, B, and O), with a fourth (AB) found a year later. In 1907, blood group O was identified as the universal blood group. Despite this, a massive problem remained: blood coagulated (clotted) as soon as it left the body, meaning it could not be stored. Because blood could not be stored in a bank, transfusions had to be carried out with the donor and the patient connected directly to each other by a tube in the exact same room. This made dealing with mass casualties on a battlefield virtually impossible.

KNOWLEDGE CHECK (Q3)

What was the main limitation of performing blood transfusions in hospital settings before the discovery of sodium citrate in 1915?

- ☐ A. Blood types had not yet been discovered, so transfusions were always fatal.
- ☐ B. Blood could not be stored because it quickly coagulated (clotted), meaning a donor had to be physically present.
- ☐ C. Syringes had not yet been invented.
- ☐ D. The Church banned the transfer of blood between two people.

Q4. Explain how artillery bombardment worsened the medical situation.

Q5. To what extent was the RAMC prepared for the medical challenges of the Western Front?

Karl Landsteiner — Biologist and Physician

Discovered the existence of different human blood groups (A, B, O) in 1901. This breakthrough explained why some blood transfusions were fatal while others succeeded, making donor matching possible and paving the way for safe transfusions.

Key actions: Discovered blood groups (A, B, and O) in 1901 (with colleagues discovering AB in 1902), proving that donor and recipient blood must be matched to prevent fatal immune reactions.

Significance: Transformed blood transfusions from a highly lethal, outlawed gamble into a safe, scientifically controlled, and routine clinical procedure.

Limitations: He did not solve the critical problem of clotting and storage. Because blood clotted immediately outside the body, transfusions still had to be performed directly on a person-to-person basis with both donor and patient present, which was logistically impossible on a chaotic battlefield.

Pair & Share Activity

Q6. Which pre-war medical development was the most critical for the survival of soldiers on the Western Front: the rise of aseptic surgery, the invention of mobile X-rays, or the progress in storing and transfusing blood?

Source A

From a lecture given by a British surgeon to the Royal College of Surgeons, 1912.

"Transfusion today remains a procedure fraught with difficulty. The donor and the recipient must lie side by side, connected directly by rubber tubing, for the blood clots within minutes of leaving the vein. Should a patient require a transfusion at short notice, we are entirely dependent on finding a willing and compatible donor already present within the building."

Source B

From the casebook of a London hospital physician, recording a transfusion carried out in 1913.

"Mrs H., having lost a dangerous quantity of blood following childbirth, was saved by a direct transfusion from her husband, who was tested beforehand using Dr Landsteiner's method and found to be of a compatible group. Without his presence in the building at the time, the procedure could not have been attempted."

Provenance Scaffolding:

Source A is a lecture given by a surgeon to expert colleagues; consider how this affects its reliability compared to a private record. Source B is a physician's private casebook entry describing one real case; what does it confirm about the limits described in Source A?

Q7. 1 (a). How useful are Sources A and B for an enquiry into the problem of storing blood for transfusions before the First World War? Explain your answer, using Sources A and B and your

knowledge of the historical context. (8 marks)

GCSE Exam Practice

Q8. Describe one feature of the First Battle of Ypres. (2 marks)

Q9. Describe one feature of the Battle of the Somme. (2 marks)

L2: KT5.2: How did the trench environment create new medical challenges?

LEARNING OBJECTIVES

- ☐ Explain the causes, symptoms, and preventive strategies developed to combat trench foot and trench fever.
- ☐ Analyze the impact of weaponry on injuries, specifically focusing on shrapnel, head injuries, and gas gangrene from soil.

The First World War on the Western Front forced the British army into a static, defensive conflict across Flanders and northern France. Instead of a fast-moving war of movement, soldiers lived and fought in a complex network of muddy trenches. This brutal environment not only produced staggering casualties in major battles like the Somme and Ypres, but the devastated terrain and narrow trenches created immense problems for the medical services trying to transport and treat the wounded.

Did you know?

- Ypres Salient: A vulnerable 'bulge' in the Allied trench line in Belgium that was surrounded by the enemy on three sides.
- Zig-zag: The pattern in which trenches were deliberately dug to prevent enemy fire and explosive blasts from travelling straight down the line.
- 2.5 miles: The length of the underground tunnel network dug into the chalk at Arras to safely shelter troops and a hospital in 1916.

Recall & Retrieval

- | | |
|---|--|
| 1. When did the First World War begin and end? | 2. What was the 'Western Front'? |
| 3. Name the three major battles fought by the British on the Western Front. | 4. What was 'No Man's Land'? |
| 5. Name the pattern in which trenches were dug. | 6. What was the purpose of the 'duckboards' in the trenches? |
| 7. Who discovered the cause of cholera in 1854? | 8. Which anesthetic was discovered by James Simpson in 1847? |
| 9. Explain why the zig-zag pattern of trenches was crucial for survival. | 10. What were the three main parallel lines of trenches? |


Vocabulary Check

The Golden Sentence: Choose TWO terms from the word bank. Write ONE grammatically sophisticated, historically accurate sentence connecting them using a causal conjunction (because, although, or consequently):

Trench Foot | Whale Oil | Trench Fever | Lice | Shell Shock | Dysentery

Teacher Model / Pupil Sentence:

Source S1 : RAMC Clinical Photograph of Trench Foot (c.1916)



A British soldier suffering from severe trench foot: swollen, blistered, and gangrenous tissue resulting from prolonged immersion in cold water.

Source S2 : British Machine Gunners in PH Gas Helmets (Somme, 1916)




Vickers machine-gun crew of the Cheshire Regiment wearing Phenate-Hexamine (PH) anti-gas hoods in a chalk trench near Ovillers.

Source S3 : Stretcher Bearers Carrying Casualty Through Deep Mud (Passchendaele, 1917)



Four exhausted British stretcher bearers wading waist-deep through liquid mud, struggling to carry a wounded soldier across the devastated battlefield.



Source A: Source A: RAMC Clinical Photograph of Trench Foot (c.1916)

Trench Environment and Illnesses The trenches were breeding grounds for disease. The most notorious environmental illness was Trench Foot, caused by standing in cold, waterlogged mud for days without changing socks. It caused feet to swell, blister, and turn gangrenous, often requiring amputation. The army eventually ordered soldiers to change socks twice a day and rub whale oil on their feet. Another major issue was Trench Fever, which caused severe joint pain and high fevers. It took years for doctors to realize it was transmitted by the millions of body lice living in the soldiers' unwashed uniforms. To combat this, the army set up massive delousing stations and bathhouses to disinfect clothing with hot steam.

The Theatre of War (Key Battles) The Ypres Salient: A vulnerable 'bulge' in the Allied line in Belgium, surrounded by the enemy on three sides, allowing the Germans to fire down from the higher ground. The First Battle of Ypres (Autumn 1914) was critical; the British held onto the salient to stop the Germans advancing to the sea, successfully retaining control of the vital English Channel ports. Major battles later included the Second Battle of Ypres (1915), which saw the first use of chlorine gas by the Germans, and the Third Battle of Ypres (Passchendaele, 1917), where heavy rain turned the ground into a deadly, waterlogged sea of mud. During the fight for Hill 60 (April 1915), the British successfully used offensive mining to tunnel under and blow up the German positions. The Somme (1916): A massive British attack aiming to relieve pressure on the French at Verdun. It became a medical nightmare, resulting in nearly 60,000 British casualties on the very first day and over 400,000 by the end of the campaign. Arras (1917): Because the ground was chalky and easy to tunnel through, British and New Zealand miners dug a vast 2.5-mile network of tunnels. This safely sheltered 25,000 troops and included a fully functioning underground hospital.

Q1. Identify the cause of Trench Foot.



Source B: Source B: British Machine Gunners in PH Gas Helmets (Somme, 1916)

The Trench System and its Organisation The frontline trench was the firing line where attacks were launched from. Soldiers stood on a firestep to shoot over the sandbag parapet. The support trench was located about 80 metres behind the frontline; troops would retreat here if the frontline came under heavy attack. The reserve trench was at least 100 metres further back, holding backup troops ready to be mobilised for a counter-attack. Communication trenches linked these rows together, allowing the safe movement of men, equipment, and wounded soldiers between the front and the rear. To prevent enemy fire or the blast wave from an exploding shell from travelling straight down the line, trenches were deliberately dug in a zig-zag pattern. The complex, narrow, and zig-zag design of the trench system made it incredibly difficult and physically exhausting for stretcher-bearers to safely manoeuvre seriously wounded men around the tight corners while under fire.

Q2. Describe the symptoms of Trench Fever.



Source C: Source C: Stretcher Bearers Carrying Casualty Through Deep Mud (Passchendaele, 1917)

Significance of the Terrain, Transport and Communications Infrastructure Constant artillery shelling destroyed the region's roads and natural drainage systems. This left the landscape heavily cratered and filled with deep, waterlogged mud. Heavy motor ambulances frequently got permanently stuck in the

deep mud, meaning the army had to rely heavily on horse-drawn ambulance wagons, which often required six horses to pull them through the sludge. These wagons were violently bumpy and often worsened severe injuries. Stretcher-bearers (working in teams of four to six) faced the exhausting, dangerous task of carrying the wounded through muddy shell-holes in No Man's Land, often under heavy fire or in the dark. Once inside the trenches, it was incredibly difficult to manoeuvre stretchers due to crowded troops moving in opposite directions. Problems with communications infrastructure severely compounded transport difficulties. Constant artillery bombardments severed buried telephone lines, cutting off frontline medical posts from Casualty Clearing Stations. Consequently, medical officers had to rely on runners, runner-dogs, or homing pigeons to send messages. Runners were frequently killed or delayed in shell craters, leaving rear hospitals with no advance warning of mass casualties and causing severe bottlenecks where wounded men lay waiting in cold mud for hours. The devastated terrain and crippled communications severely slowed evacuation; this delay caused men to die from blood loss or shock, and allowed deep infections like gas gangrene to take hold before they could reach surgical treatment.

KNOWLEDGE CHECK (Q3)

How did the British Army attempt to prevent soldiers from developing trench foot in wet and muddy conditions?

- ☐ A. By providing them with waterproof rubber wellington boots.
- ☐ B. By issuing orders for soldiers to rub their feet with whale oil and regularly change their socks.
- ☐ C. By administering a trench foot vaccine to all soldiers.
- ☐ D. By heating the trenches with underground pipes.

Q4. Explain how the army attempted to prevent Trench Fever.

Q5. Evaluate the effectiveness of military discipline in preventing environmental illnesses.

Pair & Share Activity

Q6. Which aspect of the trench environment presented the most difficult challenge for the Royal Army Medical Corps (RAMC): the physical environment (producing trench foot), the biological vectors (producing trench fever), or the psychological strain of industrial warfare (producing shell shock)?

Source A

From the personal memories of William Easton, an 18-year-old volunteer with the East Anglian Field Ambulance, describing the Passchendaele/Ypres campaign in 1917.

"During the third offensive at Ypres, we navigated paths choked with thick,

Source B

An official archival photograph taken on the Somme in 1916, showing a horse-drawn ambulance navigating a deeply rutted, waterlogged dirt road.

hip-deep mire. In the devastated sector of Hooze, we struggled to relocate casualties. The physical exertion of a single carry left us utterly spent; delivering just two or three wounded men to safety daily was our absolute limit. We worked in teams of four, using shoulder straps to stabilize the stretcher and prevent agonising falls."



Provenance Scaffolding:

Think about who wrote or produced this source. For Source A, it's an East Anglian Field Ambulance worker at Ypres. Does this personal, on-the-ground experience make it more reliable for understanding the physical toll? For Source B, it's an official photograph from the Somme. Was it meant to document reality, or perhaps carefully framed for morale?

Q7. 2 (a). How useful are Sources A and B for an enquiry into the impact of the terrain on the transport of the wounded on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

GCSE Exam Practice

Q8. Describe one feature of the trench system. (2 marks)

Q9. Describe one feature of trench foot. (2 marks)

L3: KT5.3: What new illnesses and wounds did soldiers face?

LEARNING OBJECTIVES

- ☐ Explain the connection between artillery weaponry, shrapnel, and the introduction of the steel Brodie helmet in 1915.
- ☐ Analyse how the muddy, manured environment of Flanders led to severe wound infections like gas gangrene and tetanus.

Living and fighting in the waterlogged trenches of the Western Front exposed soldiers to horrifying new medical dangers. Beyond the immediate, devastating trauma of being hit by shrapnel or rapid-fire machine guns, men faced the constant, invisible threat of deadly soil-borne infections like gas gangrene. Furthermore, the brutal, freezing environment itself caused severe illnesses like trench foot and shellshock, while the introduction of poison gas added a terrifying new psychological and physical dimension to casualty care.

Did you know?

- 58%: The percentage of all wounds on the Western Front caused by high-explosive shells and shrapnel.
- 1915: The year the steel Brodie helmet was introduced, drastically reducing fatal head injuries by 80%.
- 80,000: The estimated number of British troops who suffered from the severe psychological trauma of shellshock.

Recall & Retrieval

1. What was 'No Man's Land'?
2. Name the pattern in which trenches were dug.
3. What was the purpose of the 'duckboards' in the trenches?
4. What is 'Trench Foot'?
5. Which weapon caused the most casualties on the Western Front?
6. Why were wounds on the Western Front highly prone to severe infection?
7. Who discovered Penicillin by accident in 1928?
8. What structure did Watson and Crick discover in 1953?
9. What preventative measure was introduced to stop Trench Foot?
10. What was 'Shell Shock'?

Vocabulary Check


Conceptual Classification: Categorise the terms from the word bank into the two historical boxes below:

Shrapnel | Brodie Helmet | Gas Gangrene | Chlorine Gas | Phosgene Gas | Mustard Gas

Power, Governance & Warfare

Economy, Trade & Society

Source S1 : Diagnostic Radiograph of Lodged Bullet Localised with Lead-Wire Grid



An authentic radiograph showing a metallic bullet lodged deep in the skull, mapped precisely using an external lead-wire localization grid.

Source S2 : Brodie Steel Helmet with Shrapnel Dent (1916)



A British Mark I Brodie steel helmet displaying a deep dent from an exploding artillery shrapnel ball that would have killed an unprotected soldier.

Source S3 : John Singer Sargent: Gassed (1919)




John Singer Sargent masterwork Gassed: a line of blinded soldiers in mustard gas bandages guided toward a casualty clearing station dressing tent.

Wounds and Infections The weapons of WWI caused devastating injuries. High-explosive artillery shells caused over 60% of all wounds. When these shells exploded, hot jagged shrapnel tore through flesh, dragging the muddy, bacteria-filled uniform deep into the wound. This immediately caused deadly infections like Gas Gangrene, which could kill a man within 24 hours. Soldiers also faced the horrors of poison gas. Chlorine, Phosgene, and Mustard gas caused blinding, severe lung damage, and horrific internal and external blisters. While gas only caused about 6,000 British deaths, the psychological terror it inflicted was immense, requiring soldiers to carry heavy, uncomfortable gas masks at all times.

Illnesses from the Trench Environment Trench foot was a painful swelling caused by standing in cold mud and water for hours. It restricted blood flow, eventually causing the flesh to rot, a condition known as gangrene. Preventative orders included rubbing whale oil into the feet and changing socks regularly, but severe cases usually required amputation. Trench fever was a debilitating condition causing severe flu-like symptoms, high temperatures, and aching muscles. It was incredibly widespread, affecting an estimated half a million men. In 1918, doctors finally proved it was spread by lice living in the seams of clothing, leading to the creation of special delousing stations. Shellshock was a poorly understood psychological condition caused by the constant trauma of bombardment and the noise of the big guns. Symptoms included nightmares, loss of speech, uncontrollable shaking, and complete mental breakdown. It affected an estimated 80,000 British troops, with some being treated at the specialist Craiglockhart Hospital in Edinburgh. The army officially recorded the condition under the code NYD.N (Not Yet Diagnosed, Nervous). (Key Detail: You must use the exact military acronym 'NYD.N.' to show precise historical knowledge of how the army categorized shellshock). Because the brutal trench environment itself caused massive casualties, the medical corps had to dedicate huge resources to preventative measures just to keep the army fit enough to fight.

Q1. Identify the weapon that caused over 60% of all wounds on the Western Front.



Source A: Source A: Diagnostic Radiograph of Lodged Bullet Localised with Lead-Wire Grid

The Devastation of Shrapnel and Explosives High-explosive shells and shrapnel (jagged metal fragments packed inside artillery shells) were the greatest killers, responsible for 58% of all wounds. They scattered fast-moving metal over a massive area, causing horrific trauma. Machine guns and rifles accounted for roughly 39% of wounds. A machine gun could fire between 450 and 600 rounds a minute, instantly fracturing bones and piercing major internal organs. The sheer destructive power of modern industrial weapons meant soldiers suffered complex, jagged injuries and massive blood loss that traditional military surgeons had never encountered on this scale before.

Q2. Describe the physical and psychological effects of poison gas.

The Deadly Threat of Infection The farmland of France and Belgium had been heavily fertilised with manure before the war, meaning the soil was packed with lethal bacteria that caused tetanus and gas gangrene. When shrapnel or a bullet pierced a soldier, it dragged dirty uniform fabric and infected mud deep into the wound. While the danger of tetanus was successfully reduced by administering anti-tetanus injections from late 1914, there was no initial cure for gas gangrene. This infection rapidly produced gas inside the dying tissue, turning flesh black and emitting a foul smell, and could kill a man within a day. Because lethal bacteria were instantly driven deep into the body at the moment of injury, traditional surface antiseptics entirely failed, forcing doctors to find radically new surgical methods to stop the infection.

KNOWLEDGE CHECK (Q3)

Why were deep shrapnel wounds on the Western Front almost guaranteed to develop the deadly infection known as gas gangrene?

- ☐ A. Because the German army dipped their bullets in poison before firing them.
- ☐ B. Because the French farmland had been heavily fertilized with manure for centuries, filling the soil with deadly tetanus and gangrene bacteria.
- ☐ C. Because the British soldiers refused to wash their uniforms.
- ☐ D. Because chlorine gas from chemical attacks infected the wounds.

Q4. Explain why shrapnel wounds almost always became infected.



Source B: Source B: Brodie Steel Helmet with Shrapnel Dent (1916)

Head Injuries and the Brodie Helmet At the start of the war, British soldiers only wore soft cloth caps. Because men stood in trenches, their heads were highly exposed, and approximately 20% of all wounds were to the head, face, and neck. In 1915, the British army introduced the steel Brodie helmet, which included a strong strap to prevent it from being blown off in an explosion. The widespread introduction of the Brodie helmet was a monumental preventative success, rapidly reducing the number of fatal head wounds caused by shrapnel by an estimated 80%.

Q5. To what extent could doctors successfully treat deep infections during WWI?



Source C: *Source C: John Singer Sargent: Gassed (1919)*

The Terror of Poison Gas Chlorine gas was first used by the Germans at the Second Battle of Ypres (April 1915), causing victims to slowly suffocate as their lungs filled with fluid. Before gas masks were officially issued in July 1915, soldiers urinated on cotton pads and pressed them to their faces to neutralise the chemical. Phosgene was introduced later in 1915; it was faster-acting and more deadly, capable of killing an exposed person within two days. Mustard gas (1917) was an odourless gas that caused severe internal and external blistering and could burn the skin straight through clothing. Although poison gas was a terrifying psychological weapon, the rapid development of effective gas masks meant it was actually the cause of under 5% of British deaths (around 6,000 soldiers), making it far less lethal than artillery fire.

Pair & Share Activity

Q6. Which posed a more difficult medical challenge for the Royal Army Medical Corps (RAMC) on the Western Front: the physical injuries caused by industrial weaponry (such as shrapnel wounds and head trauma) or the environmental infections caused by the Flanders terrain (such as gas gangrene, tetanus, and trench foot)?

Source A

From a wartime notebook of Lance Sergeant Elmer Cotton in 1915, providing a private, firsthand account of a gas attack.

"A chlorine discharge floods the victim's respiratory tract, effectively suffocating them while standing on dry ground. The resulting symptoms are excruciating: a blinding headache, an overwhelming craving for water-

Source B

An official photograph capturing blinded soldiers forming a chain outside an Advanced Dressing Station.

though drinking it brings immediate fatality-and a severe, stabbing sensation in the lungs. Patients continuously expectorate a distinctive greenish fluid. It represents a truly horrific way to perish."



Provenance Scaffolding:

For Source A, consider that it is from a wartime notebook of Lance Sergeant Elmer Cotton in 1915. This is a private, unedited firsthand account. How does this affect its usefulness? For Source B, an official photograph showing blinded soldiers forming a chain. Photographs capture reality but can also be staged. Does the clear visual evidence of the gas's impact make it highly useful regardless?

Q7. 2 (a). How useful are Sources A and B for an enquiry into the effects of poison gas attacks on soldiers on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

GCSE Exam Practice

Q8. Describe one feature of the effects of poison gas. (2 marks)

Q9. Describe one feature of shrapnel wounds. (2 marks)

L4: KT5.4: How did the RAMC and FANY operate the chain of evacuation?

LEARNING OBJECTIVES

- ☐ Explain the four main stages of the Western Front chain of evacuation and the mode of transport between them.
- ☐ Evaluate the roles, significance, and contributions of the RAMC and FANY on the Western Front.

The devastating environment and weaponry of the Western Front required a highly organised system to rescue and treat casualties. The Royal Army Medical Corps (RAMC) and nursing volunteers like the FANY worked tirelessly to transport men through a multi-stage 'chain of evacuation'. From exhausted stretcher-bearers navigating deep mud to the fully equipped underground hospital at Arras, the medical services continually adapted to save lives under unimaginable pressure.

Did you know?

- Triage: The vital system used at the Casualty Clearing Station to divide wounded soldiers into three categories to prioritise medical treatment.
- 512: The number of motor ambulances purchased through The Times newspaper's public appeal in 1914.
- 700: The number of stretcher beds available inside the fully electrified underground hospital at Arras.

Recall & Retrieval

- | | |
|---|--|
| 1. What is 'Trench Foot'? | 2. Which weapon caused the most casualties on the Western Front? |
| 3. Why were wounds on the Western Front highly prone to severe infection? | 4. What does RAMC stand for? |
| 5. What was the role of the FANY? | 6. Describe the 'Chain of Evacuation' on the Western Front. |
| 7. What lifestyle choice was definitively linked to lung cancer in the 1950s? | 8. What was established in Britain in 1948 to provide free medical care? |
| 9. What was the primary role of the Casualty Clearing Station (CCS)? | 10. Where were the Base Hospitals located and what did they do? |

Vocabulary Check


Spot the Deliberate Error: Read the statement below. One historical fact or vocabulary concept has been deliberately falsified. Underline the error and explain the accurate historical reality below:

Chain of Evacuation | Regimental Aid Post (RAP) | Field Ambulance | Casualty Clearing Station (CCS)
| Base Hospitals | First Aid Nursing Yeomanry (FANY)

Challenge: Write ONE statement containing a deliberate historical misconception using **Chain of Evacuation** or **Regimental Aid Post (RAP)**. Identify and correct the error:


Teacher Model / Historical Correction:

Source S1 : Reception of the Wounded at 41st Casualty Clearing Station (1918)



Wounded soldiers on stretchers filling the marquee triage reception area at the 41st Casualty Clearing Station during the final British advance.

Source S2 : Ambulance Train Wards on the Western Front (1917)




British ambulance train interior fitted with tiered sprung bunks, heating, and pharmacy facilities to transport stabilized casualties to Base Hospitals.

Source S3 : Underground Hospital Ward at Arras (1917)



The underground hospital complex in the chalk quarries of Arras: fully protected subterranean hospital beds with electric lighting and running water.




Source A: *Source A: Reception of the Wounded at 41st Casualty Clearing Station (1918)*

The Chain of Evacuation To deal with the massive casualties, the RAMC designed a highly organized 'Chain of Evacuation'. Stretcher bearers would rescue men from No Man's Land and take them to the Regimental Aid Post (RAP) just behind the frontline trench, which provided basic first aid. From there, horse-drawn or motorized ambulances transported the wounded to Advanced Dressing Stations (ADS). Those who needed surgery were sent further back to Casualty Clearing Stations (CCS), which were large tented hospitals located near railway lines. The CCS dealt with the most critical, life-saving surgeries, particularly amputations to stop gangrene. If a soldier survived, they were put on a hospital train to a Base Hospital near the French coast, where specialists worked, before being shipped back to 'Blighty' (Britain).

The Work of the RAMC and Nurses The RAMC was the branch of the army responsible for all medical care, from treating the wounded to maintaining sanitation to keep the men healthy. Due to the sheer scale of casualties, the RAMC expanded massively from 9,000 men in 1914 to 113,000 by 1918. Professional nurses from Queen Alexandra's Imperial Military Nursing Service also grew from 300 to 10,000. The First Aid Nursing Yeomanry (FANY) was a women's voluntary organisation; they famously drove motor ambulances, transported supplies to the frontline, set up cinemas, and even operated a mobile bath unit that could bathe up to 40 men an hour. The rapid expansion of the RAMC and the vital inclusion of female volunteers were absolutely essential to prevent the medical system from entirely collapsing under the unprecedented, staggering number of casualties.

Q1. Identify the first stage of the Chain of Evacuation.



Source B: *Source B: Ambulance Train Wards on the Western Front (1917)*

Transport in the Chain of Evacuation Initially, the army relied on horse-drawn ambulance wagons, which were violently bumpy, could only carry a few men, and often worsened severe injuries. A public appeal by The Times newspaper raised money for 512 new motor ambulances, which arrived in October 1914. However, motor vehicles frequently got stuck in the deep mud, meaning horse-drawn wagons (sometimes needing six horses) and exhausted human stretcher-bearers (working in teams of four or six while under enemy fire) remained crucial. To move the wounded to Base Hospitals, specially designed ambulance trains and canal barges were used, with some trains capable of carrying up to 800 casualties at once. The devastated, cratered terrain severely hindered transport; this delay and the bumpy journeys often exacerbated injuries and caused fatal blood loss before soldiers could ever reach life-saving surgery.

Q2. Describe the role of Casualty Clearing Stations (CCS).

Stages of Treatment: RAP and Dressing Stations The Regimental Aid Post (RAP) was located within 200m of the frontline, often in communication trenches or dugouts. A Regimental Medical Officer provided immediate first aid to get walking wounded back to the fight, but could not perform surgery on serious injuries. The wounded were then moved to an Advanced Dressing Station (ADS) or Main Dressing Station (MDS), located about 400m to half a mile back in abandoned buildings, dug-outs, or tents. Here, Field Ambulance units (staffed by medical officers, orderlies, and eventually nurses) provided better bandages and pain relief, keeping men for a maximum of one week. These early stages were vital for preventing frontline trenches from becoming blocked with wounded men, quickly patching up minor injuries, and organising the transport of serious cases further back.

Q3. Explain how patients were transported between the different stages of the evacuation chain.



Source C: *Source C: Underground Hospital Ward at Arras (1917)*

Stages of Treatment: CCS and Base Hospitals The Casualty Clearing Station (CCS) was located 7 to 12 miles back, safely out of artillery range but near railway lines. Here, patients underwent triage, being sorted into three groups: the walking wounded, those needing hospital treatment, and those with no chance of recovery. Because deadly gas gangrene set in so quickly, the CCS soon took over the role of performing critical, life-saving surgery (like amputations) from the Base Hospitals. Base Hospitals, located near the French and Belgian coastal ports, eventually focused on the longer-term recovery of patients and developed specialist wards for head wounds or gas poisoning. By shifting life-saving surgical operations forward to the CCS, the medical services successfully reduced the time wounded men had to wait for surgery, drastically decreasing deaths from infection.

KNOWLEDGE CHECK (Q4)

What was the main difference between the medical treatments performed at a Regimental Aid Post (RAP) and a Casualty Clearing Station (CCS)?

- ☐ A. The RAP performed complex surgery, while the CCS only applied bandages.
- ☐ B. The RAP provided immediate first aid near the frontline, while the CCS was equipped with operating theatres and X-ray machines to perform life-saving surgery.
- ☐ C. The RAP was for officers only, while the CCS was for regular soldiers.
- ☐ D. The RAP was located in Britain, while the CCS was in France.

Q5. Evaluate the significance of the Chain of Evacuation.

The Underground Hospital at Arras In 1916, British and Commonwealth troops linked existing chalk tunnels and quarries beneath the town of Arras. They created a massive underground town that included a fully functioning hospital (nicknamed Thompson's Cave) within 800m of tunnels. It contained 700 spaces for stretchers to be used as beds, operating theatres, rest stations, a mortuary, and was supplied with electricity and piped water. The underground hospital at Arras was highly significant because it functioned as a fully equipped medical facility incredibly close to the frontline, while remaining completely safe from enemy artillery bombardment.

Pair & Share Activity

Q6. Was the medical service's success on the Western Front driven more by the highly organized military logistics of the RAMC, or by the courage and adaptability of voluntary organizations like the FANY?

Source A

From the 1919 published autobiography of Pat Beauchamp, a volunteer FANY ambulance driver.

"With the Somme offensive raging, dozens of transport cars would form a convoy to meet incoming hospital trains at the railway siding. Once positioned, stretcher bearers loaded our vehicles with four severely injured, non-walking cases. Traversing the ruined, potholed cobblestone roads was an agonizing ordeal. The violent jolts caused immense suffering, yet we were forced to drive slowly to protect those inside."

Source B

A 1916 archival photograph from the Somme showing a motor ambulance navigating a flooded road.



Provenance Scaffolding:

Source A is from the 1919 published autobiography of a FANY ambulance driver. It's a memoir published right after the war. How might writing for a public audience affect her description of the difficulties?
Source B is a 1916 photograph from the Somme. Official photographs were subject to censorship. Does its depiction of flooded roads match your own knowledge of the Somme battlefield?

Q7. 2 (a). How useful are Sources A and B for an enquiry into the problems of transporting the wounded on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

GCSE Exam Practice

Q8. Describe one feature of the FANY. (2 marks)

Q9. Describe one feature of the Chain of Evacuation. (2 marks)

L5: KT5.5: What incredible medical advances were forged on the Western Front?

LEARNING OBJECTIVES

- ☐ Explain how the Thomas splint and mobile X-ray units solved critical evacuation and surgical challenges.
- ☐ Analyze the development of blood transfusion and the significance of the first blood bank at the Battle of Cambrai.

The horrific new weapons of the First World War created devastating, complex injuries that traditional 19th-century surgeons simply did not know how to cure. Faced with thousands of men rapidly dying from gas gangrene, massive blood loss, and shattered limbs, the medical services were forced to turn the Western Front into a vast laboratory. Driven by sheer desperation, they successfully pioneered radical new surgical techniques, mobile diagnostic technology, and miraculous methods for storing blood directly on the battlefield.

Did you know?

- 82%: The dramatically increased survival rate for soldiers with compound leg fractures after the Thomas Splint was introduced in late 1915.
- 1917: The year Oswald Hope Robertson established the world's first 'blood depot' (blood bank) before the Battle of Cambrai.
- Debridement: The rapid surgical technique of cutting away dead, damaged, and infected tissue from around a wound to prevent the spread of gas gangrene.

Recall & Retrieval

- | | |
|---|---|
| 1. What does RAMC stand for? | 2. What was the role of the FANY? |
| 3. Describe the 'Chain of Evacuation' on the Western Front. | 4. What was the Thomas Splint? |
| 5. What chemical was discovered in 1915 to prevent blood from clotting? | 6. What did the discovery of blood storage allow surgeons to do? |
| 7. Who discovered the structure of DNA in 1953? | 8. Who introduced antiseptic surgery using carbolic acid in 1865? |
| 9. How did WWI affect the use of X-rays in medicine? | 10. What was the 'Carrel-Dakin' method? |

Vocabulary Check

The Odd One Out: Circle *ONE* term that does not belong with the others. Explain your reasoning: What historical connection links the other terms, and why is your chosen term different?

Thomas Splint | Mobile X-ray Unit | Carrel-Dakin Method | Sodium Citrate | Blood Depot | Brain Surgery

Teacher Model / Pupil Reasoning:

Source S1 : The Thomas Splint Demonstrated on a Wounded Soldier (1916)



Demonstration of the Thomas Splint: a metal and leather traction frame that pulled the broken femur straight, preventing severed arteries and shock.

Source S2 : Geoffrey Keynes & Robertson's Portable Blood Transfusion Kit (1917)



A portable wooden field transfusion case designed by Lieutenant Geoffrey Keynes (RAMC) with calibrated glass bottles and anticoagulant regulators.

Source S3 : Walter Yeo's Facial Reconstruction by Harold Gillies (1917)



Walter Yeo before and after facial reconstruction by Harold Gillies using the 'tubed pedicle' skin flap technique at Queen's Hospital, Sidcup.

Medical Advances on the Western Front The sheer volume of casualties on the Western Front forced doctors to innovate rapidly. One of the greatest breakthroughs was the Thomas Splint, a metal frame designed by Hugh Owen Thomas. Before 1915, 80% of soldiers with broken femurs died from shock or bleeding. The Thomas Splint kept the leg perfectly still during transport, increasing the survival rate to an incredible 82%. Another major advance was in blood transfusions. At the start of the war, transfusions had to be done arm-to-arm. However, in 1915, Richard Lewisohn discovered that adding sodium citrate to blood stopped it from clotting, allowing it to be stored on ice. This led to the creation of the first blood banks at the Battle of Cambrai in 1917, allowing surgeons to treat shock and perform complex surgeries.

Hugh Owen Thomas — Orthopaedic Surgeon

Invented the Thomas Splint in the late 19th century. During WWI, his invention was widely adopted to immobilize shattered femurs on the Western Front, stopping the broken bone from causing fatal internal bleeding and raising survival rates from 20% to 82%.

Key actions: Designed the Thomas splint—a rigid metal leg frame featuring a leather ring at the hip to pull the leg under traction and keep the femur fully immobilized.

Significance: When adopted on the Western Front in 1916, his splint revolutionized the treatment of femur fractures, dropping the mortality rate from compound leg wounds from 80% to just 18% by stopping internal bleeding, bone movement, and shock.

Limitations: He designed and used the splint in his civilian practice decades before the First World War, but his work was entirely ignored by the British military and medical establishment during his lifetime due to professional conservatism. It was only the desperate, high-mortality crisis of trench warfare that forced its retrospective, widespread adoption in 1916.

New Techniques in the Treatment of Wounds and Infection Because traditional aseptic surgery was impossible in dirty casualty stations and carbolic acid simply did not kill gas gangrene bacteria effectively, radical new solutions had to be found. Debridement (Wound Excision): Surgeons began aggressively cutting away all dead, damaged, and infected tissue from around a wound to physically remove the gangrene. This had to be done incredibly quickly before the wound was stitched up to stop the infection from spreading into healthy flesh. The Carrel-Dakin method: If debridement wasn't enough, surgeons used a system where tubes pumped a chemical sterilised salt solution directly into deep wounds to continually wash out and kill the infection-causing bacteria. The method was effective but highly difficult, as the solution only stayed fresh for six hours and had to be made as it was needed. Amputation: If antiseptics or wound excision failed to stop the spread of infection, the limb was amputated to save the man's life. By 1918, over 240,000 men had lost limbs. Because infection spread so

rapidly in the heavily manured soil of the Western Front, surgeons were forced to abandon slow peacetime methods and instead pioneer radical, invasive techniques to physically flush out or cut away deadly bacteria before it killed the patient.

Q1. Identify the device designed by Hugh Owen Thomas.



Source A: Source A: The Thomas Splint Demonstrated on a Wounded Soldier (1916)

The Thomas Splint In 1914 and 1915, a soldier with a gunshot or shrapnel wound that fractured their femur (thigh bone) only had a 20% chance of survival. Traditional splints did not keep the leg straight, meaning the jagged bone caused massive internal bleeding and shock during bumpy transport. In December 1915, Robert Jones introduced the Thomas Splint (a device originally designed by his uncle, Hugh Thomas) to the Western Front. This splint rigidly pulled the leg lengthways, completely stopping the broken bones from grinding on each other. The introduction of the Thomas Splint was one of the greatest medical successes of the war. By keeping the leg perfectly still, it prevented fatal blood loss and shock, miraculously increasing the survival rate for thigh fractures from 20% to 82%.

KNOWLEDGE CHECK (Q2)

How did the implementation of the Thomas splint on the Western Front in 1916 directly reduce mortality rates among wounded soldiers?

- ☐ A. It stopped the spread of infection by killing the bacteria that caused gas gangrene.
- ☐ B. It cured shell shock by holding the soldier firmly in place.
- ☐ C. It kept fractured leg bones rigidly in place during transport, preventing fatal blood loss and shock.
- ☐ D. It provided a steady drip of blood for transfusions on the battlefield.

Q3. Describe how Richard Lewisohn improved blood transfusions.

The Use of Mobile X-Ray Units X-rays were absolutely essential for accurately locating bullets and deeply embedded shrapnel so they could be removed before infection set in. While static machines were used in Base Hospitals, six mobile x-ray units (vans) were developed to operate in the British sector so they could be driven directly to Casualty Clearing Stations closer to the fighting. There were limitations:

the glass tubes overheated very quickly (meaning machines could only be used for an hour before needing to cool down), and they could not detect fragments of dirty clothing driven into wounds. Even though the mobile images were not quite as clear as static machines and they overheated, they were incredibly significant because they allowed life-saving diagnostic technology to be transported directly to the frontline, heavily reducing deaths from infection.

Q4. Explain how the Thomas Splint improved survival rates.



Source B: Source B: Geoffrey Keynes & Robertson's Portable Blood Transfusion Kit (1917)

Blood Transfusions and the Blood Bank at Cambrai Severe blood loss caused victims' bodies to go into fatal shock. Dr Lawrence Bruce Robertson pioneered the use of the indirect blood transfusion method (using a syringe and tube) at Boulogne in 1915. However, early transfusions were heavily limited because blood could not be stored and would quickly clot. Scientists rapidly discovered chemical solutions: In 1915, Richard Lewisohn added sodium citrate to stop blood from clotting. Richard Weil then discovered this allowed blood to be refrigerated for up to two days. In 1916, Francis Rous and James Turner added citrate glucose, a monumental breakthrough that allowed blood to be safely stored for up to four weeks. These chemical discoveries allowed the American doctor Oswald Hope Robertson to establish the world's first 'blood depot' (blood bank) before the Battle of Cambrai in 1917. He stored 22 units of pre-donated universal Type O blood in glass bottles packed with ice and sawdust, successfully treating severely shocked soldiers and proving that stored blood could save lives on the battlefield.

Q5. To what extent did the Western Front accelerate medical progress?

Oswald Hope Robertson — Medical Researcher

An American medical researcher who served with the US Army Medical Corps attached to the British Army during WWI. He successfully developed the first functional blood bank.

Key actions: Set up a forward supply depot of preserved donor blood before the Battle of Cambrai in 1917.

Significance: Created the world's first functional blood bank. Utilized sodium citrate to prevent clotting and dextrose/glucose to preserve red blood cells, allowing universal Group O blood to be safely stockpiled for up to 21 days.

Limitations: Preserved blood could only be kept fresh for a maximum of three weeks and required delicate, constant refrigeration in basic wooden crates packed with ice, which was exceptionally difficult to maintain on a chaotic, mud-soaked battlefield.

The Work of Harvey Cushing and Brain Surgery Head wounds caused by high-explosive shrapnel and sniper fire were among the most fatal injuries on the Western Front. At the beginning of the war, penetrating brain wounds carried a mortality rate of over 50%, as removing metal fragments often caused lethal brain swelling, haemorrhaging, or fatal cerebrospinal infections. Dr. Harvey Cushing, an American neurosurgeon working in British base hospitals and Casualty Clearing Stations, pioneered revolutionary surgical techniques for treating head trauma: 1. **Local Anaesthetic:** Cushing discovered that using general anaesthetics caused the brain to swell and bleed uncontrollably. By operating under local anaesthetic, brain pressure remained low, significantly reducing haemorrhage during delicate cranial procedures. 2. **Electromagnets and Suction:** Instead of probing deeply into brain tissue with forceps, which caused irreversible damage, Cushing used a surgical suction tube to clean damaged tissue and carefully drew out metal shrapnel fragments using a powerful electromagnet. 3. **Rapid Forward Surgery:** Cushing demonstrated that operating quickly before infection set in dramatically increased survival. At the Third Battle of Ypres (Passchendaele) in 1917, Cushing operated on 45 consecutive patients with penetrating head wounds, achieving a remarkable survival rate of 71% (reducing mortality to 29%).

Q6. Explain how Harvey Cushing reduced the mortality rate for brain surgery on the Western Front.



Source C: Source C: Walter Yeo's Facial Reconstruction by Harold Gillies (1917)

The Work of Harold Gillies and Plastic Surgery Modern industrial weapons, particularly exploding artillery shells, shrapnel, and machine-gun fire, caused catastrophic, disfiguring facial trauma that standard battlefield surgery could not repair. Soldiers surviving with shattered jaws, missing noses, or destroyed facial features suffered intense psychological trauma and social isolation. Dr. Harold Gillies, a New Zealand-born doctor serving in the Royal Army Medical Corps (RAMC), was horrified by the neglect of facial reconstruction and persuaded the army to establish a dedicated specialist hospital. In August 1917, the Queen's Hospital in Sidcup, Kent, was opened as the world's first specialized facial reconstructive surgery centre: 1. **The 'Tube Pedicle' Technique:** Gillies pioneered the 'waltzing tube pedicle' to transplant living skin. A flap of healthy skin from the chest, arm, or neck was cut and rolled into an enclosed tube, leaving one end attached to maintain its own blood supply while the free end was grafted onto the damaged facial area. Once new blood vessels developed, the remaining end was freed and shaped into a new nose, jaw, or cheek, preventing tissue death and infection. 2. **Detailed Medical Records:** Gillies worked closely with artists, photographers, and sculptors (such as Henry Tonks) to create meticulous anatomical drawings and wax models of each patient before, during, and after

reconstruction, establishing standardized plastic surgery methodologies. By the end of the war, Gillies and his surgical team at Sidcup had performed over 11,000 plastic surgery operations on more than 5,000 soldiers, laying the foundation for modern reconstructive and plastic surgery.

Q7. Describe the 'tube pedicle' technique pioneered by Harold Gillies.

Pair & Share Activity

Q8. Which medical advance on the Western Front represented a more impressive leap forward: the mechanical simplicity of the Thomas Splint which stabilized fractures, or the complex chemical engineering of blood preservation and storage?


Source A

From an official clinical report written by Captain Arthur Vaughan, a surgeon serving with the Royal Army Medical Corps (RAMC) at a Casualty Clearing Station (CCS) behind the frontline, October 1916.

"Our medical team at the Casualty Clearing Station has fully adopted the new leg traction splint. Before we received this iron frame, almost every compound femur fracture arrived at our ward in a state of terminal shock due to the jagged bones grating inside the flesh during transit. Since implementing this immobilization frame last month, we have kept the fractured limbs completely rigid. Out of forty admissions with thigh wounds, we have lost only six patients."

Source B

A studio photograph taken to train medical orderlies, showing a soldier's leg secured in a Thomas splint.



Provenance Scaffolding:

Source A is a clinical report by a surgeon; does this make its statistics reliable? Source B is a staged training photograph; while not taken on the chaotic frontline, does this actually make it better for demonstrating how the device technically worked?

Q9. 2 (a). How useful are Sources A and B for an enquiry into the effectiveness of new medical developments used to treat wounded soldiers on the Western Front? Explain your answer, using Sources A and B and your knowledge of the historical context. (8 marks)

