

# Lecture 22: Convex Optimization in Control With Powered Descent Guidance Example

Textbook Sections 4.3 & 12.7

Dr. Jordan D. Larson

# Introduction

- Optimal control problems: can be solved using generalized calculus of variations
  - Pontryagin's principle
  - Hamilton-Jacobi-Bellman equation

# Introduction

- Optimal control problems: can be solved using generalized calculus of variations
  - Pontryagin's principle
  - Hamilton-Jacobi-Bellman equation
- Linear optimal control problems:
  - Simpler solutions possible if convex optimization
  - Special convex optimization: semidefinite programming
  - Non-convex constraints may make solution more difficult

# Introduction

- Optimal control problems: can be solved using generalized calculus of variations
  - Pontryagin's principle
  - Hamilton-Jacobi-Bellman equation
- Linear optimal control problems:
  - Simpler solutions possible if convex optimization
  - Special convex optimization: semidefinite programming
  - Non-convex constraints may make solution more difficult
- Lecture: introduce concepts behind standard semidefinite programming solvers for optimal control

# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity

# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity
- To explain, consider two points in vector space  $\mathcal{V}$ , i.e.  $\vec{v}_1 \in \mathcal{V}$  and  $\vec{v}_2 \in \mathcal{V}$

# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity
- To explain, consider two points in vector space  $\mathcal{V}$ , i.e.  $\vec{v}_1 \in \mathcal{V}$  and  $\vec{v}_2 \in \mathcal{V}$
- Line segment  $L(\vec{v}_1, \vec{v}_2)$  between them contains set of points:

$$L(\vec{v}_1, \vec{v}_2) = \{\vec{v} \in \mathcal{V} : \vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2 \text{ for some } \mu \in [0, 1]\} \quad (1)$$

# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity
- To explain, consider two points in vector space  $\mathcal{V}$ , i.e.  $\vec{v}_1 \in \mathcal{V}$  and  $\vec{v}_2 \in \mathcal{V}$
- Line segment  $L(\vec{v}_1, \vec{v}_2)$  between them contains set of points:

$$L(\vec{v}_1, \vec{v}_2) = \{\vec{v} \in \mathcal{V} : \vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2 \text{ for some } \mu \in [0, 1]\} \quad (1)$$

- Let  $\mathcal{Q} \subset \mathcal{V}$  be nonempty, then  $\mathcal{Q}$  **convex set** of  $\mathcal{V}$  if for any  $\vec{v}_1, \vec{v}_2 \in \mathcal{Q}$ ,  $L(\vec{v}_1, \vec{v}_2) \subset \mathcal{Q}$



# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity
- To explain, consider two points in vector space  $\mathcal{V}$ , i.e.  $\vec{v}_1 \in \mathcal{V}$  and  $\vec{v}_2 \in \mathcal{V}$
- Line segment  $L(\vec{v}_1, \vec{v}_2)$  between them contains set of points:

$$L(\vec{v}_1, \vec{v}_2) = \{\vec{v} \in \mathcal{V} : \vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2 \text{ for some } \mu \in [0, 1]\} \quad (1)$$

- Let  $\mathcal{Q} \subset \mathcal{V}$  be nonempty, then  $\mathcal{Q}$  **convex set** of  $\mathcal{V}$  if for any  $\vec{v}_1, \vec{v}_2 \in \mathcal{Q}$ ,  $L(\vec{v}_1, \vec{v}_2) \subset \mathcal{Q}$
- Consider expression  $\vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2$  as weighed average, i.e. if  $\mu_1, \mu_2 \in [0, 1]$  with  $\mu_1 + \mu_2 = 1$ , then  $\vec{v} = \mu_1 \vec{v}_1 + \mu_2 \vec{v}_2$

# OCPs as Convex Optimizations

- Many OCPs can be set up convex optimization problems
  - Use concept of convexity
- To explain, consider two points in vector space  $\mathcal{V}$ , i.e.  $\vec{v}_1 \in \mathcal{V}$  and  $\vec{v}_2 \in \mathcal{V}$
- Line segment  $L(\vec{v}_1, \vec{v}_2)$  between them contains set of points:

$$L(\vec{v}_1, \vec{v}_2) = \{\vec{v} \in \mathcal{V} : \vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2 \text{ for some } \mu \in [0, 1]\} \quad (1)$$


- Let  $\mathcal{Q} \subset \mathcal{V}$  be nonempty, then  $\mathcal{Q}$  **convex set** of  $\mathcal{V}$  if for any  $\vec{v}_1, \vec{v}_2 \in \mathcal{Q}$ ,  $L(\vec{v}_1, \vec{v}_2) \subset \mathcal{Q}$
- Consider expression  $\vec{v} = \mu \vec{v}_1 + (1 - \mu) \vec{v}_2$  as weighed average, i.e. if  $\mu_1, \mu_2 \in [0, 1]$  with  $\mu_1 + \mu_2 = 1$ , then  $\vec{v} = \mu_1 \vec{v}_1 + \mu_2 \vec{v}_2$
- Generalized to average of  $n$  points as  $\vec{v} = \mu_1 \vec{v}_1 + \cdots + \mu_n \vec{v}_n$  with  $\mu_1, \dots, \mu_n \in [0, 1]$  and  $\mu_1 + \cdots + \mu_n = 1$

# Convex Set of Points

- Set comprised of all weighted averages of points  $\vec{v}_1, \dots, \vec{v}_n$ : convex hull,  $co(\{\vec{v}_1, \dots, \vec{v}_n\})$


# Convex Set of Points

- Set comprised of all weighted averages of points  $\vec{v}_1, \dots, \vec{v}_n$ : convex hull,  $\text{co}(\{\vec{v}_1, \dots, \vec{v}_n\})$
- Following diagram shows two-dimensional example of convex set defined by finite number of points & convex hull



# Convex Set of Points


- Set comprised of all weighted averages of points  $\vec{v}_1, \dots, \vec{v}_n$ : convex hull,  $co(\{\vec{v}_1, \dots, \vec{v}_n\})$
- Following diagram shows two-dimensional example of convex set defined by finite number of points & convex hull



- More generally, given set  $\mathcal{Q}$ ,  
**Convex hull**,  $co(\mathcal{Q})$ , by set  $\{\vec{v} \in \mathcal{V} : \text{there exists } n \text{ and } \vec{v}_1, \dots, \vec{v}_n \in \mathcal{Q} \text{ such that } \vec{v} \in co(\{\vec{v}_1, \dots, \vec{v}_n\})\}$ 
  - I.e. convex hull of  $\mathcal{Q}$ : collection of all possible weighted averages of points in  $\mathcal{Q}$


## 2D Example of Convex Hull

- Set  $\mathcal{Q}$



## 2D Example of Convex Hull

- Set  $\mathcal{Q}$




- Can show:

- Subset  $\mathcal{Q} \subset \text{co}(\mathcal{Q})$  satisfied
- Convex hull  $\text{co}(\mathcal{Q})$  convex
- $\text{co}(\mathcal{Q}) = \text{co}(\text{co}(\mathcal{Q}))$
- Set  $\mathcal{Q}$  convex if and only if  $\text{co}(\mathcal{Q}) = \mathcal{Q}$

## 2D Example of Convex Hull

- Set  $\mathcal{Q}$



- Can show:
  - Subset  $\mathcal{Q} \subset \text{co}(\mathcal{Q})$  satisfied
  - Convex hull  $\text{co}(\mathcal{Q})$  convex
  - $\text{co}(\mathcal{Q}) = \text{co}(\text{co}(\mathcal{Q}))$
  - Set  $\mathcal{Q}$  convex if and only if  $\text{co}(\mathcal{Q}) = \mathcal{Q}$
- Note: by definition, intersection of convex sets always convex



# Convex Cones

- Set  $\mathcal{Q} \subset \mathcal{V}$  called **cone** if closed under positive scalar multiplication, i.e. if

$$\vec{v} \in \mathcal{Q} \text{ implies } t\vec{v} \in \mathcal{Q} \text{ for every } t > 0 \quad (2)$$

# Convex Cones

- Set  $\mathcal{Q} \subset \mathcal{V}$  called **cone** if closed under positive scalar multiplication, i.e. if

$$\vec{v} \in \mathcal{Q} \text{ implies } t\vec{v} \in \mathcal{Q} \text{ for every } t > 0 \quad (2)$$

- Subspaces are cones, but cones broader set, e.g. half-line  $\mathcal{C}_v = \{\alpha \vec{v} : \alpha > 0\}$  for fixed vector  $\vec{v}$

# Convex Cones

- Set  $\mathcal{Q} \subset \mathcal{V}$  called **cone** if closed under positive scalar multiplication, i.e. if

$$\vec{v} \in \mathcal{Q} \text{ implies } t\vec{v} \in \mathcal{Q} \text{ for every } t > 0 \quad (2)$$


- Subspaces are cones, but cones broader set, e.g. half-line  $\mathcal{C}_v = \{\alpha \vec{v} : \alpha > 0\}$  for fixed vector  $\vec{v}$
- **Convex cone**: cones closed under addition, i.e.  $\vec{v}_1, \vec{v}_2 \in \mathcal{Q}$  implies  $\vec{v}_1 + \vec{v}_2 \in \mathcal{Q}$

# Convex Cones


- Set  $\mathcal{Q} \subset \mathcal{V}$  called **cone** if closed under positive scalar multiplication, i.e. if

$$\vec{v} \in \mathcal{Q} \text{ implies } t\vec{v} \in \mathcal{Q} \text{ for every } t > 0 \quad (2)$$

- Subspaces are cones, but cones broader set, e.g. half-line  $\mathcal{C}_v = \{\alpha \vec{v} : \alpha > 0\}$  for fixed vector  $\vec{v}$
- Convex cone**: cones closed under addition, i.e.  $\vec{v}_1, \vec{v}_2 \in \mathcal{Q}$  implies  $\vec{v}_1 + \vec{v}_2 \in \mathcal{Q}$
- Two-dimensional example of convex and nonconvex cone



Convex Cone



Nonconvex Cone

# Linear Programs (LP)



# Quadratic Programs (QP)



# Second-Order Cone Programs (SOCP)



# Semidefinite Programs (SDP)

- Many generalized OCPs formulated as **semidefinite programs (SDP)**: type of convex optimization
  - Use LMI constraints on solution space  $\mathcal{X}$ , e.g. ARIs



# Semidefinite Programs (SDP)

- Many generalized OCPs formulated as **semidefinite programs (SDP)**: type of convex optimization
  - Use LMI constraints on solution space  $\mathcal{X}$ , e.g. ARIs
- General form of SDP formulation:

$$\begin{aligned} X^{opt} = \operatorname{argmin}_{X \in \mathcal{X}} \quad & c(X) \\ \text{subject to: } \quad & F(X) \leq Q \\ & X \in \mathcal{X} \end{aligned} \tag{3}$$

- $c(X)$ : *linear* functional on vector space  $\mathcal{X}$
- A.k.a. type of **linear objective problem**

# Feasibility Problem

- For this SDP, LMI constraint must be feasible

# Feasibility Problem

- For this SDP, LMI constraint must be feasible
- Simplest SDP may be formulated as **feasibility problem**, if there exists  $X \in \mathcal{X}$  satisfying  $F(X) < Q$  also recast as linear objective problem

$$\begin{aligned} J &= \inf t \\ \text{subject to: } F(X) - tI &\leq Q \end{aligned} \tag{4}$$

- If and only if  $J < 0$ , then LMI  $F(X) < Q$  feasible

# Feasibility Problem

- For this SDP, LMI constraint must be feasible
- Simplest SDP may be formulated as **feasibility problem**, if there exists  $X \in \mathcal{X}$  satisfying  $F(X) < Q$  also recast as linear objective problem

$$\begin{aligned} J &= \inf t \\ \text{subject to: } F(X) - tI &\leq Q \end{aligned} \tag{4}$$

- If and only if  $J < 0$ , then LMI  $F(X) < Q$  feasible
- Feasibility question can be made part of SDP and focus of solving SDPs on solving linear objective problems with LMI constraints

# Important Convex Set

- To derive solutions for SDPs, note set  $\mathcal{C} = \{X \in \mathcal{X} : F(X) < Q\}$ : **convex set** in  $\mathcal{X}$

# Important Convex Set

- To derive solutions for SDPs, note set  $\mathcal{C} = \{X \in \mathcal{X} : F(X) < Q\}$ : **convex set** in  $\mathcal{X}$
- Proof: let  $X_1, X_2 \in \mathcal{C}$ , satisfies  $F(X_1) < Q$  and  $F(X_2) < Q$

# Important Convex Set

- To derive solutions for SDPs, note set  $\mathcal{C} = \{X \in \mathcal{X} : F(X) < Q\}$ : **convex set** in  $\mathcal{X}$
- Proof: let  $X_1, X_2 \in \mathcal{C}$ , satisfies  $F(X_1) < Q$  and  $F(X_2) < Q$
- Then, consider any point  $X_3 \in L(X_1, X_2)$ , i.e.  $X_3 = \mu X_1 + (1 - \mu)X_2$  for some value  $\mu \in [0, 1]$

# Important Convex Set

- To derive solutions for SDPs, note set  $\mathcal{C} = \{X \in \mathcal{X} : F(X) < Q\}$ : **convex set** in  $\mathcal{X}$
- Proof: let  $X_1, X_2 \in \mathcal{C}$ , satisfies  $F(X_1) < Q$  and  $F(X_2) < Q$
- Then, consider any point  $X_3 \in L(X_1, X_2)$ , i.e.  $X_3 = \mu X_1 + (1 - \mu)X_2$  for some value  $\mu \in [0, 1]$
- Using linearity of  $F()$ :

$$F(X_3) = \mu F(X_1) + (1 - \mu)F(X_2) < \mu Q + (1 - \mu)Q = Q \quad (5)$$

- Inequality follows from fact that positive definite matrices: convex cones
- Therefore:  $X_3 \in \mathcal{C}$



# Linear Objective Problem Example

- For  $X = [x_1 \ x_2]^T \in \mathbb{R}^2$  with

$$c(X) = Y^T X = y_1 x_1 + y_2 x_2 \quad (6)$$

- For some fixed  $Y \in \mathbb{R}^2$

# Linear Objective Problem Example

- For  $X = [x_1 \ x_2]^T \in \mathbb{R}^2$  with

$$c(X) = Y^T X = y_1 x_1 + y_2 x_2 \quad (6)$$

- For some fixed  $Y \in \mathbb{R}^2$
- Represent feasibility set  $\mathcal{C} = \{X \in \mathbb{R}^2 : F(X) < Q\}$  for linear objective problem:
  - $X_n$ : current guess
  - $X_{min}$ : element of  $\mathcal{C}$  with most negative projection in direction of  $Y$  and solution to problem

# Global Minimum for Convex Set

- Note: diagram correctly suggests no other local minima for function in set
  - Namely, for every other point exists “descent” direction
  - This property, fundamental to convex optimization problems, stated precisely as follows.

# Global Minimum for Convex Set

- Note: diagram correctly suggests no other local minima for function in set
  - Namely, for every other point exists “descent” direction
  - This property, fundamental to convex optimization problems, stated precisely as follows.
- If  $X_0$  *local* minimum of linear objective problem, i.e.  $c(X_0) \leq c(X)$  for every  $X$  in some neighborhood  $\mathcal{N}(X_0) \cap \mathcal{C}$ , then  $X_0$ : *global* minimum of problem over  $\mathcal{C}$

# Global Minimum for Convex Set

- Note: diagram correctly suggests no other local minima for function in set
  - Namely, for every other point exists “descent” direction
  - This property, fundamental to convex optimization problems, stated precisely as follows.
- If  $X_0$  *local* minimum of linear objective problem, i.e.  $c(X_0) \leq c(X)$  for every  $X$  in some neighborhood  $\mathcal{N}(X_0) \cap \mathcal{C}$ , then  $X_0$ : *global* minimum of problem over  $\mathcal{C}$
- Proof, let  $X_1$  be any other point in  $\mathcal{C}$

# Global Minimum for Convex Set

- Note: diagram correctly suggests no other local minima for function in set
  - Namely, for every other point exists “descent” direction
  - This property, fundamental to convex optimization problems, stated precisely as follows.
- If  $X_0$  *local* minimum of linear objective problem, i.e.  $c(X_0) \leq c(X)$  for every  $X$  in some neighborhood  $\mathcal{N}(X_0) \cap \mathcal{C}$ , then  $X_0$ : *global* minimum of problem over  $\mathcal{C}$
- Proof, let  $X_1$  be any other point in  $\mathcal{C}$
- Since  $\mathcal{C}$  convex: contains  $L(X_0, X_1)$  and by definition, neighborhood  $\mathcal{N}(X_0)$  contains points  $\mu X_1 + (1 - \mu)X_0$  for some sufficiently small  $\mu$ , i.e.  $\mu \in [0, \epsilon)$

# Global Minimum for Convex Set

- Note: diagram correctly suggests no other local minima for function in set
  - Namely, for every other point exists “descent” direction
  - This property, fundamental to convex optimization problems, stated precisely as follows.
- If  $X_0$  *local* minimum of linear objective problem, i.e.  $c(X_0) \leq c(X)$  for every  $X$  in some neighborhood  $\mathcal{N}(X_0) \cap \mathcal{C}$ , then  $X_0$ : *global* minimum of problem over  $\mathcal{C}$
- Proof, let  $X_1$  be any other point in  $\mathcal{C}$
- Since  $\mathcal{C}$  convex: contains  $L(X_0, X_1)$  and by definition, neighborhood  $\mathcal{N}(X_0)$  contains points  $\mu X_1 + (1 - \mu)X_0$  for some sufficiently small  $\mu$ , i.e.  $\mu \in [0, \epsilon)$
- Function  $f(\mu) = c(\mu X_1 + (1 - \mu)X_0)$  linear in  $\mu \in [0, \epsilon)$  and  $f(0) \leq f(\mu)$  for  $\mu \in [0, \epsilon)$  by hypothesis
  - $f(\mu)$  non-decreasing and  $f(0) \leq f(1)$ , or equivalently,  $c(X_0) \leq c(X_1)$

# Interior Point Method

- Property generalizes to any convex optimization problem and ensures these problems can be solved globally, not just locally, by numerical search algorithms



# Interior Point Method

- Property generalizes to any convex optimization problem and ensures these problems can be solved globally, not just locally, by numerical search algorithms
- At first glance, think minimum if it exists must lie on boundary of feasible set which would restrict search to boundary
  - Case for **linear programming**
  - Boundary in general SDPs complicated, thus, **interior point methods** favored in practice

# Interior Point Method

- Property generalizes to any convex optimization problem and ensures these problems can be solved globally, not just locally, by numerical search algorithms
- At first glance, think minimum if it exists must lie on boundary of feasible set which would restrict search to boundary
  - Case for **linear programming**
  - Boundary in general SDPs complicated, thus, **interior point methods** favored in practice
- Example of convex optimization method: suppose point  $X_n$  in feasible set

# Interior Point Method

- Property generalizes to any convex optimization problem and ensures these problems can be solved globally, not just locally, by numerical search algorithms
- At first glance, think minimum if it exists must lie on boundary of feasible set which would restrict search to boundary
  - Case for **linear programming**
  - Boundary in general SDPs complicated, thus, **interior point methods** favored in practice
- Example of convex optimization method: suppose point  $X_n$  in feasible set
- Immediate consequence: one only needs to keep set

$$\{X \in \mathcal{C} : c(X) \leq c(X_n)\} \quad (7)$$

- For remaining search for global minimum
- Amounts to setting  $\mathcal{C}$  with half-space

# Interior Point Method

- Property generalizes to any convex optimization problem and ensures these problems can be solved globally, not just locally, by numerical search algorithms
- At first glance, think minimum if it exists must lie on boundary of feasible set which would restrict search to boundary
  - Case for **linear programming**
  - Boundary in general SDPs complicated, thus, **interior point methods** favored in practice
- Example of convex optimization method: suppose point  $X_n$  in feasible set
- Immediate consequence: one only needs to keep set

$$\{X \in \mathcal{C} : c(X) \leq c(X_n)\} \quad (7)$$

- For remaining search for global minimum
  - Amounts to setting  $\mathcal{C}$  with half-space
- Can progressively shrink feasibility region to zero, provided able to successively generate “good” feasible point, e.g.  $X_{n+1}$

# Ellipsoid Algorithm

- Many optimization methods based on principle, one of simplest known as **ellipsoid algorithm**
  - Alternates between “cutting” and bounding resulting set by ellipsoid
  - $X_{n+1}$  would be center of such ellipsoid

# Ellipsoid Algorithm

- Many optimization methods based on principle, one of simplest known as **ellipsoid algorithm**
  - Alternates between “cutting” and bounding resulting set by ellipsoid
  - $X_{n+1}$  would be center of such ellipsoid
- More efficient methods for SDPs: based on **barrier functions** to impose feasibility constraint.
  - Idea: minimize function

$$c(X) + \alpha\phi(X) \tag{8}$$

- Where  $\alpha > 0$  and barrier function  $\phi(X)$  convex and approaches infinity on boundary of feasible set
- E.g. for set  $\mathcal{C}$

$$\phi(X) = -\log(\det[Q - F(X)]) \tag{9}$$

Serve as barrier function

# Convex Optimization Algorithms

- Provided one starts from feasible point, minimization can be globally solved by unconstrained optimization methods, e.g. Newton's method

# Convex Optimization Algorithms

- Provided one starts from feasible point, minimization can be globally solved by unconstrained optimization methods, e.g. Newton's method
- By successively reducing weight of barrier function: iteration produced converges to global minimum
  - Computational complexity of polynomial growth with problem size characterized by dimension of  $\mathcal{X}$  and constraint set  $\mathbb{H}^n$



# Convex Optimization Algorithms

- Provided one starts from feasible point, minimization can be globally solved by unconstrained optimization methods, e.g. Newton's method
- By successively reducing weight of barrier function: iteration produced converges to global minimum
  - Computational complexity of polynomial growth with problem size characterized by dimension of  $\mathcal{X}$  and constraint set  $\mathbb{H}^n$
- Additional details on convex optimization algorithms beyond scope of textbook and left to reader

# Algebraic Riccati Equation

- *Will be shown:* solutions to infinite-horizon OCPs found by setting up algebraic Riccati equations (ARE) of general form:

$$A^T P + PA + Q + PRP = 0 \quad (10)$$

- Typically, required to find particular solution  $P = P^T \in \mathbb{R}^{n_x \times n_x}$  such that  $A + RP$  stable, i.e. all its eigenvalues have strict negative real part

# Algebraic Riccati Equation

- *Will be shown:* solutions to infinite-horizon OCPs found by setting up algebraic Riccati equations (ARE) of general form:

$$A^T P + PA + Q + PRP = 0 \quad (10)$$

- Typically, required to find particular solution  $P = P^T \in \mathbb{R}^{n_x \times n_x}$  such that  $A + RP$  stable, i.e. all its eigenvalues have strict negative real part
- For such problems, define Hamiltonian matrix of ARE of  $(A, Q, R)$ :

$$H = \begin{bmatrix} A & R \\ -Q & -A^T \end{bmatrix} \quad (11)$$

# Algebraic Riccati Equation (continued)

- Alternate ARE:

$$\begin{bmatrix} P & -I \end{bmatrix} H \begin{bmatrix} I \\ P \end{bmatrix} = 0 \quad (12)$$

- Eigenvalue decomposition of  $H$ : key to solving ARE

# Algebraic Riccati Equation (continued)

- Alternate ARE:

$$\begin{bmatrix} P & -I \end{bmatrix} H \begin{bmatrix} I \\ P \end{bmatrix} = 0 \quad (12)$$

- Eigenvalue decomposition of  $H$ : key to solving ARE
- In this case, state if:
  - $A$ ,  $Q = Q^T$  and  $R = R^T$  given
  - $H$  has no purely imaginary axis eigenvalues
  - $R \geq 0$  or  $R \leq 0$
  - $(A, R)$  stabilizable

Then ARE of  $(A, Q, R)$  has unique solution  $P = P^T$  such that  $A + RP$  stable

# Algebraic Riccati Inequality

- For more advanced control synthesis, instead use **algebraic Riccati inequality (ARI)**:

$$A^T P + PA + Q + PRP < 0 \quad (13)$$

- Type of linear matrix inequality discussed in following section.

# Algebraic Riccati Inequality

- For more advanced control synthesis, instead use **algebraic Riccati inequality (ARI)**:

$$A^T P + PA + Q + PRP < 0 \quad (13)$$

- Type of linear matrix inequality discussed in following section.
- For example, consider inequality

$$A^T P + PA + C^T C + \gamma^{-2} P B B^T P < 0 \quad (14)$$

- Quadratic in  $P \in \mathbb{S}^n > 0$

# Algebraic Riccati Inequality

- Rewriting this as

$$\begin{bmatrix} A^T P + PA + C^T C + \gamma^{-2} P B B^T P & 0 \\ 0 & -\gamma^{-2} I \end{bmatrix} < 0 \quad (15)$$



# Algebraic Riccati Inequality

- Rewriting this as

$$\begin{bmatrix} A^T P + PA + C^T C + \gamma^{-2} P B B^T P & 0 \\ 0 & -\gamma^{-2} I \end{bmatrix} < 0 \quad (15)$$

- By Schur Complement Lemma, form equivalent LMI for  $P$

$$\begin{bmatrix} A^T P + PA + C^T C & PB \\ B^T P & -\gamma^2 I \end{bmatrix} < 0 \quad (16)$$

# Bounded Real Lemma

$$\begin{bmatrix} A^T P + PA & PB \\ B^T P & \gamma^{-2} I \end{bmatrix} < \begin{bmatrix} -C^T C & 0 \\ 0 & 0 \end{bmatrix} \quad (17)$$

- LMI in  $P$  as left side can be assigned as  $F(P) : \mathbb{S}^n \rightarrow \mathbb{S}^n$
- Right side assigned as  $Q \in \mathbb{S}$

# Bounded Real Lemma

$$\begin{bmatrix} A^T P + PA & PB \\ B^T P & \gamma^{-2} I \end{bmatrix} < \begin{bmatrix} -C^T C & 0 \\ 0 & 0 \end{bmatrix} \quad (17)$$

- LMI in  $P$  as left side can be assigned as  $F(P) : \mathbb{S}^n \rightarrow \mathbb{S}^n$
- Right side assigned as  $Q \in \mathbb{S}$
- In this case, relate solutions to AREs and ARIs for OCPs

# Bounded Real Lemma

$$\begin{bmatrix} A^T P + PA & PB \\ B^T P & \gamma^{-2} I \end{bmatrix} < \begin{bmatrix} -C^T C & 0 \\ 0 & 0 \end{bmatrix} \quad (17)$$

- LMI in  $P$  as left side can be assigned as  $F(P) : \mathbb{S}^n \rightarrow \mathbb{S}^n$
- Right side assigned as  $Q \in \mathbb{S}$
- In this case, relate solutions to AREs and ARIs for OCPs
- One of important lemmas between AREs/ARIs: known as **Bounded Real Lemma**, a.k.a. **Kalman-Yacubovich-Popov (KYP) lemma** stated as follows

# Bounded Real Lemma

$$\begin{bmatrix} A^T P + PA & PB \\ B^T P & \gamma^{-2} I \end{bmatrix} < \begin{bmatrix} -C^T C & 0 \\ 0 & 0 \end{bmatrix} \quad (17)$$

- LMI in  $P$  as left side can be assigned as  $F(P) : \mathbb{S}^n \rightarrow \mathbb{S}^n$
- Right side assigned as  $Q \in \mathbb{S}$
- In this case, relate solutions to AREs and ARIs for OCPs
- One of important lemmas between AREs/ARIs: known as **Bounded Real Lemma**, a.k.a. **Kalman-Yacubovich-Popov (KYP) lemma** stated as follows
- Consider following LTI system,  $F_L(G, K)$ :

$$\begin{aligned} \dot{\vec{x}}(t) &= A_L \vec{x}(t) + B_1 \vec{d}(t) \\ \vec{e}(t) &= C_1 \vec{x}(t) \end{aligned} \quad (18)$$

- Let  $\gamma > 0$  be given

## Bounded Real Lemma (continued)

- Following three statements equivalent

①  $F_L(G, K)$  stable, i.e.  $A_L$  stable, and  $\|F_L(G, K)\|_\infty < \gamma^2$

② Exists unique  $P_1 \geq 0$  such that  $A_L + \gamma^{-2}P_1B_1B_1^TP_1$  stable and satisfies ARE:

$$A^TP_1 + P_1A + C_1^TC_1 + \gamma^{-2}P_1B_1B_1^TP_1 = 0 \quad (19)$$

③ Exists  $P_2 > 0$  satisfying strict ARI:

$$A^TP_2 + P_2A + C_1^TC_1 + \gamma^{-2}P_2B_1B_1^TP_2 < 0 \quad (20)$$

# Bounded Real Lemma (continued)

- Following three statements equivalent

①  $F_L(G, K)$  stable, i.e.  $A_L$  stable, and  $\|F_L(G, K)\|_\infty < \gamma^2$

② Exists unique  $P_1 \geq 0$  such that  $A_L + \gamma^{-2}P_1B_1B_1^TP_1$  stable and satisfies ARE:


$$A^TP_1 + P_1A + C_1^TC_1 + \gamma^{-2}P_1B_1B_1^TP_1 = 0 \quad (19)$$

③ Exists  $P_2 > 0$  satisfying strict ARI:

$$A^TP_2 + P_2A + C_1^TC_1 + \gamma^{-2}P_2B_1B_1^TP_2 < 0 \quad (20)$$

- To solve ARIs for  $P_2$  requires **semidefinite programming (SDP)**
  - Class of convex optimization
  - Discussed in later subsection and then applied to both  $\mathcal{H}_2$  and  $\mathcal{H}_\infty$  OCPs in later lectures

# Generalized Feedback Control System



- Generalized plant,  $G(s)$ :

$$\begin{aligned}\dot{\vec{x}}(t) &= A\vec{x}(t) + B_1\vec{d}(t) + B_2\vec{u}(t) \\ \vec{e}(t) &= C_1\vec{x}(t) + D_{11}\vec{d}(t) + D_{12}\vec{u}(t)\end{aligned}\tag{21}$$

- State feedback control policy for  $K(s)$ :

$$\vec{u}(t) = D_K\vec{x}(t)\tag{22}$$



# Generalized Feedback Control System (continued)

- Results in closed-loop system:

$$\begin{aligned}\dot{\vec{x}}(t) &= (A + B_2 D_K) \vec{x}(t) + B_1 \vec{d}(t) \\ \vec{e}(t) &= (C_1 + D_{12} D_K) \vec{x}(t) + D_{11} \vec{d}(t)\end{aligned}\tag{23}$$

- Stable if and only if  $A_L = A + B_2 D_K$  stable

## Generalized Feedback Control System (continued)

- Results in closed-loop system:

$$\begin{aligned}\dot{\vec{x}}(t) &= (A + B_2 D_K) \vec{x}(t) + B_1 \vec{d}(t) \\ \vec{e}(t) &= (C_1 + D_{12} D_K) \vec{x}(t) + D_{11} \vec{d}(t)\end{aligned}\tag{23}$$

- Stable if and only if  $A_L = A + B_2 D_K$  stable
- LMI for stabilizing  $D_K$ , enforce stable condition by matrix Lyapunov inequality:

$$A_L P + P A_L^T < 0\tag{24}$$

- Solution  $P > 0$

## Generalized Feedback Control System (continued)

- Results in closed-loop system:

$$\begin{aligned}\dot{\vec{x}}(t) &= (A + B_2 D_K) \vec{x}(t) + B_1 \vec{d}(t) \\ \vec{e}(t) &= (C_1 + D_{12} D_K) \vec{x}(t) + D_{11} \vec{d}(t)\end{aligned}\tag{23}$$

- Stable if and only if  $A_L = A + B_2 D_K$  stable
- LMI for stabilizing  $D_K$ , enforce stable condition by matrix Lyapunov inequality:

$$A_L P + P A_L^T < 0\tag{24}$$

- Solution  $P > 0$
- Substituting for  $A_L$ :

$$(A + B_2 D_K) P + P (A + B_2 D_K)^T < 0\tag{25}$$

# LMI Construction

- Expanded:

$$AP + PA^T + B_2(D_K P) + (PD_K^T)B_2^T < 0 \quad (26)$$

# LMI Construction

- Expanded:

$$AP + PA^T + B_2(D_K P) + (PD_K^T)B_2^T < 0 \quad (26)$$

- Substituting  $Y = D_K P$ , then LMI in  $P$  and  $Y$ :

$$\begin{bmatrix} A & B_2 \end{bmatrix} \begin{bmatrix} P \\ Y \end{bmatrix} - \begin{bmatrix} P & Y^T \end{bmatrix} \begin{bmatrix} A^T \\ B_2^T \end{bmatrix} < 0 \quad (27)$$

- Solved for  $P$  and  $Y$  and obtain stabilizing feedback controller as  $D_K = YP^{-1}$

# LMI Construction

- Expanded:

$$AP + PA^T + B_2(D_K P) + (PD_K^T)B_2^T < 0 \quad (26)$$

- Substituting  $Y = D_K P$ , then LMI in  $P$  and  $Y$ :

$$\begin{bmatrix} A & B_2 \end{bmatrix} \begin{bmatrix} P \\ Y \end{bmatrix} - \begin{bmatrix} P & Y^T \end{bmatrix} \begin{bmatrix} A^T \\ B_2^T \end{bmatrix} < 0 \quad (27)$$

- Solved for  $P$  and  $Y$  and obtain stabilizing feedback controller as  $D_K = YP^{-1}$
- Note: process can be expanded to general LMI characterizations for non-state feedback controllers  $D_K$

# Summary

- Linear optimal control problems:
  - Finite-horizon, quadratic cost: Riccati differential equation
  - Infinite-horizon, quadratic-like costs: algebraic Riccati equation & algebraic Riccati inequality
  - Inequalities for stability and constraints: semidefinite programming

# Summary

- Linear optimal control problems:
  - Finite-horizon, quadratic cost: Riccati differential equation
  - Infinite-horizon, quadratic-like costs: algebraic Riccati equation & algebraic Riccati inequality
  - Inequalities for stability and constraints: semidefinite programming
- Convex optimal control problems:
  - Interior point methods: search iteratively for minimum in convex set
  - Local minimum = global minimum



# Summary

- Linear optimal control problems:
  - Finite-horizon, quadratic cost: Riccati differential equation
  - Infinite-horizon, quadratic-like costs: algebraic Riccati equation & algebraic Riccati inequality
  - Inequalities for stability and constraints: semidefinite programming
- Convex optimal control problems:
  - Interior point methods: search iteratively for minimum in convex set
  - Local minimum = global minimum
- Bounded real lemma: stabilize and bound  $\mathcal{H}_\infty$ -norm for generalized feedback control system

# Summary

- Linear optimal control problems:
  - Finite-horizon, quadratic cost: Riccati differential equation
  - Infinite-horizon, quadratic-like costs: algebraic Riccati equation & algebraic Riccati inequality
  - Inequalities for stability and constraints: semidefinite programming
- Convex optimal control problems:
  - Interior point methods: search iteratively for minimum in convex set
  - Local minimum = global minimum
- Bounded real lemma: stabilize and bound  $\mathcal{H}_\infty$ -norm for generalized feedback control system
- Simple LMI for stabilizing state feedback control