

# Lecture 24: Time-Varying Systems Theory and Control

## Textbook Section 6.1 & 6.2

Dr. Jordan D. Larson

# Introduction

- Course to this point: used LTI systems theory for system analysis and control design

# Introduction

- Course to this point: used LTI systems theory for system analysis and control design
- Flight vehicles inherently nonlinear systems
  - Linearization depends on many flight conditions

# Introduction

- Course to this point: used LTI systems theory for system analysis and control design
- Flight vehicles inherently nonlinear systems
  - Linearization depends on many flight conditions
- Another approach: linear, parameter-varying (LPV) systems
  - Utilize linearization dependence on non-state flight conditions, i.e. parameters
  - Requires additional dynamical systems theory

# Introduction

- Course to this point: used LTI systems theory for system analysis and control design
- Flight vehicles inherently nonlinear systems
  - Linearization depends on many flight conditions
- Another approach: linear, parameter-varying (LPV) systems
  - Utilize linearization dependence on non-state flight conditions, i.e. parameters
  - Requires additional dynamical systems theory
- Introduces concepts used in nonlinear systems as well

# Time-Varying Dynamical System

- **Time-varying dynamical system:**

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}, \vec{u}) \\ \vec{y}(t) &= h(t, \vec{x}, \vec{u})\end{aligned}\tag{1}$$

# Time-Varying Dynamical System

- **Time-varying dynamical system:**

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}, \vec{u}) \\ \vec{y}(t) &= h(t, \vec{x}, \vec{u})\end{aligned}\tag{1}$$

- Unforced, i.e.

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}) \\ \vec{y}(t) &= h(t, \vec{x})\end{aligned}\tag{2}$$

# Time-Varying Dynamical System

- **Time-varying dynamical system:**

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}, \vec{u}) \\ \vec{y}(t) &= h(t, \vec{x}, \vec{u})\end{aligned}\tag{1}$$

- Unforced, i.e.

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}) \\ \vec{y}(t) &= h(t, \vec{x})\end{aligned}\tag{2}$$

- **Initial value problem (IVP):** system at time  $t_0 \geq 0$  as some  $\vec{x}(t_0) = \vec{x}_0 \in \mathbb{R}^n$ 
  - A.k.a. **Cauchy problem**
  - May have many solutions, one unique solution, or no existing solution



# Time-Varying Dynamical System

- **Time-varying dynamical system:**

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}, \vec{u}) \\ \vec{y}(t) &= h(t, \vec{x}, \vec{u})\end{aligned}\tag{1}$$

- Unforced, i.e.

$$\begin{aligned}\dot{\vec{x}}(t) &= f(t, \vec{x}) \\ \vec{y}(t) &= h(t, \vec{x})\end{aligned}\tag{2}$$

- **Initial value problem (IVP):** system at time  $t_0 \geq 0$  as some  $\vec{x}(t_0) = \vec{x}_0 \in \mathbb{R}^n$ 
  - A.k.a. **Cauchy problem**
  - May have many solutions, one unique solution, or no existing solution
- Existence and uniqueness of solutions to IVPs for non-LTI dynamical systems not always guaranteed

# Existence and Uniqueness of Solutions

- **Cauchy-Peano Theorem:** sufficient conditions for IVP to admit solution, may not be unique:

If, for some  $T > 0$ , some  $\epsilon > 0$ , and  $f(t, \vec{x}) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  continuous in closed region, i.e.

$$B = \{(t, \vec{x}) : |t - t_0| \leq T, \|\vec{x} - \vec{x}_0\| \leq \epsilon\} \subseteq \mathbb{R} \times \mathbb{R}^n \quad (3)$$

Then, exists  $t_0 < t_1 \leq T$  s.t. IVP has at least one continuously differentiable solution  $\vec{x}(t)$  on interval  $[t_0, T]$

- Assumed continuity of  $f(t, \vec{x})$  in arguments,  $t$  and  $\vec{x}$  ensure at least one solution of IVP
- Does not guarantee uniqueness of solution

# Existence and Uniqueness of Solutions

- **Cauchy-Peano Theorem:** sufficient conditions for IVP to admit solution, may not be unique:  
If, for some  $T > 0$ , some  $\epsilon > 0$ , and  $f(t, \vec{x}) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  continuous in closed region, i.e.

$$B = \{(t, \vec{x}) : |t - t_0| \leq T, \|\vec{x} - \vec{x}_0\| \leq \epsilon\} \subseteq \mathbb{R} \times \mathbb{R}^n \quad (3)$$

Then, exists  $t_0 < t_1 \leq T$  s.t. IVP has at least one continuously differentiable solution  $\vec{x}(t)$  on interval  $[t_0, T]$

- Assumed continuity of  $f(t, \vec{x})$  in arguments,  $t$  and  $\vec{x}$  ensure at least one solution of IVP
- Does not guarantee uniqueness of solution
- Key constraint that yields uniqueness: **Lipschitz condition** whereby  $f(t, \vec{x})$  satisfies inequality

$$\|f(t, \vec{x}) - f(t, \vec{y})\| \leq L \|\vec{x} - \vec{y}\| \quad (4)$$

for all  $(t, \vec{x})$  and  $(t, \vec{y})$  in some “neighborhood” of  $(t_0, \vec{x}_0)$  with finite constant  $L > 0$

## Existence and Uniqueness of Solutions (continued)

- Sufficient conditions for IVP to admit *local* existence and uniqueness of solution:  
If, for some  $\epsilon > 0$ ,  $f(t, \vec{x}) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$  and satisfies Lipschitz condition, i.e.

$$\forall \vec{x}, \vec{y} \in B = \{\vec{x} \in \mathbb{R}^n : \|\vec{x} - \vec{x}_0\| \leq \epsilon\}, \forall t \in [t_0, t_1] \quad (5)$$

then, exists some  $\delta > 0$  such that IVP for state equation  $\dot{\vec{x}} = f(t, \vec{x})$  with  $\vec{x}(t_0) = \vec{x}_0$  has unique solution over  $[t_0, t_0 + \delta]$ .

# Existence and Uniqueness of Solutions (continued)

- Sufficient conditions for IVP to admit *local* existence and uniqueness of solution:  
If, for some  $\epsilon > 0$ ,  $f(t, \vec{x}) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$  and satisfies Lipschitz condition, i.e.

$$\forall \vec{x}, \vec{y} \in B = \{\vec{x} \in \mathbb{R}^n : \|\vec{x} - \vec{x}_0\| \leq \epsilon\}, \forall t \in [t_0, t_1] \quad (5)$$

then, exists some  $\delta > 0$  such that IVP for state equation  $\dot{\vec{x}} = f(t, \vec{x})$  with  $\vec{x}(t_0) = \vec{x}_0$  has unique solution over  $[t_0, t_0 + \delta]$ .

- Sufficient conditions for IVP to admit *global* existence and uniqueness of solution:  
If, for some finite  $L > 0$  and  $f(t, \vec{x}) : \mathbb{R} \times \mathbb{R}^n \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$  and globally Lipschitz in  $\vec{x}$ , i.e.

$$\|f(t, \vec{x}) - f(t, \vec{y})\| \leq L\|\vec{x} - \vec{y}\|, \quad \forall \vec{x}, \vec{y} \in \mathbb{R}^n, \quad \forall t \in [t_0, t_1] \quad (6)$$

then, IVP has unique solution over  $[t_0, t_1]$  where final time  $t_1$  may be arbitrarily large

# Equilibrium for Unforced Time-Varying System

- Unforced, time-varying dynamics:  $f : [0, \infty) \times D \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$

# Equilibrium for Unforced Time-Varying System

- Unforced, time-varying dynamics:  $f : [0, \infty) \times D \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$
- Locally Lipschitz in  $\vec{x}$  with domain  $D \subset \mathbb{R}^n$  that contains origin  $\vec{x} = \vec{0}$

## Equilibrium for Unforced Time-Varying System

- Unforced, time-varying dynamics:  $f : [0, \infty) \times D \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$
- Locally Lipschitz in  $\vec{x}$  with domain  $D \subset \mathbb{R}^n$  that contains origin  $\vec{x} = \vec{0}$
- Origin  $\mathbb{R}^n$ : **equilibrium point** for unforced, time-varying system at  $t_0 = 0$  if

$$f(t, \vec{0}) = \vec{0}, \quad \forall t \geq 0 \quad (7)$$



# Equilibrium for Unforced Time-Varying System

- Unforced, time-varying dynamics:  $f : [0, \infty) \times D \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$
- Locally Lipschitz in  $\vec{x}$  with domain  $D \subset \mathbb{R}^n$  that contains origin  $\vec{x} = \vec{0}$
- Origin  $\mathbb{R}^n$ : **equilibrium point** for unforced, time-varying system at  $t_0 = 0$  if

$$f(t, \vec{0}) = \vec{0}, \quad \forall t \geq 0 \quad (7)$$

- Without loss of generality, extended beyond origin and initial time by defining nonzero vector  $\vec{x} \in \mathbb{R}^n$  to be equilibrium point of  $\dot{\vec{x}} = f(t, \vec{x})$  at nonzero initial time  $t = t_0$ , thus

$$f(t, \vec{x}) = \vec{0}, \quad \forall t \geq t_0 \quad (8)$$

# Equilibrium for Unforced Time-Varying System

- Unforced, time-varying dynamics:  $f : [0, \infty) \times D \rightarrow \mathbb{R}^n$  piece-wise continuous in  $t$
- Locally Lipschitz in  $\vec{x}$  with domain  $D \subset \mathbb{R}^n$  that contains origin  $\vec{x} = \vec{0}$
- Origin  $\mathbb{R}^n$ : **equilibrium point** for unforced, time-varying system at  $t_0 = 0$  if

$$f(t, \vec{0}) = \vec{0}, \quad \forall t \geq 0 \quad (7)$$

- Without loss of generality, extended beyond origin and initial time by defining nonzero vector  $\vec{x} \in \mathbb{R}^n$  to be equilibrium point of  $\dot{\vec{x}} = f(t, \vec{x})$  at nonzero initial time  $t = t_0$ , thus

$$f(t, \vec{x}) = \vec{0}, \quad \forall t \geq t_0 \quad (8)$$

- Define: new time as  $\tau = t - t_0$  and new state as  $\vec{z}(\tau) = \vec{x}(\tau + t_0) - \vec{x}$

$$\frac{d\vec{z}(\tau)}{d\tau} = \frac{d\vec{x}(\tau + t_0)}{dt} = f(\tau + t_0, \vec{z}(\tau) + \vec{x}) = g(\tau, \vec{z}(\tau)) \quad (9)$$

- With  $g(0, \vec{0}) = f(t_0, \vec{x}) = 0$
- Can shift equilibrium point to origin and initial time to zero

## Zero and Origin as Arbitrary Equilibrium

- Suppose one has state trajectory  $\vec{x}(t)$  that starts at  $t = t_0$ , i.e.

$$\dot{\vec{x}}^*(t) = f(t, \vec{x}(t)), \quad \forall t \geq t_0 \quad (10)$$

# Zero and Origin as Arbitrary Equilibrium

- Suppose one has state trajectory  $\vec{x}(t)$  that starts at  $t = t_0$ , i.e.

$$\dot{\vec{x}}^*(t) = f(t, \vec{x}(t)), \quad \forall t \geq t_0 \quad (10)$$

- Define new time as  $\tau = t - t_0$  and new state this time as  $\vec{z}(\tau) = \vec{x}(\tau + t_0) - \vec{x}(t_0)$ :

$$\begin{aligned} \frac{d\vec{z}(\tau)}{d\tau} &= \frac{d\vec{x}(\tau + t_0)}{d\tau} - \frac{d\vec{x}(t_0)}{d\tau} \\ &= f(\tau + t_0, \vec{z}(\tau) + \vec{x}(t_0)) - f(t_0, \vec{x}(t_0)) = g(\tau, \vec{z}(\tau)) \end{aligned} \quad (11)$$

- With  $g(0, \vec{0}) = \vec{0}$

# Zero and Origin as Arbitrary Equilibrium

- Suppose one has state trajectory  $\vec{x}(t)$  that starts at  $t = t_0$ , i.e.

$$\dot{\vec{x}}^*(t) = f(t, \vec{x}(t)), \quad \forall t \geq t_0 \quad (10)$$

- Define new time as  $\tau = t - t_0$  and new state this time as  $\vec{z}(\tau) = \vec{x}(\tau + t_0) - \vec{x}^*(\tau + t_0)$ :

$$\begin{aligned} \frac{d\vec{z}(\tau)}{d\tau} &= \frac{d\vec{x}(\tau + t_0)}{dt} - \frac{d\vec{x}^*(\tau + t_0)}{dt} \\ &= f(\tau + t_0, \vec{z}(\tau) + \vec{x}^*(\tau + t_0)) - f(\tau + t_0, \vec{x}^*(\tau + t_0)) = g(\tau, \vec{z}(\tau)) \end{aligned} \quad (11)$$

- With  $g(0, \vec{0}) = \vec{0}$
- Analyzing these new dynamics around origin as equilibrium point while starting at  $t_0$ : determine original system behavior around original nonzero equilibrium  $\vec{x}^*$ 
  - I.e. can assess system relative dynamics w.r.t. any time-dependent trajectory  $\vec{x}^*(t)$ , starting at arbitrary initial time  $t_0 \geq 0$
  - Key for stability analysis

# Lyapunov Stability

- Lyapunov stability theory provides analysis tool for assessing behavior of system trajectories near equilibrium points, but *without* explicit computation of these trajectories

# Lyapunov Stability

- Lyapunov stability theory provides analysis tool for assessing behavior of system trajectories near equilibrium points, but *without* explicit computation of these trajectories
- System stability can be interpreted as continuity of system trajectories, w.r.t. initial conditions, over *infinite* time interval

# Lyapunov Stability

- Lyapunov stability theory provides analysis tool for assessing behavior of system trajectories near equilibrium points, but *without* explicit computation of these trajectories
- System stability can be interpreted as continuity of system trajectories, w.r.t. initial conditions, over *infinite* time interval
- Infinite time interval highlights primary notion of stability as continuity property of Lipschitz-continuous differential equations holding infinitely in time
  - Let  $\vec{x}(t, \vec{x}_0)$  define unique solution of  $\dot{\vec{x}} = f(t, \vec{x})$  with initial condition  $\vec{x}(t_0) = \vec{x}_0$  which exists on finite, possibly open-ended interval  $[t_0, T)$
- Continuity property of  $\vec{x}(t, \vec{x}_0)$  due to changes in  $\vec{x}_0$ :
  - Given any constant  $\epsilon > 0$ , there must exist sufficiently small constant  $\delta > 0$  such that for all perturbed initial conditions  $\vec{x}_0 + \Delta \vec{x}_0$  with  $\|\Delta \vec{x}_0\| \leq \delta$   
Corresponding perturbed solution  $\vec{x}(t, \vec{x}_0 + \Delta \vec{x}_0)$  deviates from original  $\vec{x}_0$  by no more than  $\epsilon$ 
    - I.e.  $\|\vec{x}(t, \vec{x}_0 + \Delta \vec{x}_0) - \vec{x}(t, \vec{x}_0)\| \leq \epsilon$ , for all  $t_0 \leq t < T$



# Lyapunov Stability of Equilibrium Points

- **Lyapunov stability of equilibrium point**,  $\vec{x} = \vec{0}$ , i.e. origin, for time-varying unforced dynamics:
  - *Stable*: if for any  $\epsilon > 0$  and  $t \geq 0$  there exists some  $\delta(\epsilon, t_0) > 0$  such that for all initial conditions  $\|\vec{x}_0\| < \delta$  and for all  $t \geq t_0 \geq 0$ , the corresponding system trajectories bounded, i.e.  $\|\vec{x}(t)\| < \epsilon$
  - Otherwise, *unstable*
  - Given outer “hyper-sphere”  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ , can find inner “hyper-sphere”  $B_\delta = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \delta\}$ , such that any trajectory that starts inside  $B_\delta$  will evolve inside  $B_\epsilon$  for *all* future times

# Lyapunov Stability of Equilibrium Points

- **Lyapunov stability of equilibrium point**,  $\vec{x} = \vec{0}$ , i.e. origin, for time-varying unforced dynamics:
  - *Stable*: if for any  $\epsilon > 0$  and  $t \geq 0$  there exists some  $\delta(\epsilon, t_0) > 0$  such that for all initial conditions  $\|\vec{x}_0\| < \delta$  and for all  $t \geq t_0 \geq 0$ , the corresponding system trajectories bounded, i.e.  $\|\vec{x}(t)\| < \epsilon$
  - Otherwise, *unstable*
  - Given outer “hyper-sphere”  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ , can find inner “hyper-sphere”  $B_\delta = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \delta\}$ , such that any trajectory that starts inside  $B_\delta$  will evolve inside  $B_\epsilon$  for *all* future times
- Nonlinear dynamical systems may display completely different behavior in various domains of state: distinguish between local and global stability
  - Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **global stability** if stable and  $\lim_{\epsilon \rightarrow \infty} \delta(\epsilon, t_0) = \infty$ 
    - I.e. trajectory will not deviate “too far” from any arbitrary initial conditions

# Lyapunov Stability of Equilibrium Points

- **Lyapunov stability of equilibrium point**,  $\vec{x} = \vec{0}$ , i.e. origin, for time-varying unforced dynamics:
  - *Stable*: if for any  $\epsilon > 0$  and  $t \geq 0$  there exists some  $\delta(\epsilon, t_0) > 0$  such that for all initial conditions  $\|\vec{x}_0\| < \delta$  and for all  $t \geq t_0 \geq 0$ , the corresponding system trajectories bounded, i.e.  $\|\vec{x}(t)\| < \epsilon$
  - Otherwise, *unstable*
  - Given outer “hyper-sphere”  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ , can find inner “hyper-sphere”  $B_\delta = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \delta\}$ , such that any trajectory that starts inside  $B_\delta$  will evolve inside  $B_\epsilon$  for *all* future times
- Nonlinear dynamical systems may display completely different behavior in various domains of state: distinguish between local and global stability
  - Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **global stability** if stable and  $\lim_{\epsilon \rightarrow \infty} \delta(\epsilon, t_0) = \infty$ 
    - I.e. trajectory will not deviate “too far” from any arbitrary initial conditions
- System trajectories of time-varying dynamical systems depend on initial time  $t_0$ 
  - Stability of equilibrium point for time-varying systems may depend on  $t_0$
  - Equilibrium point has **uniform stability** if stable and  $\delta$  does not depend on  $t_0$

# Stricter Lyapunov Stability of Equilibrium Points

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **asymptotic stability**:

If stable and there exists constant  $c = c(t_0) > 0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$

# Stricter Lyapunov Stability of Equilibrium Points

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **asymptotic stability**:

If stable and there exists constant  $c = c(t_0) > 0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **uniform asymptotic stability**:

If uniformly stable and there exists constant  $c > 0$  independent of  $t_0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$  uniformly in  $t_0$

# Stricter Lyapunov Stability of Equilibrium Points

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **asymptotic stability**:

If stable and there exists constant  $c = c(t_0) > 0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **uniform asymptotic stability**:

If uniformly stable and there exists constant  $c > 0$  independent of  $t_0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$  uniformly in  $t_0$

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **global uniform asymptotic stability**:

If uniformly asymptotically stable and  $\lim_{\epsilon \rightarrow \infty} \delta(\epsilon) = \infty$

# Stricter Lyapunov Stability of Equilibrium Points

- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **asymptotic stability**:  
If stable and there exists constant  $c = c(t_0) > 0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$
- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **uniform asymptotic stability**:  
If uniformly stable and there exists constant  $c > 0$  independent of  $t_0$  such that  $\vec{x} \rightarrow 0$  as  $t \rightarrow \infty$  for all  $\|\vec{x}\| \leq c$  uniformly in  $t_0$
- Equilibrium point,  $\vec{x} = \vec{0}$ , i.e. origin, has **global uniform asymptotic stability**:  
If uniformly asymptotically stable and  $\lim_{\epsilon \rightarrow \infty} \delta(\epsilon) = \infty$
- Uniform asymptotic stability typically highly desirably property of control system design: able to maintain their closed-loop performance in presence of state perturbations and disturbances

# Lyapunov Indirect and Direct Methods

- In dissertation on stability of motion, Lyapunov developed indirect and direct methods for assessing stability of nominal solutions to dynamical systems governed by finite number of coupled ODEs
  - Provide verifiable sufficient conditions for stability of nominal trajectory without explicit knowledge of system solutions



# Lyapunov Indirect and Direct Methods

- In dissertation on stability of motion, Lyapunov developed indirect and direct methods for assessing stability of nominal solutions to dynamical systems governed by finite number of coupled ODEs
  - Provide verifiable sufficient conditions for stability of nominal trajectory without explicit knowledge of system solutions
- **Lyapunov's indirect method:** can determine stability of equilibrium point, i.e. origin, for nonlinear, time-invariant  $n$ -dimensional systems by linearizing system dynamics about equilibrium point
  - Utilized in previous systems theory discussions

# Lyapunov Indirect and Direct Methods

- In dissertation on stability of motion, Lyapunov developed indirect and direct methods for assessing stability of nominal solutions to dynamical systems governed by finite number of coupled ODEs
  - Provide verifiable sufficient conditions for stability of nominal trajectory without explicit knowledge of system solutions
- **Lyapunov's indirect method:** can determine stability of equilibrium point, i.e. origin, for nonlinear, time-invariant  $n$ -dimensional systems by linearizing system dynamics about equilibrium point
  - Utilized in previous systems theory discussions
- **Lyapunov's direct method** requires concepts:
  - Positive and negative definite functions
  - Time derivative of scalar function along state trajectories of differential equation, i.e. possible solutions

# Positive and Negative Definite Functions

- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V > 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive definite**

# Positive and Negative Definite Functions

- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V > 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive semi-definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V \geq 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive semi-definite**

# Positive and Negative Definite Functions

- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V > 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive semi-definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V \geq 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive semi-definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally negative definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V < 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally negative definite**

# Positive and Negative Definite Functions

- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V > 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally positive semi-definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V \geq 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally positive semi-definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally negative definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V < 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally negative definite**
- $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$  of vector argument  $\vec{x} \in \mathbb{R}^n$ : **locally negative semi-definite**  
If  $V(\vec{0}) = 0$  and there exists constant  $\epsilon$  such that  $V \leq 0$  for all  $\vec{x} \in \mathbb{R}^n$  in neighborhood of origin, i.e.  $B_\epsilon = \{\vec{x} \in \mathbb{R}^n : \|\vec{x}\| \leq \epsilon\}$ 
  - If  $\epsilon = \infty$ , then  $V(\vec{x})$  **globally negative semi-definite**

# Time Derivative Along State Trajectories

- Scalar continuously differentiable function  $V(\vec{x})$ 
  - $\vec{x}(t) \in \mathbb{R}^n$  represents time-varying trajectory of time-varying system

# Time Derivative Along State Trajectories

- Scalar continuously differentiable function  $V(\vec{x})$ 
  - $\vec{x}(t) \in \mathbb{R}^n$  represents time-varying trajectory of time-varying system
- Time derivative of  $V(\vec{x}(t))$  along system solution  $\vec{x}(t) = [x_1(t), \dots, x_n(t)]^T$ :

$$\dot{V}(\vec{x}) = \sum_{i=1}^n \frac{\partial V}{\partial x_i} \dot{x}_i = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(t, \vec{x}) = \nabla V(\vec{x}) f(t, \vec{x}) \quad (12)$$

- $\nabla V(\vec{x}) = [\frac{\partial V}{\partial x_1}, \dots, \frac{\partial V}{\partial x_n}]$ : row vector gradient of  $V(\vec{x})$  w.r.t.  $\vec{x}$
- Time derivative of  $V(\vec{x})$  depends not only on function  $V(\vec{x})$  but also on system dynamics under consideration
- Changing latter while keeping same  $V(\vec{x})$  may result in different  $\dot{V}(\vec{x})$



# Time Derivative Along State Trajectories

- Scalar continuously differentiable function  $V(\vec{x})$ 
  - $\vec{x}(t) \in \mathbb{R}^n$  represents time-varying trajectory of time-varying system
- Time derivative of  $V(\vec{x}(t))$  along system solution  $\vec{x}(t) = [x_1(t), \dots, x_n(t)]^T$ :

$$\dot{V}(\vec{x}) = \sum_{i=1}^n \frac{\partial V}{\partial x_i} \dot{x}_i = \sum_{i=1}^n \frac{\partial V}{\partial x_i} f_i(t, \vec{x}) = \nabla V(\vec{x}) f(t, \vec{x}) \quad (12)$$

- $\nabla V(\vec{x}) = [\frac{\partial V}{\partial x_1}, \dots, \frac{\partial V}{\partial x_n}]$ : row vector gradient of  $V(\vec{x})$  w.r.t.  $\vec{x}$
- Time derivative of  $V(\vec{x})$  depends not only on function  $V(\vec{x})$  but also on system dynamics under consideration
- Changing latter while keeping same  $V(\vec{x})$  may result in different  $\dot{V}(\vec{x})$
- Lyapunov direct method: let  $\vec{\bar{x}} = \vec{0} \in \mathbb{R}^n$  as equilibrium point for time-varying dynamics, initial conditions drawn from domain  $D \subset \mathbb{R}^n$  with  $\vec{\bar{x}} \in D$  and  $t_0 = 0$

# Lyapunov Direct Method

- If on domain  $D$ , there exists continuously differentiable locally positive definite function  $V(\vec{x}) : D \rightarrow \mathbb{R}$ , whose time derivative along system trajectories locally negative semi-definite, i.e.

$$\dot{V}(\vec{x}) = \nabla V(\vec{x})f(t, \vec{x}) \leq 0 \quad (13)$$

for all  $t \geq 0$  and for all  $\vec{x} \in D$ , then  $\vec{x} = \vec{0}$  locally uniformly stable

- If  $\dot{V}(\vec{x}) < 0$  for all  $t \geq 0$ , i.e. time derivative along the system trajectories locally negative definite, then  $\vec{x} = \vec{0}$  locally uniformly asymptotically stable

# Lyapunov Direct Method

- If on domain  $D$ , there exists continuously differentiable locally positive definite function  $V(\vec{x}) : D \rightarrow \mathbb{R}$ , whose time derivative along system trajectories locally negative semi-definite, i.e.

$$\dot{V}(\vec{x}) = \nabla V(\vec{x})f(t, \vec{x}) \leq 0 \quad (13)$$

for all  $t \geq 0$  and for all  $\vec{x} \in D$ , then  $\vec{x} = \vec{0}$  locally uniformly stable

- If  $\dot{V}(\vec{x}) < 0$  for all  $t \geq 0$ , i.e. time derivative along the system trajectories locally negative definite, then  $\vec{x} = \vec{0}$  locally uniformly asymptotically stable
- Any locally positive definite  $V(\vec{x})$ : **Lyapunov function candidate**, if satisfies time derivative condition: **Lyapunov function**
  - Existence of Lyapunov function sufficient to claim uniform stability for equilibrium point, if one cannot be found, nothing can be stated about stability of equilibrium point
  - Lyapunov functions not unique

# Lyapunov Functions and Definitions

- Lyapunov function viewable as “energy-like” function for testing system stability
  - If values of  $V$  do not increase along system trajectories, then origin uniformly stable
  - If  $V$  strictly decreases, then, in addition, system trajectories will approach origin asymptotically
  - Uniform asymptotic stability requires subset of  $D$  known as **region of attraction**, i.e. starting there, system solutions will converge to origin
  - If region of attraction of uniformly asymptotically stable equilibrium:  $\mathbb{R}^n$ , then equilibrium globally uniformly asymptotically stable

# Lyapunov Functions and Definitions

- Lyapunov function viewable as “energy-like” function for testing system stability
  - If values of  $V$  do not increase along system trajectories, then origin uniformly stable
  - If  $V$  strictly decreases, then, in addition, system trajectories will approach origin asymptotically
  - Uniform asymptotic stability requires subset of  $D$  known as **region of attraction**, i.e. starting there, system solutions will converge to origin
  - If region of attraction of uniformly asymptotically stable equilibrium:  $\mathbb{R}^n$ , then equilibrium globally uniformly asymptotically stable
- Define property  $\lim_{\|\vec{x}\| \rightarrow \infty} V(\vec{x}) = \infty$  where  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$ : **radially unbounded**  
Lyapunov function candidate

# Lyapunov Functions and Definitions

- Lyapunov function viewable as “energy-like” function for testing system stability
  - If values of  $V$  do not increase along system trajectories, then origin uniformly stable
  - If  $V$  strictly decreases, then, in addition, system trajectories will approach origin asymptotically
  - Uniform asymptotic stability requires subset of  $D$  known as **region of attraction**, i.e. starting there, system solutions will converge to origin
  - If region of attraction of uniformly asymptotically stable equilibrium:  $\mathbb{R}^n$ , then equilibrium globally uniformly asymptotically stable
- Define property  $\lim_{\|\vec{x}\| \rightarrow \infty} V(\vec{x}) = \infty$  where  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$ : **radially unbounded**  
Lyapunov function candidate
- Define  $V_c = \{\vec{x} \in \mathbb{R}^n : V(\vec{x}) = c\}$  as **level set**
  - i.e.  $V_c$  has constant value  $c$ , of radially unbounded Lyapunov function candidate  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$

# Lyapunov Functions and Definitions

- Lyapunov function viewable as “energy-like” function for testing system stability
  - If values of  $V$  do not increase along system trajectories, then origin uniformly stable
  - If  $V$  strictly decreases, then, in addition, system trajectories will approach origin asymptotically
  - Uniform asymptotic stability requires subset of  $D$  known as **region of attraction**, i.e. starting there, system solutions will converge to origin
  - If region of attraction of uniformly asymptotically stable equilibrium:  $\mathbb{R}^n$ , then equilibrium globally uniformly asymptotically stable
- Define property  $\lim_{\|\vec{x}\| \rightarrow \infty} V(\vec{x}) = \infty$  where  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$ : **radially unbounded**  
Lyapunov function candidate
- Define  $V_c = \{\vec{x} \in \mathbb{R}^n : V(\vec{x}) = c\}$  as **level set**
  - i.e.  $V_c$  has constant value  $c$ , of radially unbounded Lyapunov function candidate  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$
- Define  $\Omega_c = \{\vec{x} \in \mathbb{R}^n : V(\vec{x}) \leq c\}$  as union of interior set of  $V_c$  and  $V_c$  itself

## Lyapunov Functions (continued)

- Consider converging sequence  $\lim_{k \rightarrow \infty} \vec{x}_k = \vec{a}$  with all  $\vec{x}$  from  $\Omega_c$ , then limit point  $\vec{a}$  must also be in  $\Omega_c$



# Lyapunov Functions (continued)

- Consider converging sequence  $\lim_{k \rightarrow \infty} \vec{x}_k = \vec{a}$  with all  $\vec{x}$  from  $\Omega_c$ , then limit point  $\vec{a}$  must also be in  $\Omega_c$ 
  - $\Omega_c$ : **closed set**
    - Since  $V(\vec{x})$  continuous on  $\mathbb{R}^n$  and  $V(\vec{x}) \leq c$  for all  $k = 1, 2, \dots \rightarrow c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = V(\vec{a})$ , and consequently  $\vec{a} \in \Omega_c$ : every converging sequence has limit point in same set

# Lyapunov Functions (continued)

- Consider converging sequence  $\lim_{k \rightarrow \infty} \vec{x}_k = \vec{a}$  with all  $\vec{x}$  from  $\Omega_c$ , then limit point  $\vec{a}$  must also be in  $\Omega_c$ 
  - $\Omega_c$ : **closed set**
    - Since  $V(\vec{x})$  continuous on  $\mathbb{R}^n$  and  $V(\vec{x}) \leq c$  for all  $k = 1, 2, \dots \rightarrow c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = V(\vec{a})$ , and consequently  $\vec{a} \in \Omega_c$ : every converging sequence has limit point in same set
  - $\Omega_c$ : **bounded set**
    - If not, then there must exist sequence of points  $\{\vec{x}_k\} \in \Omega_c$  with limit  $\infty$   
Since  $V(\vec{x})$  continuous and radially unbounded, then  $c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = \infty \rightarrow$  contradiction

# Lyapunov Functions (continued)

- Consider converging sequence  $\lim_{k \rightarrow \infty} \vec{x}_k = \vec{a}$  with all  $\vec{x}$  from  $\Omega_c$ , then limit point  $\vec{a}$  must also be in  $\Omega_c$ 
  - $\Omega_c$ : **closed set**
    - Since  $V(\vec{x})$  continuous on  $\mathbb{R}^n$  and  $V(\vec{x}) \leq c$  for all  $k = 1, 2, \dots \rightarrow c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = V(\vec{a})$ , and consequently  $\vec{a} \in \Omega_c$ : every converging sequence has limit point in same set
  - $\Omega_c$ : **bounded set**
    - If not, then there must exist sequence of points  $\{\vec{x}_k\} \in \Omega_c$  with limit  $\infty$   
Since  $V(\vec{x})$  continuous and radially unbounded, then  $c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = \infty \rightarrow$  contradiction
  - $\Omega_c$  closed, bounded, and belongs to  $\mathbb{R}^n$ : **compact set**

# Lyapunov Functions (continued)

- Consider converging sequence  $\lim_{k \rightarrow \infty} \vec{x}_k = \vec{a}$  with all  $\vec{x}$  from  $\Omega_c$ , then limit point  $\vec{a}$  must also be in  $\Omega_c$ 
  - $\Omega_c$ : **closed set**
    - Since  $V(\vec{x})$  continuous on  $\mathbb{R}^n$  and  $V(\vec{x}) \leq c$  for all  $k = 1, 2, \dots \rightarrow c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = V(\vec{a})$ , and consequently  $\vec{a} \in \Omega_c$ : every converging sequence has limit point in same set
  - $\Omega_c$ : **bounded set**
    - If not, then there must exist sequence of points  $\{\vec{x}_k\} \in \Omega_c$  with limit  $\infty$   
Since  $V(\vec{x})$  continuous and radially unbounded, then  $c \geq \lim_{k \rightarrow \infty} V(\vec{x}_k) = \infty \rightarrow$  contradiction
  - $\Omega_c$  closed, bounded, and belongs to  $\mathbb{R}^n$ : **compact set**
- Krasovskii-LaSalle theorem:**  
If  $\vec{x} = \vec{0}$ : equilibrium point of  $\dot{\vec{x}} = f(t, \vec{x})$  and  $V(\vec{x}) : \mathbb{R}^n \rightarrow \mathbb{R}$ : radially unbounded Lyapunov function  
Then  $\vec{x}$  globally uniformly asymptotically stable equilibrium point
  - Simple example of radially unbounded Lyapunov function candidates include quadratic form  $V(\vec{x}) = \vec{x}^T P \vec{x}$  where  $P$  symmetric positive definite matrix, i.e.  $P = P^T > 0$

# Linear, Time-Varying Dynamical System

- **Linear, time-varying (LTV) systems:**

$$\begin{aligned}\dot{\vec{x}}(t) &= A(t)\vec{x}(t) + B(t)\vec{u}(t) \\ \vec{y}(t) &= C(t)\vec{x}(t) + D(t)\vec{u}(t)\end{aligned}\tag{14}$$

- Similar to LTI, general solution of LTV/LPV system

$$\vec{x}(t) = \Phi(t, t_0)\vec{x}(t_0) + \int_{t_0}^t \Phi(t, \tau)B(\tau)\vec{u}(\tau)d\tau\tag{15}$$

- $\Phi(t, t_0)$ : state transition matrix of LTV/LPV system

$$\Phi(t, t_0) = I + \int_{t_0}^t A(\sigma_1)d\sigma_1 + \int_{t_0}^t A(\sigma_1) \int_{t_0}^{\sigma_1} A(\sigma_2)d\sigma_2 d\sigma_1 + \cdots\tag{16}$$

- $\Phi(t, t_0)$  converges uniformly and absolutely to solution that exists and unique

# Lyapunov Stability of LTV Systems

- Lyapunov candidate function with  $P > 0$

$$V(\vec{x}) = \vec{x}^T P \vec{x} \quad (17)$$

# Lyapunov Stability of LTV Systems

- Lyapunov candidate function with  $P > 0$

$$V(\vec{x}) = \vec{x}^T P \vec{x} \quad (17)$$

- Time derivative:

$$\dot{V}(\vec{x}) = \dot{\vec{x}}^T P \vec{x} + \vec{x}^T P \dot{\vec{x}} \quad (18)$$

# Lyapunov Stability of LTV Systems

- Lyapunov candidate function with  $P > 0$

$$V(\vec{x}) = \vec{x}^T P \vec{x} \quad (17)$$

- Time derivative:

$$\dot{V}(\vec{x}) = \dot{\vec{x}}^T P \vec{x} + \vec{x}^T P \dot{\vec{x}} \quad (18)$$

- Evaluated along state trajectories:

$$\dot{V}(\vec{x}) = \vec{x}^T (A^T(t)P + PA(t)) \vec{x} \quad (19)$$



# Lyapunov Stability of LTV Systems

- Lyapunov candidate function with  $P > 0$

$$V(\vec{x}) = \vec{x}^T P \vec{x} \quad (17)$$

- Time derivative:

$$\dot{V}(\vec{x}) = \dot{\vec{x}}^T P \vec{x} + \vec{x}^T P \dot{\vec{x}} \quad (18)$$

- Evaluated along state trajectories:

$$\dot{V}(\vec{x}) = \vec{x}^T (A^T(t)P + PA(t)) \vec{x} \quad (19)$$

- $\dot{V}$  negative definite if and only if

$$A^T(t)P + PA(t) < 0 \quad (20)$$

for all  $t$

# Lyapunov Stability of LTV Systems

- Lyapunov candidate function with  $P > 0$

$$V(\vec{x}) = \vec{x}^T P \vec{x} \quad (17)$$

- Time derivative:

$$\dot{V}(\vec{x}) = \dot{\vec{x}}^T P \vec{x} + \vec{x}^T P \dot{\vec{x}} \quad (18)$$

- Evaluated along state trajectories:

$$\dot{V}(\vec{x}) = \vec{x}^T (A^T(t)P + PA(t)) \vec{x} \quad (19)$$

- $\dot{V}$  negative definite if and only if

$$A^T(t)P + PA(t) < 0 \quad (20)$$

for all  $t$

- Stable LTV systems: globally uniformly asymptotically stable

# Controllability and Observability

- **Linear, time-varying controllability Gramian** using state-transition matrix:

$$W_C(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t_0, t) B(t) B(t)^T \Phi(t_0, t)^T dt \quad (21)$$

- $W_C(t_0, t_1)$ : symmetric, positive semi-definite, satisfies

$$W_C(t_0, t_1) = W_C(t_0, t) + \Phi(t_0, t) W(t, t_1) \Phi(t_0, t)^T \quad (22)$$

# Controllability and Observability

- **Linear, time-varying controllability Gramian** using state-transition matrix:

$$W_C(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t_0, t) B(t) B(t)^T \Phi(t_0, t)^T dt \quad (21)$$

- $W_C(t_0, t_1)$ : symmetric, positive semi-definite, satisfies

$$W_C(t_0, t_1) = W_C(t_0, t) + \Phi(t_0, t) W(t, t_1) \Phi(t_0, t)^T \quad (22)$$

- If there exists some  $t_1$  such that  $W_C(t_0, t_1)$  full rank, then LTV system state controllable

# Controllability and Observability

- **Linear, time-varying controllability Gramian** using state-transition matrix:

$$W_C(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t_0, t) B(t) B(t)^T \Phi(t_0, t)^T dt \quad (21)$$

- $W_C(t_0, t_1)$ : symmetric, positive semi-definite, satisfies

$$W_C(t_0, t_1) = W_C(t_0, t) + \Phi(t_0, t) W(t, t_1) \Phi(t_0, t)^T \quad (22)$$

- If there exists some  $t_1$  such that  $W_C(t_0, t_1)$  full rank, then LTV system state controllable

- **Linear, time-varying observability Gramian** using state-transition matrix:

$$W_O(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t, t_0)^T C(t)^T C(t) \Phi(t, t_0) dt \quad (23)$$

- $W_O(t_0, t_1)$  symmetric, positive semi-definite, satisfies

$$W_O(t_0, t_1) = W_O(t_0, t) + \Phi(t, t_0)^T W(t, t_1) \Phi(t, t_0) \quad (24)$$

# Controllability and Observability

- **Linear, time-varying controllability Gramian** using state-transition matrix:

$$W_C(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t_0, t) B(t) B(t)^T \Phi(t_0, t)^T dt \quad (21)$$

- $W_C(t_0, t_1)$ : symmetric, positive semi-definite, satisfies

$$W_C(t_0, t_1) = W_C(t_0, t) + \Phi(t_0, t) W(t, t_1) \Phi(t_0, t)^T \quad (22)$$

- If there exists some  $t_1$  such that  $W_C(t_0, t_1)$  full rank, then LTV system state controllable

- **Linear, time-varying observability Gramian** using state-transition matrix:

$$W_O(t_0, t_1) = \int_{t_0}^{t_1} \Phi(t, t_0)^T C(t)^T C(t) \Phi(t, t_0) dt \quad (23)$$

- $W_O(t_0, t_1)$  symmetric, positive semi-definite, satisfies

$$W_O(t_0, t_1) = W_O(t_0, t) + \Phi(t, t_0)^T W(t, t_1) \Phi(t, t_0) \quad (24)$$

- If there exists some  $t_1$  such that  $W_O(t_0, t_1)$  full rank, then LTV system observable

# Linear, Parameter-Varying Systems

- Special class of LTV systems: **linear, parameter-varying (LPV) systems**:

$$\begin{aligned}\dot{\vec{x}}(t) &= A(\vec{\beta}(t))\vec{x}(t) + B(\vec{\beta}(t))\vec{u}(t) \\ \vec{y}(t) &= C(\vec{\beta}(t))\vec{x}(t) + D(\vec{\beta}(t))\vec{u}(t)\end{aligned}\tag{25}$$

- $\vec{\beta}(t)$ : **parameter vector** must be continuously differentiable function of time
- State-space matrices continuous functions of  $\vec{\beta}$
- Typically arise when linearizing nonlinear system about equilibrium specified by  $\vec{\beta}$

# Linear, Parameter-Varying Systems

- Special class of LTV systems: **linear, parameter-varying (LPV) systems**:

$$\begin{aligned}\dot{\vec{x}}(t) &= A(\vec{\beta}(t))\vec{x}(t) + B(\vec{\beta}(t))\vec{u}(t) \\ \vec{y}(t) &= C(\vec{\beta}(t))\vec{x}(t) + D(\vec{\beta}(t))\vec{u}(t)\end{aligned}\tag{25}$$

- $\vec{\beta}(t)$ : **parameter vector** must be continuously differentiable function of time
  - State-space matrices continuous functions of  $\vec{\beta}$
  - Typically arise when linearizing nonlinear system about equilibrium specified by  $\vec{\beta}$
- $\vec{\beta}(t)$  typically also has restricted admissible trajectories for any particular realization
  - Typically at least restricted by some upper and lower bounds, e.g.  $\vec{\beta}_U$  and  $\vec{\beta}_L$
  - May also be rate restricted by some upper and lower bounds, e.g.  $\dot{\vec{\beta}}_U$  and  $\dot{\vec{\beta}}_L$
  - $\mathcal{B}$ : set of admissible trajectories for  $\vec{\beta}$

$$\mathcal{B} = \left\{ \vec{\beta}(t) : \mathbb{R}_+ \rightarrow \mathbb{R}^{n_\beta} \text{ such that: } \vec{\beta}(t) \text{ continuously differentiable,} \right. \\ \left. \vec{\beta}_L \leq \vec{\beta}(t) \leq \vec{\beta}_U \forall t \geq 0, \quad \dot{\vec{\beta}}_L \leq \dot{\vec{\beta}}(t) \leq \dot{\vec{\beta}}_U \forall t \geq 0 \right\} \tag{26}$$



# LPV Control Design Approaches

- Common LPV control design approach: **gain scheduling**
  - Compute array of optimal LTI controllers at variety of *different* equilibrium points across ranges of admissible  $\vec{\beta}$ , e.g. obtain *different* optimal LTI feedback gain matrices
  - Blend various LTI controllers to obtain single, integrated control design, e.g. interpolate between different optimal LTI feedback gain matrices if operating away from equilibrium points
  - Works well for locally “smooth” systems i.e. linearized dynamics reasonably accurate near given equilibrium condition
  - Nonlinearities have significant effect over entire range of  $\vec{\beta}$

# LPV Control Design Approaches


- Common LPV control design approach: **gain scheduling**
  - Compute array of optimal LTI controllers at variety of *different* equilibrium points across ranges of admissible  $\vec{\beta}$ , e.g. obtain *different* optimal LTI feedback gain matrices
  - Blend various LTI controllers to obtain single, integrated control design, e.g. interpolate between different optimal LTI feedback gain matrices if operating away from equilibrium points
  - Works well for locally “smooth” systems i.e. linearized dynamics reasonably accurate near given equilibrium condition
  - Nonlinearities have significant effect over entire range of  $\vec{\beta}$
- More advanced approaches for LPV systems proposed based on specific structure of state-space matrices, e.g. linear or rational dependence on  $\vec{\beta}$

# LPV Control Design Approaches

- Common LPV control design approach: **gain scheduling**
  - Compute array of optimal LTI controllers at variety of *different* equilibrium points across ranges of admissible  $\vec{\beta}$ , e.g. obtain *different* optimal LTI feedback gain matrices
  - Blend various LTI controllers to obtain single, integrated control design, e.g. interpolate between different optimal LTI feedback gain matrices if operating away from equilibrium points
  - Works well for locally “smooth” systems i.e. linearized dynamics reasonably accurate near given equilibrium condition
  - Nonlinearities have significant effect over entire range of  $\vec{\beta}$
- More advanced approaches for LPV systems proposed based on specific structure of state-space matrices, e.g. linear or rational dependence on  $\vec{\beta}$
- For arbitrary dependence, specify performance of LPV system,  $G_{\beta}$  in terms of maximum possible induced  $\mathcal{L}_{2 \leftarrow 2}$  gain from input  $\vec{u}$  to output  $\vec{y}$ , i.e.

$$\|G_{\beta}\|_{2 \leftarrow 2} = \max_{\vec{\beta} \in \mathcal{B}, 0 \neq \|\vec{u}\|_2 \leq \infty, \vec{x}(0)=0} \frac{\|\vec{y}\|_2}{\|\vec{u}\|_2} \quad (27)$$

# State Feedback Linear, Parameter-Varying OCP



- LPV plant:

$$\begin{aligned} \dot{\vec{x}}(t) &= A(\vec{\beta})\vec{x}(t) + \begin{bmatrix} B_1(\vec{\beta}) & B_2(\vec{\beta}) \end{bmatrix} \begin{bmatrix} \vec{d}(t) \\ \vec{u}(t) \end{bmatrix} \\ \begin{bmatrix} \vec{e}(t) \\ \vec{y}(t) \end{bmatrix} &= \begin{bmatrix} C_1(\vec{\beta}) \\ I \end{bmatrix} \vec{x}(t) + \begin{bmatrix} 0 & D_{12}(\vec{\beta}) \\ 0 & 0 \end{bmatrix} \begin{bmatrix} \vec{d}(t) \\ \vec{u}(t) \end{bmatrix} \end{aligned} \quad (28)$$

- State feedback LPV controller:

$$\vec{u} = D_K(\vec{\beta})\vec{x} \quad (29)$$

- Could also use output feedback or observer feedback

# Generalized Bounded Real Lemma

- As  $G_\beta$  time-varying, *cannot* interpret as  $\mathcal{H}_\infty$ -norm in frequency domain
  - **Generalized bounded real lemma:** derive sufficient conditions for upper bound on  $\|G_\beta\|_{2\leftarrow 2}$

# Generalized Bounded Real Lemma

- As  $G_\beta$  time-varying, *cannot* interpret as  $\mathcal{H}_\infty$ -norm in frequency domain
  - Generalized bounded real lemma:** derive sufficient conditions for upper bound on  $\|G_\beta\|_{2 \leftarrow 2}$
- Rate-unbounded LPV system globally uniformly asymptotically stable and  $\|G_\beta\|_{2 \leftarrow 2} \leq \gamma$ :  
 If there exists  $P > 0$  such that  $\forall \vec{\beta} \in [\vec{\beta}_L, \vec{\beta}_U]$ :

$$\begin{bmatrix} A^T(\vec{\beta})P + PA(\vec{\beta}) & PB(\vec{\beta}) \\ B^T(\vec{\beta})P & -I \end{bmatrix} + \frac{1}{\gamma^2} \begin{bmatrix} C^T(\vec{\beta}) \\ D^T(\vec{\beta}) \end{bmatrix} \begin{bmatrix} C(\vec{\beta}) & D(\vec{\beta}) \end{bmatrix} < 0 \quad (30)$$

# Generalized Bounded Real Lemma

- As  $G_\beta$  time-varying, *cannot* interpret as  $\mathcal{H}_\infty$ -norm in frequency domain
  - Generalized bounded real lemma:** derive sufficient conditions for upper bound on  $\|G_\beta\|_{2\leftarrow 2}$
- Rate-unbounded LPV system globally uniformly asymptotically stable and  $\|G_\beta\|_{2\leftarrow 2} \leq \gamma$ :  
If there exists  $P > 0$  such that  $\forall \vec{\beta} \in [\vec{\beta}_L, \vec{\beta}_U]$ :

$$\begin{bmatrix} A^T(\vec{\beta})P + PA(\vec{\beta}) & PB(\vec{\beta}) \\ B^T(\vec{\beta})P & -I \end{bmatrix} + \frac{1}{\gamma^2} \begin{bmatrix} C^T(\vec{\beta}) \\ D^T(\vec{\beta}) \end{bmatrix} \begin{bmatrix} C(\vec{\beta}) & D(\vec{\beta}) \end{bmatrix} < 0 \quad (30)$$

- To show  $\|G_\beta\|_{2\leftarrow 2} \leq \gamma$ : left and right multiply by  $[\vec{x}^T \ \vec{u}^T]^T$ :

$$\dot{\vec{x}}^T P \vec{x} + \vec{x}^T P \dot{\vec{x}} + \frac{1}{\gamma^2} \vec{y}^T \vec{y} - \vec{u}^T \vec{u} \leq 0 \quad (31)$$

$$\dot{V} + \frac{1}{\gamma^2} \vec{y}^T \vec{y} - \vec{u}^T \vec{u} \leq 0 \quad (32)$$

## Generalized Bounded Real Lemma (continued)

- Integrate from  $t = 0$  to  $t = t_f$ :

$$V(\vec{x}(t_f)) - V(\vec{x}(0)) + \frac{1}{\gamma^2} \int_0^{t_f} \vec{y}^T \vec{y} dt - \int_0^{t_f} \vec{u}^T \vec{u} dt \leq 0 \quad (33)$$



## Generalized Bounded Real Lemma (continued)

- Integrate from  $t = 0$  to  $t = t_f$ :

$$V(\vec{x}(t_f)) - V(\vec{x}(0)) + \frac{1}{\gamma^2} \int_0^{t_f} \vec{y}^T \vec{y} dt - \int_0^{t_f} \vec{u}^T \vec{u} dt \leq 0 \quad (33)$$

- $V(\vec{x}(t_f)) > 0$  and  $V(\vec{x}(0)) = 0$ :

$$\int_0^{t_f} \vec{y}^T \vec{y} dt \leq \gamma^2 \int_0^{t_f} \vec{u}^T \vec{u} dt \quad (34)$$

# Generalized Bounded Real Lemma (continued)

- Integrate from  $t = 0$  to  $t = t_f$ :

$$V(\vec{x}(t_f)) - V(\vec{x}(0)) + \frac{1}{\gamma^2} \int_0^{t_f} \vec{y}^T \vec{y} dt - \int_0^{t_f} \vec{u}^T \vec{u} dt \leq 0 \quad (33)$$

- $V(\vec{x}(t_f)) > 0$  and  $V(\vec{x}(0)) = 0$ :

$$\int_0^{t_f} \vec{y}^T \vec{y} dt \leq \gamma^2 \int_0^{t_f} \vec{u}^T \vec{u} dt \quad (34)$$

- As  $t_f \rightarrow \infty$ :

$$\|\vec{y}\|_2^2 \leq \gamma^2 \|\vec{u}\|_2^2 \quad (35)$$

$$\|\mathbf{G}_\beta\|_{2 \leftarrow 2} \leq \gamma \quad (36)$$

## Generalized Bounded Real Lemma (continued)

- Matrix inequality as parameterized LMI condition: one LMI for each value of  $\vec{\beta} \in \mathcal{B}$ 
  - In practice: infinite LMI conditions approximated by enforcing only on finite grid of points
  - Finite dimensional LMI conditions directly obtained to bound  $\|G_{\beta}\|_{2 \leftarrow 2}$  without approximation if state matrices have rational dependence on  $\vec{\beta}$

# Generalized Bounded Real Lemma (continued)

- Matrix inequality as parameterized LMI condition: one LMI for each value of  $\vec{\beta} \in \mathcal{B}$ 
  - In practice: infinite LMI conditions approximated by enforcing only on finite grid of points
  - Finite dimensional LMI conditions directly obtained to bound  $\|G_{\beta}\|_{2 \leftarrow 2}$  without approximation if state matrices have rational dependence on  $\vec{\beta}$
- Rate-bounded LPV system: alter matrix inequality with variation on Lyapunov theory
  - Check  $2^{n_{\beta}}$  LMI conditions evaluated at endpoints defined by hypercube of elements of  $\dot{\vec{\beta}}_U$  and  $\dot{\vec{\beta}}_L$
  - Also need to search over infinitely dimensional space of functions,  $P(\beta)$ , must be restricted to finite dimensional subspace
  - In practice: specify collection of scalar basis functions,  $g_i(\vec{\beta})$ , and use linear combination

$$P(\vec{\beta}) = \sum_{i=1}^N g_i(\vec{\beta}) P_i \quad (37)$$

- $P_i$ : symmetric matrices that form finite collection of decision variables, often polynomials chosen as basis functions

# State Feedback LPV Optimal Control

- Closed-loop LPV system:

$$\begin{aligned}\dot{\vec{x}}(t) &= \left( A(\vec{\beta}) + B_2(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t) + B_1(\vec{\beta})\vec{d}(t) \\ \vec{e}(t) &= \left( C_1(\vec{\beta}) + D_{12}(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t)\end{aligned}\tag{38}$$

# State Feedback LPV Optimal Control

- Closed-loop LPV system:

$$\begin{aligned}\dot{\vec{x}}(t) &= \left( A(\vec{\beta}) + B_2(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t) + B_1(\vec{\beta})\vec{d}(t) \\ \vec{e}(t) &= \left( C_1(\vec{\beta}) + D_{12}(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t)\end{aligned}\tag{38}$$

- By generalized bounded real lemma, closed-loop LPV system internally stable and  $\|G_\beta\|_{2 \leftarrow 2} \leq \gamma$  if there exists  $P > 0$  such that  $\forall \vec{\beta} \in [\vec{\beta}_L, \vec{\beta}_U]$ :

$$\begin{bmatrix} A_L^T(\vec{\beta})P + PA_L(\vec{\beta}) & PB_L(\vec{\beta}) \\ B_L^T(\vec{\beta})P & -I \end{bmatrix} + \frac{1}{\gamma^2} \begin{bmatrix} C_L^T(\vec{\beta}) \\ D_L^T(\vec{\beta}) \end{bmatrix} \begin{bmatrix} C_L(\vec{\beta}) & D_L(\vec{\beta}) \end{bmatrix} < 0 \tag{39}$$

# State Feedback LPV Optimal Control

- Closed-loop LPV system:

$$\begin{aligned}\dot{\vec{x}}(t) &= \left( A(\vec{\beta}) + B_2(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t) + B_1(\vec{\beta})\vec{d}(t) \\ \vec{e}(t) &= \left( C_1(\vec{\beta}) + D_{12}(\vec{\beta})D_K(\vec{\beta}) \right) \vec{x}(t)\end{aligned}\quad (38)$$

- By generalized bounded real lemma, closed-loop LPV system internally stable and  $\|G_\beta\|_{2 \leftarrow 2} \leq \gamma$  if there exists  $P > 0$  such that  $\forall \vec{\beta} \in [\vec{\beta}_L, \vec{\beta}_U]$ :

$$\begin{bmatrix} A_L^T(\vec{\beta})P + PA_L(\vec{\beta}) & PB_L(\vec{\beta}) \\ B_L^T(\vec{\beta})P & -I \end{bmatrix} + \frac{1}{\gamma^2} \begin{bmatrix} C_L^T(\vec{\beta}) \\ D_L^T(\vec{\beta}) \end{bmatrix} \begin{bmatrix} C_L(\vec{\beta}) & D_L(\vec{\beta}) \end{bmatrix} < 0 \quad (39)$$

- Use change of variables:

$$Q = P^{-1} \quad (40)$$

$$R(\vec{\beta}) = D_K(\vec{\beta})Q \quad (41)$$

# State Feedback LPV Optimal Control (continued)

- By Schur Complement Lemma:

$$\begin{bmatrix} QA^T(\vec{\beta}) + A(\vec{\beta})Q + R^T(\vec{\beta})B_2^T(\vec{\beta}) + B_2(\vec{\beta})R(\vec{\beta}) & B_1(\vec{\beta}) & (C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}))^T \\ B_1^T(\vec{\beta}) & \gamma^{-2}I & 0 \\ C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}) & 0 & -I \end{bmatrix} < 0 \quad (42)$$



# State Feedback LPV Optimal Control (continued)

- By Schur Complement Lemma:

$$\begin{bmatrix} QA^T(\vec{\beta}) + A(\vec{\beta})Q + R^T(\vec{\beta})B_2^T(\vec{\beta}) + B_2(\vec{\beta})R(\vec{\beta}) & B_1(\vec{\beta}) & (C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}))^T \\ B_1^T(\vec{\beta}) & \gamma^{-2}I & 0 \\ C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}) & 0 & -I \end{bmatrix} < 0 \quad (42)$$

- State feedback LPV OCP as SDP in  $(Q, R, \gamma)$ :

$$\begin{aligned} (Q, R(\vec{\beta}), \gamma)^{opt} = & \underset{Q \in \mathbb{S}^{n_x}, R \in \mathbb{R}^{n_u \times n_x}, \gamma > 0}{\operatorname{argmin}} \quad \gamma \\ \text{subject to: } & \begin{bmatrix} QA^T(\vec{\beta}) + A(\vec{\beta})Q + R^T(\vec{\beta})B_2^T(\vec{\beta}) + B_2(\vec{\beta})R(\vec{\beta}) & B_1(\vec{\beta}) & (C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}))^T \\ B_1^T(\vec{\beta}) & -\gamma^{-2}I & 0 \\ C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}) & 0 & -I \end{bmatrix} < 0 \\ & Q > 0 \end{aligned} \quad (43)$$

# State Feedback LPV Optimal Control (continued)

- By Schur Complement Lemma:

$$\begin{bmatrix} QA^T(\vec{\beta}) + A(\vec{\beta})Q + R^T(\vec{\beta})B_2^T(\vec{\beta}) + B_2(\vec{\beta})R(\vec{\beta}) & B_1(\vec{\beta}) & (C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}))^T \\ B_1^T(\vec{\beta}) & \gamma^{-2}I & 0 \\ C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}) & 0 & -I \end{bmatrix} < 0 \quad (42)$$

- State feedback LPV OCP as SDP in  $(Q, R, \gamma)$ :

$$\begin{aligned} (Q, R(\vec{\beta}), \gamma)^{opt} = & \underset{Q \in \mathbb{S}^{n_x}, R \in \mathbb{R}^{n_u \times n_x}, \gamma > 0}{\operatorname{argmin}} \gamma \\ \text{subject to: } & \begin{bmatrix} QA^T(\vec{\beta}) + A(\vec{\beta})Q + R^T(\vec{\beta})B_2^T(\vec{\beta}) + B_2(\vec{\beta})R(\vec{\beta}) & B_1(\vec{\beta}) & (C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}))^T \\ B_1^T(\vec{\beta}) & -\gamma^{-2}I & 0 \\ C_1(\vec{\beta})Q + D_{12}(\vec{\beta})R(\vec{\beta}) & 0 & -I \end{bmatrix} < 0 \\ & Q > 0 \end{aligned} \quad (43)$$

- State feedback LPV optimal controller as  $\vec{u}(t) = D_K(\vec{\beta})\vec{x}(t)$  reconstructed with

$$D_K(\vec{\beta}) = R(\vec{\beta})Q^{-1} \quad (44)$$

- Solved approximately by gridding over  $[\vec{\beta}_L, \vec{\beta}_U]$

# Gain-Scheduling



# Summary

- Non-LTI systems requires additional dynamical systems theory
  - Need to show existence and uniqueness: Lipschitz sufficient
  - Need Lyapunov direct stability method for various definitions of stability for equilibrium points

# Summary

- Non-LTI systems requires additional dynamical systems theory
  - Need to show existence and uniqueness: Lipschitz sufficient
  - Need Lyapunov direct stability method for various definitions of stability for equilibrium points
- Linear, Time-Varying (LTV) systems exhibit excellent stability properties using Lyapunov theory
  - General solution, controllability, and observability analysis similar to LTI systems using state-transition matrix