

UThwigl - an R package for closed- and open-system uranium-thorium dating

Anthony Dosseto¹, Ben Marwick¹

^a *Wollongong Isotope Geochronology Laboratory, School of Earth, Atmosphere & Life Sciences, University of Wollongong, Wollongong NSW Australia*

^b *Department of Anthropology, University of Washington, Seattle, WA, USA*

Abstract

For several decades, uranium-thorium (U-Th) dating has allowed geochronologists to precisely date geological materials, providing invaluable geochronological constraints on Quaternary processes. Open-system dating of bones and teeth has also provided ages of human and faunal remains of archaeological significance.

To facilitate access of open-system U-Th dating to the broad scientific community, here we provide an R package, named *UThwigl*, that implements the Diffusion-Adsorption-Decay model of Sambridge et al. (2012). Description of input and output parameters is given, as well as a guide for running the model. The package can be used three different ways: (i) as a web application, (ii) through a web browser with an internet connection, or (iii) in R (most efficiently with RStudio). Examples of application of the model are also provided, showing that it yields ages within error of previously published values.

*Corresponding Author

1 Introduction

2 Uranium-thorium (U-Th) dating has revolutionised Quaternary science and
3 archaeology. Dating uses the decay of ^{238}U into ^{230}Th , with ^{234}U and a few
4 short-lived nuclides as intermediary products. It is based on the principle that
5 the age of formation of a material can be dated as it incorporates U and no or
6 little Th at the time of formation, so all the ^{230}Th in the sample comes from
7 decay of ^{238}U . If detrital Th is included to the sample, a correction must be
8 included to account for the fraction of ^{230}Th which is detrital and not derived
9 from ^{238}U decay. Another requirement is that there is no gain or loss of ^{230}Th ,
10 ^{234}U or ^{238}U after formation of the material (*closed system*).

11 Closed-system U-Th dating has been successfully applied to a range of
12 carbonates, from corals (Edwards, Gallup, and Cheng 2003) to speleothems
13 (Richards and Dorale 2003). In corals and most speleothems, detrital correc-
14 tion is minimal; however, it can be significant when dating pedogenic carbon-
15 ates, for instance (Ludwig and Paces 2002). In this case, detrital correction can
16 be performed using the measured or assumed composition of the detrital frac-
17 tion (e.g. K. Ludwig 2003). Alternatively, isochron techniques can be applied
18 (Ludwig and Titterton 1994); the latter are beyond the scope of this article
19 but IsoPlot is a commonly used software for isochron calculations and other
20 geochronological applications (K. R. Ludwig 2003).

21 Closed-system conditions are seldom met in teeth and bones (although enamel
22 can sometimes be quite impervious to isotope gain or loss). Thus, for teeth and
23 bone, U-Th dating requires to take into account open system behaviour. The
24 diffusion-adsorption-decay (DAD) model developed by Sambridge et al. (2012)
25 was instrumental to implement successfully open-system U-Th dating. It allows
26 for advective and diffusive transport of uranium and thorium isotopes, while
27 include synchronous radioactive decay. The software implementation was writ-
28 ten in Fortran and is available as a Java GUI ([http://www.earth.org.au/codes/](http://www.earth.org.au/codes/iDaD/)
29 [iDaD/](http://www.earth.org.au/codes/iDaD/)).

30 Open-system U-Th dating of teeth and bones, while challenging, has pro-
31 vided quantitative ages for human and faunal remains (Eggins et al. 2005; Grün
32 et al. 2014; Sambridge, Grün, and Eggins 2012). Thus, this approach has signif-
33 icantly improved our understanding of human evolution (e.g. Dirks et al. 2017;
34 Sutikna et al. 2016).

35 In this article, we propose a R package which offers functions to perform
36 closed-system, `csUTh()`, and open-system, `osUTh()`, U-Th age calculations. The
37 former implements formulations given in Ludwig (2003) while the latter applies
38 the Diffusion-Adsorption-Decay (DAD) model of Sambridge et al. (2012). The
39 motivation for providing an R package is to increase the transparency, repro-
40 ducibility, and flexibility of the analytical workflow for computing U-Th ages. In
41 particular, for open-system Currently it is difficult to include the Java GUI in a
42 fully scripted data analysis so the method for computing the DAD model is not
43 highly transparent. This can obscure steps where key decisions are made that
44 are important for others to see to verify the reliability of the analysis. Enabling
45 a scripted workflow for computational analysis of geoscience data is important

for improving the reproducibility of results. Reproducibility refers the ability to recreate the results or retest the hypotheses leading to a scientific claim, either by rerunning the same code used by the original authors, or by writing new code. High rates of irreproducibility of research results have been estimated in several fields and disciplines (Medical Sciences 2015; Freedman, Cockburn, and Simcoe 2015; Institute 2013; Ioannidis 2005; Collaboration and others 2015; Camerer et al. 2018; Camerer et al. 2016). Consequently, the transparency, openness, and reproducibility of results and methods are receiving increased attention, and the norms of research in many fields are changing (Nosek et al. 2015; Miguel et al. 2014; Marwick 2016).

There is strong interest in open, transparent, and reusable research in the geoscience community (Gil et al. 2016) and substantial progress toward open data has been made in the geosciences with the widespread use of data services of NASA, USGS, NOAA and community-built data portals such as OneGeology, EarthChem, RRUFF, PANGAEA, PaleoBioDB, and others (Kattge, Díaz, and Wirth 2014; Ma 2018). However, the use of open source software such as R (Pebesma, Nüst, and Bivand 2012), and sharing of scripted data analysis workflows with research publications is not yet widespread (Hutton et al. 2016). With this R package our goal is to make scripted and reproducible data analysis easy for open-system uranium-thorium dating. This will improve the transparency of geochronology research, and provide a more credible and robust foundation for scientific advancement (Hutton et al. 2016).

To enable re-use of our materials and improve reproducibility and transparency, all the results and visualisations in this paper can be reproduced using the RMarkdown vignette document included with the UThwgl package. We have archived these files at <http://doi.org/10.17605/OSF.IO/D5P7S> to ensure long-term accessibility. Our code is released under the MIT licence, our data as CC-0, and our figures as CC-BY, to enable maximum re-use (for more details, see Marwick 2016).

Methods

Closed-system dating

Pending closed-system behaviour can be assessed, it is possible to derive an age for each U-Th analysis. The closed-system function `csUTh()` requires that for each analysis to yield an age, $(^{234}\text{U}/^{238}\text{U})$, $(^{230}\text{Th}/^{238}\text{U})$ and $(^{232}\text{Th}/^{238}\text{U})$ activity ratios are measured (parentheses denote activity ratios throughout this article). The $(^{232}\text{Th}/^{238}\text{U})$ activity ratio is required for detrital correction (although it is needed to use `csUTh()` whether the detrital correction is performed or not).

Open-system dating

Data required for the DAD model are $(^{230}\text{Th}/^{238}\text{U})$ and $(^{234}\text{U}/^{238}\text{U})$ activity ratios collected along a transect perpendicular to the surface of the tooth or bone. Sampling for analysis can be done by micro-drilling or laser ablation.