

Many-Body Theory I

01.17.23

I. Review of single-particle quantum mechanics + Stat. mech. of a quantum system.

1. Wave function. Max Born interpretation: prob. density of finding a particle at point \vec{r} is

$$P(\vec{r}, t) = |\psi(\vec{r}, t)|^2$$

2. Single-particle Schrödinger equation:

$$i\hbar \frac{\partial \psi}{\partial t} = \hat{\mathcal{H}} \psi, \quad \hat{\mathcal{H}} = \frac{\hat{p}^2}{2m} + V(\hat{r})$$

In coordinate rep.: $\vec{p} = -i\hbar \vec{\nabla}; \quad \hat{\mathcal{H}} = -\frac{\hbar^2}{2m}(\vec{\nabla})^2 + V(\vec{r})$

3. Linear superposition of waves:

If $\psi_a(\vec{r}, t)$, $\psi_b(\vec{r}, t)$ are solutions of SE, then


$\psi_c = a\psi_a + b\psi_b$ is also a solution

- 3a. Stationary states, eigenstates

$$\psi_n(\vec{r}, t) = e^{-i\varepsilon_n t/\hbar} \psi_n(\vec{r}) \Rightarrow \hat{\mathcal{H}} \psi_n(\vec{r}) = \varepsilon_n \psi_n(\vec{r})$$

$$\psi(\vec{r}, t) = \sum_n c_n e^{-i\varepsilon_n t/\hbar} \psi_n(\vec{r})$$

4. Quantum Gibbs distribution



$$\Psi(r_1, r_2, \dots, r_N)$$

$$\left(\prod_{i=1}^N \int d\vec{r}_i \right) \Psi^*(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N)$$

$$\equiv P(\vec{r}, \vec{r}') : \text{density matrix (coord. rep.)}$$

$$P(r_i) = P(r_i, r_i)$$

We may expand $\rho(\vec{r}, \vec{r}')$ in eigenfunctions $\psi_n(\vec{r})$ of a single-particle Hamiltonian (describing an isolated particle)

$$\rho(r, r') = \sum_{nm} w_{mn} \psi_n^*(r) \psi_m(r') ; \quad \hat{\rho} = \sum_{nm} |n\rangle w_{mn} \langle n|$$

density matrix (Dirac notation)

Quantum Gibbs distribution :
(equilibrium distribution)

$$w_{mn} = \frac{1}{Z} e^{-\beta \epsilon_n} \delta_{mn}$$

$$\beta = \frac{1}{k_B T} \quad (\text{we will mostly stick to } k_B = 1)$$

Partition function: $Z = \sum_n e^{-\beta \epsilon_n}$ (so that $\sum_n w_{nn} = 1$)

Quantum Gibbs distribution, statistical operator

$$\hat{\rho}_G = \frac{1}{Z} \sum_n |n\rangle e^{-\beta \epsilon_n} \langle n| = \frac{1}{Z} e^{-\beta \hat{H}}$$

$$\text{Partition function: } Z = \sum_n e^{-\beta \epsilon_n} = \sum_n \langle n | e^{-\beta \hat{H}} | n \rangle = \text{Tr } e^{-\beta \hat{H}}$$

The traditional point of view at microcanonical (fixed energy) and thermal (Gibbs) distributions is based on the notion of statistical ensemble, see Huang, Stat. Mech. Ch. 8 and also employs the notion of ergodicity in evolution of a classical system.

A modern point of view is based on the eigenstate thermalization hypothesis (ETH), see Mark Srednicki (Chaos and Quantum Therm.) Phys. Rev. E, 50, 888 (1994)

01.19.23

5. Classical limit in Statistical Mechanics:

$$Z_{cl} = \frac{1}{(2\pi\hbar)^d} \int d\vec{p} \int d\vec{x} e^{-\beta \hat{H}(p, x)} \quad \text{vs. } Z = \text{Tr } e^{-\beta \hat{H}}$$

Example: 1D Harmonic oscillator $\hat{H} = \frac{\hat{p}^2}{2m} + \frac{m\omega_0^2}{2} x^2$ (1D)

$$\epsilon_n = (n + \frac{1}{2})\hbar\omega_0$$

$$Z = \sum_{n=0}^{\infty} e^{-\beta \hbar \omega_0 (n + \frac{1}{2})} = \frac{e^{-\beta \hbar \omega_0 / 2}}{1 - e^{-\beta \hbar \omega_0}} = \frac{1}{2 \sinh(\beta \hbar \omega_0 / 2)}$$

Classical limit: $\beta \hbar \omega_0 \ll 1 \Rightarrow Z \approx \frac{1}{\beta \hbar \omega_0}$

$$Z_{cl} = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dp \int_{-\infty}^{\infty} dx e^{-\beta \left(\frac{p^2}{2m} + \frac{m\omega_0^2}{2} x^2 \right)} = \frac{1}{2\pi\hbar} \int_{-\infty}^{\infty} dp e^{-\frac{\beta p^2}{2m}} \int_{-\infty}^{\infty} dx e^{-\frac{\beta m\omega_0^2 x^2}{2}} \\ = \frac{1}{\beta \hbar \omega_0}$$

$\frac{1}{2\pi\hbar}$ in Z_{cl} gives the correct state counting
($\Delta p \Delta x = 2\pi\hbar$ per state)

6. Free energy

$$F = -T \ln Z$$

Classical description of harmonic oscillator:

$$F_{cl} = -T \ln Z_{cl} = -T \ln \left(\frac{T}{\hbar \omega_0} \right)$$

$$S = -\frac{\partial F}{\partial T} \quad (\text{entropy, as defined in thermodynamics})$$

$$S_{cl} = -\frac{\partial F_{cl}}{\partial T} = 1 + \ln \left(\frac{T}{\hbar \omega_0} \right) \quad \text{problem at } T \rightarrow 0!$$

full quantum result for oscillator (with $Z = \sum_n e^{-\beta \epsilon_n}$):

$$F = \frac{1}{2} \hbar \omega_0 + T \ln(1 - e^{-\beta \hbar \omega_0})$$

$$S = -\ln(1 - e^{-\frac{\hbar \omega_0}{T}}) + \frac{\frac{\hbar \omega_0}{T}}{e^{\frac{\hbar \omega_0}{T}} - 1}$$

$$S \propto \frac{\hbar \omega_0}{T} e^{-\frac{\hbar \omega_0}{T}} \quad \text{at } T \rightarrow 0$$

Entropy (the density matrix definition):

$$S = \ln [\text{number of available states}] = - \sum_n w_n \ln w_n;$$

w_n : is the probability of state n occupation

Yields the thermodynamic def. of S for Gibbs distr. w_n

7. Expectation values

\hat{O} : operator of a (measurable) quantity
expectation value for a given pure state: $\langle \psi | \hat{O} | \psi \rangle$

i.e., average of \hat{O} over a state $|\psi\rangle$

In the eigenstates of a Hamiltonian representation: $\Psi = \sum_n \alpha_n |n\rangle$

$$\langle \psi | \hat{O} | \psi \rangle = \sum_{mn} \alpha_m^* \alpha_n \underbrace{\langle m | \hat{O} | n \rangle}_{\substack{\uparrow \\ \text{matrix element of } \hat{O} \text{ on the eigenstate basis}}} \equiv \sum_{mn} \alpha_m^* \alpha_n \underbrace{O_{mn}}_{\substack{\uparrow \\ \Psi = \sum_n \langle n | \alpha_n^*}}$$

Time - averaged expectation value for a pure state:

$$\Psi = \sum_n \alpha_n |n\rangle \Rightarrow \Psi(t) = \sum_n \alpha_n e^{-i\epsilon_n t/\hbar} |n\rangle$$

$$\langle \Psi(t) | \hat{O} | \Psi(t) \rangle = \sum_{nm} \alpha_m^* \alpha_n e^{i \frac{\epsilon_m - \epsilon_n}{\hbar} t} O_{mn}$$

time - averaged value : $\bar{A} = \lim_{t_0 \rightarrow \infty} \frac{1}{t_0} \int_0^{t_0} dt A(t)$

$$\overline{\langle \Psi(t) | \hat{O} | \Psi(t) \rangle} = \sum_{nm} \alpha_m^* \alpha_n e^{i \frac{\epsilon_m - \epsilon_n}{\hbar} t} O_{mn} = \sum_n |\alpha_n|^2 O_{nn}$$

In general, depends on the initial state (via α_n).

ETH states that for a large (infinite in the limit) system and a local quantity \hat{O} , $|\alpha_n|^2$ depend only on the state's energy E_n

Expectation value of \hat{O} for a mixed state described by a density matrix

Return for a minute to the def. of expectation value for a one-particle system and write it out using the real-space coordinate rep.:

$$\langle \psi | \hat{O} | \psi \rangle = \int d\vec{r} \psi^*(\vec{r}) \hat{O}(\vec{r}, \vec{p}) \psi(\vec{r})$$

Now suppose we have a many-particle system in a pure state, $\Psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N)$, but still are interested in the expectation value of an observable associated with one specific particle, namely particle 1: $\hat{O}(\vec{r}_1, \vec{p}_1)$

We will start now from the already introduced definition of the expectation value, $\langle \Psi | \hat{O} | \Psi \rangle$, and see how to abbreviate it using the notion of the defined earlier density matrix $\rho(\vec{r}, \vec{r}')$,

$$\rho(\vec{r}_i, \vec{r}_i) \equiv \left(\prod_{i=2}^N \int d\vec{r}_i \right) \psi^*(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N).$$

Indeed,

$$\begin{aligned} \langle \Psi | \hat{O} | \Psi \rangle &= \int d\vec{r}_1 \int \prod_{i=2}^N d\vec{r}_i \psi^*(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \hat{O}(\vec{r}_1, \hat{p}_1) \psi(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \\ &= \int d\vec{r}_1 \int \prod_{i=2}^N d\vec{r}_i \psi^*(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \hat{O}(\vec{r}_1, \hat{p}_1) \int d\vec{r}'_1 \delta(\vec{r}_1 - \vec{r}'_1) \psi(\vec{r}'_1, \vec{r}_2, \dots, \vec{r}_N) \\ &= \int d\vec{r}'_1 \int d\vec{r}_1 \int \prod_{i=2}^N d\vec{r}_i \psi^*(\vec{r}_1, \vec{r}_2, \dots, \vec{r}_N) \psi(\vec{r}'_1, \vec{r}_2, \dots, \vec{r}_N) \hat{O}(\vec{r}_1, \hat{p}_1) \delta(\vec{r}_1 - \vec{r}'_1) \\ &= \int d\vec{r}'_1 \int d\vec{r}_1 \rho(\vec{r}_1, \vec{r}'_1) \hat{O}(\vec{r}_1, \hat{p}_1) \delta(\vec{r}_1 - \vec{r}'_1) \quad (\text{vector signs omitted}) \end{aligned}$$

Now we can use a suitable complete basis of single-particle wave functions:

$$\rho(r, r') = \sum_{nm} w_{mn} \psi_n^*(r) \psi_m(r') ; \quad \hat{p} = \sum_{nm} \langle nm | w_{mn} | n \rangle$$

Substitution of $\rho(r, r')$ in this form to the last line above yields

$$\langle \Psi | \hat{O} | \Psi \rangle = \int d\vec{r}'_1 \int d\vec{r}_1 \sum_{nm} w_{mn} \psi_n^*(r_1) \psi_m(r'_1) \hat{O}(r_1, \hat{p}_1) \delta(r_1 - r'_1)$$


Noting that \hat{p}_1 acts on r_1 but not on r'_1 we may re-write

$$\begin{aligned} \langle \Psi | \hat{O} | \Psi \rangle &= \int d\vec{r}_1 \sum_{nm} w_{mn} \psi_n^*(r_1) \hat{O}(r_1, \hat{p}_1) \int d\vec{r}'_1 \delta(r_1 - r'_1) \psi_m(r'_1) \\ &= \sum_{nm} w_{mn} \underbrace{\int d\vec{r}_1 \psi_n^*(r_1) \hat{O}(r_1, \hat{p}_1) \psi_m(r_1)}_{D_{nm}} = \sum_{nm} w_{mn} D_{nm} \end{aligned}$$

$$\langle\langle \hat{O} \rangle\rangle \equiv \sum_{nm} \omega_{mn} O_{nm} = \sum_m \sum_n \omega_{mn} O_{nm} = \text{Tr}(\hat{\rho} \hat{O})$$

Unless the function $\Psi(\bar{r}_1, \bar{r}_2, \dots, \bar{r}_N)$ is factorizable, $\Psi(\bar{r}_1, \bar{r}_2, \dots, \bar{r}_N) = \psi(\bar{r}_1) \cdot \psi(\bar{r}_2, \dots, \bar{r}_N)$, the density matrix $\hat{\rho}$ corresponds to a mixed state of particle 1. The density matrix definition of the expectation value of $\hat{O}(r_1, \hat{\rho}_1)$ is agnostic wrt states of other particles.

Within ETH, a thermal ensemble average of a local quantity can be obtained as an expectation value over a pure state with energy $\propto T \cdot N$ of a large system ($N \gg 1$).



We may consider a system with particle 1 confined to volume V , while other $N-1$ particles reside in volume V_{out} . Assuming that particle 1 interacts with other particles, but that interaction is weak compared to single-particle energies E_m and

assuming the ETH is valid, we may use

$$\omega_{mn} = \frac{1}{Z} e^{-\beta E_m} \delta_{mn} \quad (\text{Gibbs distr.}) \text{ to evaluate } \langle\langle \hat{O} \rangle\rangle:$$

$$\langle\langle \hat{O} \rangle\rangle = \frac{1}{Z} \sum_m e^{-\beta E_m} O_m = \frac{1}{Z} \text{Tr}(e^{-\beta \hat{H}} \hat{O})$$

In the conventional language, this is operator \hat{O} averaged over the Gibbs (i.e. thermal) ensemble,

$$\hat{P}_G = \frac{1}{Z} \cdot \sum_m \ln m e^{-\beta E_m} \langle m | \hat{O} | m \rangle$$