

Modernizing the Diagnosis and Treatment of Gender Dysphoria: A Neurobiologically Grounded Policy Framework

Joseph Ben-Simon

March 2025

© 2025 Joseph Ben-Simon™

© 2025 Joseph Ben-Simon™

All rights reserved. No part of this publication may be reproduced, distributed, or transmitted
in any form or by any means—
including photocopying, recording, or other electronic or mechanical methods—
without prior written permission of the author, except in the case of brief quotations used for
critical review or scholarly analysis.

For permissions, contact: GDFPolicyFeedback@bensimonlabs.com

DOI: 10.5281/zenodo.15098234

Author Attribution and Registration

Title:

Modernizing the Diagnosis and Treatment of Gender Dysphoria: A Neurobiologically Grounded Policy Framework

Author:

Joseph Ben-Simon

Affiliation:

Independent Researcher and Policy Consultant

Position:

Neuroevolutionary Psychologist, Policy Advisor

Email:

GDFPolicyFeedback@bensimonlabs.com

Date of Completion:

March 2025

Version:

Final Public Policy Draft (v1.0)

Status:

Submitted for institutional review and policy consultation

DOI: 10.5281/zenodo.15098234

Executive Summary

The current clinical and policy frameworks for diagnosing and treating gender dysphoria rely predominantly on self-reported identity and psychological assessments. While personal identity is a critical dimension of patient care, the absence of objective biological criteria introduces substantial risks across medical, legal, and insurance domains. These include the potential for misdiagnosis, premature or unnecessary medicalization, and public health policy decisions that lack empirical grounding.

This white paper proposes a modernized, neurobiologically grounded diagnostic and treatment framework for gender dysphoria. Drawing from a growing body of neuroscientific research, we identify consistent morphological and functional differences in specific brain regions associated with persistent gender identity incongruence. These findings provide a foundation for the development of standardized diagnostic criteria that are verifiable, reproducible, and developmentally appropriate.

Central to this proposal is the requirement that irreversible gender-affirming medical interventions—such as hormone therapy or surgery—be authorized only for adults who have demonstrated stable neuroanatomical markers over a longitudinal period. Specifically, eligibility would require:

- A minimum age of 18 years;
- Evidence of completed neurodevelopment in gender-related brain regions;
- At least three years of consistent neuroimaging data demonstrating morphological alignment with reported gender identity;
- Final evaluation by a multidisciplinary panel including specialists in neuropsychology, psychiatry, endocrinology, and neurology.

This framework aims to reduce the risks associated with overdiagnosis and premature intervention, while preserving access to medically necessary care for individuals whose identities are substantiated by durable neurological evidence.

By incorporating objective neurobiological standards into the diagnostic and treatment process, this policy model enhances clinical accuracy, supports insurance accountability, and protects both patient welfare and institutional integrity. It offers a balanced, non-ideological path forward for public health systems, medical boards, and policymakers navigating the complexities of gender-affirming care.

1 Introduction

The diagnosis and treatment of gender dysphoria have undergone rapid transformation over the past two decades. Shifting cultural attitudes, growing visibility of transgender individuals, and evolving standards of care have led to broader access to gender-affirming medical interventions. However, this acceleration has not been matched by the development of rigorous diagnostic criteria grounded in neuroscience or developmental biology.

Current diagnostic models rely heavily on subjective self-identification and psychosocial assessment. While these factors are essential components of understanding a person's lived experience, they are insufficient as the sole basis for initiating permanent medical interventions, such as hormone replacement therapy or surgical transition. In nearly every other domain of medicine—particularly those involving lifelong or irreversible treatments—diagnoses are confirmed through objective, reproducible biological data.

In parallel, the field of neuroscience has matured to the point where structural and functional brain differences related to gender identity can be observed and measured. Emerging research identifies distinct patterns in specific brain regions that are more aligned with an individual's experienced gender than their sex assigned at birth. These include areas involved in hormonal regulation, interoception, body image, and emotional processing. This growing body of evidence makes it both feasible and necessary to incorporate neuroimaging into the diagnostic process for gender dysphoria.

At the same time, ethical and developmental concerns have intensified regarding the provision of irreversible treatments to minors and individuals in early stages of identity formation. Adolescence is a period marked by neuroplasticity, heightened suggestibility, and often unresolved mental health challenges. Introducing permanent medical pathways during this time—without biological confirmation—carries significant risks, both for the individual and the institutions that authorize treatment.

This white paper proposes a new diagnostic framework that integrates neuroscientific evidence into clinical and policy standards. It calls for the use of neuroimaging to confirm gender dysphoria in adults whose brain development has stabilized, and whose brain morphology shows consistent alignment with their experienced gender over a three-year period. This approach balances individual autonomy with developmental caution, offering a scientifically grounded, ethically defensible model for diagnosis, treatment, and healthcare policy.

2 Current Challenges

The increasing accessibility of gender-affirming care has exposed critical weaknesses in the existing diagnostic and treatment infrastructure. These challenges are not rooted in malice or neglect, but in the rapid pace of social change outstripping the development of clinical standards. Without a robust biological framework, both patients and institutions remain vulnerable to misdiagnosis, overreach, and inconsistent policy enforcement.

2.1 Diagnostic Overreach

Present diagnostic guidelines, such as those in the DSM-5, emphasize subjective experience and psychological discomfort as primary indicators of gender dysphoria. While these factors are important in understanding personal identity, they are insufficient on their own to justify lifelong medical intervention.

In contrast to most chronic or irreversible medical conditions, the diagnosis of gender dysphoria currently lacks a requirement for objective biomarkers or physiological evidence. This diagnostic looseness can lead to over-classification—treating individuals whose distress may stem from unrelated sources, including trauma, developmental confusion, or social influence.

2.2 Risk of Misdiagnosis and Premature Intervention

The combination of subjective diagnostics and the rising social visibility of gender transition pathways has created an environment where some individuals may be fast-tracked into irreversible interventions without adequate developmental, psychological, or neurological assessment.

Adolescents are particularly vulnerable. During this critical period of neurodevelopment, identity formation is fluid, emotional regulation is still maturing, and the influence of peers and social media is exceptionally high. Without longitudinal data or biological corroboration, some individuals may receive life-altering treatments that they later come to regret—raising ethical concerns about informed consent, autonomy, and clinical responsibility.

2.3 Policy Vulnerability

Public institutions, insurance providers, and national health systems are now tasked with making high-stakes decisions about access to gender-affirming care—often in the absence

of scientifically robust diagnostic criteria. This creates an unstable policy environment where decisions may be driven more by ideology, political pressure, or litigation risk than by clinical evidence.

In such a vacuum, institutions face a lose-lose scenario: either approve all claims, regardless of medical necessity, or risk being accused of discrimination for applying scrutiny. A standardized, neurobiologically informed diagnostic model would give policymakers a defensible, evidence-based toolset to guide access, oversight, and long-term planning.

3 Scientific Basis for a Neurobiological Framework

Advances in neuroscience over the past two decades have revealed that gender identity is not exclusively a psychosocial construct, but in many cases has measurable neurobiological correlates. Structural and functional neuroimaging studies have identified specific brain regions that differ in individuals experiencing gender dysphoria, often aligning more closely with their experienced gender than their sex assigned at birth (Frigerio et al., 2021).

These findings provide a critical opportunity: to elevate the diagnosis of gender dysphoria from a purely subjective classification to a condition supported by identifiable, reproducible neuroanatomical patterns. The following brain regions have been most consistently implicated in the emerging literature.

3.1 Hypothalamus

Function: Regulation of hormonal balance, sexual behavior, and homeostasis.

Findings: Recent neuroimaging studies have identified structural differences in the hypothalamus among transgender individuals, suggesting a neurobiological basis for gender identity (Frigerio et al., 2021). Additionally, developmental research indicates significant interactions between treatment status, sex, and gender, with notable volume changes observed in the hypothalamus over time (Burke et al., 2021).

Implication: These findings underscore the hypothalamus's critical role in gender identity and its potential involvement in the experience of gender dysphoria.

3.2 Inferior Front-Occipital Fasciculus (IFOF)

Function: Facilitates communication between the frontal and occipital lobes, playing a role in visual processing and cognitive functions.

Findings: Diffusion tensor imaging studies have revealed that transgender individuals exhibit unique white matter microstructures in the IFOF, aligning more closely with their

experienced gender rather than their sex assigned at birth. (Burke et al., 2017)

Implication: These findings suggest that the IFOF may serve as a neural correlate of gender identity, independent of sexual orientation.

3.3 Insular Cortex (Insula)

Function: Involved in interoception, self-awareness, and emotional regulation.

Findings: Recent neuroimaging studies have identified variations in gray matter volume and altered functional connectivity in the insular cortex among transgender individuals. For instance, increased gray matter volumes in the right insular and inferior frontal cortex have been observed in male-to-female transgender individuals compared to cisgender controls (Savic & Arver, 2011). Additionally, changes in functional connectivity involving the insular cortex have been noted following gender-affirming hormone therapy (Reed et al., 2023).

Implication: These findings suggest that structural and functional variations in the insular cortex may contribute to altered body perception and the persistent discomfort with sex-assigned anatomy experienced in gender dysphoria.

3.4 Anterior Cingulate Cortex and Ventral Striatum

Function: Involved in cognitive control, decision-making, and reward processing.

Findings: The anterior cingulate cortex and ventral striatum have been implicated in the experience of cognitive dissonance related to gender identity. (Altinay & Anand, 2020)

Implication: Suggests that gender dysphoria may involve neural mechanisms associated with conflict monitoring and reward evaluation.

3.5 Precentral Gyrus (Primary Motor Cortex)

Function: Responsible for voluntary motor control and movement planning.

Findings: Neuroimaging studies have identified structural differences in the precentral gyrus among transgender individuals. For instance, smaller gray and white matter volumes in the precentral gyrus have been observed in male-to-female transgender individuals compared to heterosexual women (Savic & Arver, 2011). Additionally, differential activations in the precentral gyrus have been noted in homosexual males during specific tasks (Frigerio et al., 2021).

Implication: These findings suggest that structural variations in the precentral gyrus may relate to embodied motor behavior and gendered movement patterns, contributing to the understanding of gender dysphoria.

3.6 Superior Parietal Lobule (SPL)

Function: Plays a crucial role in spatial orientation, body image integration, and coordination of sensory information.

Findings: Neuroimaging studies have identified notable differences in the SPL among transgender individuals. Increased centrality in the primary somatosensory cortex and the SPL has been observed in transgender individuals (Lin et al., 2020). Additionally, greater intrinsic connectivity in body representation areas, including the SPL, has been reported (Manzouri & Savic, 2017).

Implication: These findings suggest that structural and functional variations in the SPL may contribute to disrupted body image and persistent discomfort with sex-assigned anatomy, characteristic of gender dysphoria.

3.7 White Matter Microstructure

Function: Neural communication between brain regions.

Findings: Diffusion tensor imaging (DTI) studies have revealed that white matter tracts in transgender individuals often resemble those of their experienced gender. For instance, mean diffusivity in specific white matter regions differs between trans individuals and cisgender controls, aligning more closely with experienced gender (Kranz et al., 2014). In youth with gender dysphoria, MRI studies have similarly found sex-atypical white matter organization even before the onset of puberty (Bakker, 2017).

Implication: These findings reinforce the hypothesis that gender identity is embedded in global brain architecture and not confined to localized anatomical features.

3.8 Default Mode Network (DMN) and Salience Network

Function: Self-referential thought (DMN) and prioritization of salient stimuli (Salience Network).

Findings: Neuroimaging studies have identified altered functional connectivity patterns within these networks in individuals with gender dysphoria. For example, disruptions in the DMN and SN have been associated with differences in own-body perception (Feusner et al., 2017). Additionally, models propose that these network alterations contribute to cognitive dissonance experienced in gender dysphoria (Altinay & Anand, 2020).

Implication: These findings suggest that core mechanisms of identity processing are neurologically aligned with gender identity.

3.9 Amygdala and Limbic System

Function: Emotional regulation, fear response, and stress processing.

Findings: Neuroimaging studies have identified variations in the amygdala and limbic system among individuals with gender dysphoria. For instance, structural differences in these regions have been associated with gender identity and sexual orientation (Frigerio et al., 2021). Additionally, models propose that the limbic system plays a crucial role in processing the cognitive dissonance experienced in gender dysphoria (Altinay & Anand, 2020).

Implication: These findings suggest a neurobiological component to the emotional distress often observed in untreated or unsupported gender dysphoria.

4 Proposed Diagnostic and Treatment Framework

Given the growing body of neuroscientific evidence supporting a biological basis for gender identity, it is now possible to establish a diagnostic framework that incorporates objective data, protects patients, and provides clear criteria for access to irreversible medical interventions.

This section outlines a proposed model in which medical eligibility for gender-affirming treatment is contingent upon demonstrable neurodevelopmental stability and verified brain morphology consistent with the individual's experienced gender. This model does not prohibit care, but instead ensures that care is grounded in long-term biological patterns rather than temporary psychological states or social influences.

4.1 Eligibility Criteria for Irreversible Medical Intervention

To reduce the risks of overdiagnosis, premature treatment, and long-term regret, the following eligibility conditions are proposed:

- 1. Minimum Age of 18 Years:**

All candidates for irreversible gender-affirming treatments must be legal adults. This aligns with established neurodevelopmental timelines, which show that core executive functions—including cognitive control, working memory, and risk evaluation—reach adult-like stability around this age (Luna et al., 2015; Paulsen et al., 2015).

- 2. Neurodevelopmental Stability:**

Neuroimaging must confirm that the relevant brain regions associated with gender identity—particularly the hypothalamus, insula, anterior cingulate cortex, and white

matter networks—have completed maturation. This ensures that treatment is not initiated during ongoing neural remodeling (Paus, 2024; Shaw et al., 2008).

3. **Three-Year Longitudinal Neuroimaging Record:**

The individual must present three consecutive years of consistent neuroimaging data (e.g., MRI, fMRI, DTI) demonstrating stable morphological alignment with their reported gender identity, without regression or ambiguity in brain structure development (Frigerio et al., 2021; Kreukels & Guillamon, 2016).

4. **Multidisciplinary Review Panel:**

A formal diagnosis and treatment clearance must be issued by a qualified panel composed of specialists in neuropsychology, psychiatry, endocrinology, and neurology. This team is responsible for evaluating imaging data, developmental history, comorbid conditions, and psychological readiness (Coleman et al., 2012; Hembree et al., 2017).

5. **Ethical and Medical Standards for Irreversible Intervention:**

Irreversible gender-affirming interventions must comply with established ethical and medical guidelines. These standards emphasize comprehensive evaluations, informed consent, and individualized care to ensure patient safety and autonomy (Coleman et al., 2012; Hembree et al., 2017; Spencer & Spencer, 2016).

4.2 **Scope of Application**

This framework applies specifically to irreversible gender-affirming interventions—defined as treatments that permanently alter physiology, anatomy, or legal classification in ways not easily reversed. These include:

- Hormone replacement therapy (HRT) with permanent physiological effects
- Surgical procedures altering primary or secondary sex characteristics
- Legal or medical reclassification that affects access to sex-specific services or spaces

Supportive therapies, counseling, and non-invasive forms of care remain fully accessible to all individuals, regardless of age or imaging status. Diagnostic safeguards must never impede access to emotional support, psychosocial stability, or gender exploration.

These protections are consistent with medical and legal precedents in other domains. For example, elective sterilization in individuals under 25 is often discouraged or restricted due to the irreversible nature of the procedure and risk of future regret (American College of Obstetricians and Gynecologists, 2017; U.S. Department of Health and Human Services, 2023). Such age-based caution reflects a broader ethical principle: permanent medical

interventions should only proceed when identity stability and neurodevelopmental maturity are sufficiently demonstrated.

All decisions must occur within a voluntary, informed process, with full respect for patient autonomy and multidisciplinary review.

4.3 Rationale for Developmental Safeguards

These safeguards reflect a commitment to medical ethics, evidence-based care, and long-term patient protection. By requiring neurologically confirmed, stable gender identity prior to irreversible intervention, the framework mitigates the risk of overmedicalization, misdiagnosis, and treatment regret—especially in vulnerable or developmentally immature populations.

This approach aligns with the standards set forth by major clinical organizations, including the World Professional Association for Transgender Health (WPATH), the Endocrine Society, and the American Medical Association, all of which emphasize informed consent, multidisciplinary review, and diagnostic clarity before proceeding with permanent treatments (Coleman et al., 2012; Hembree et al., 2017; Spencer & Spencer, 2016).

In addition to protecting patients, these criteria offer medical institutions and insurers a defensible standard for clinical judgment and resource allocation. They reduce liability exposure, clarify the threshold for irreversible care, and bolster public trust in the integrity of gender-affirming healthcare systems.

5 Policy and Insurance Recommendations

The implementation of a neurobiologically grounded diagnostic framework for gender dysphoria offers substantial benefits for insurers, healthcare providers, and policymakers. By defining objective thresholds for irreversible interventions, this model strengthens clinical accountability, minimizes legal and administrative liability, and safeguards patient autonomy through scientifically validated protocols.

These recommendations align with established legal and medical precedents. For example, federal regulations governing sterilization in federally funded programs require enhanced informed consent protections for individuals under 21 (U.S. Department of Health and Human Services, 2023), and medical ethics organizations such as the American College of Obstetricians and Gynecologists advise caution around elective sterilization in individuals under 25 due to risks of long-term regret and inadequate future self-projection (American College of Obstetricians and Gynecologists, 2017).

From an institutional perspective, the use of objective neurodevelopmental criteria also provides insurers and medical boards with a clear standard for clinical eligibility. This reduces the risk of malpractice claims, regulatory challenges, or public trust erosion arising from allegations of negligent or premature intervention (Spencer & Spencer, 2016).

The following recommendations outline actionable steps for aligning institutional policies with emerging neuroscientific standards in gender-affirming care.

5.1 5.1 Insurance Providers

- **Require objective verification of diagnosis:** Authorize irreversible gender-affirming interventions only when supported by neuroimaging data and multidisciplinary panel review, as outlined in Section 4. This aligns with the World Professional Association for Transgender Health (WPATH) Standards of Care, which emphasize comprehensive assessments prior to medical interventions (World Professional Association for Transgender Health, 2012).
- **Differentiate between psychological distress and neurologically substantiated dysphoria:** Align coverage decisions with biological evidence to avoid approving treatments for individuals whose distress may stem from other psychiatric or developmental causes. Incorporating neurobiological assessments can aid in distinguishing genuine gender dysphoria from other conditions (Altinay & Anand, 2020).
- **Standardize documentation requirements:** Ensure that submitted claims include longitudinal imaging reports, panel findings, and neurodevelopmental evaluations to validate medical necessity. Establishing clear documentation standards is consistent with practices outlined by major healthcare providers (Permanente, 2025).
- **Implement fraud prevention measures:** Reduce vulnerability to ideologically driven or misrepresented claims by enforcing evidence-based eligibility criteria. Adherence to standardized protocols, such as those recommended by WPATH, can help mitigate the risk of fraudulent claims (World Professional Association for Transgender Health, 2012).

5.2 5.2 Medical Boards and Institutions

- **Revise clinical guidelines:** Update diagnostic and treatment protocols to consider emerging neurobiological research related to gender dysphoria. While neuroimaging studies have provided insights into potential neurobiological markers, their clinical application remains exploratory and should be integrated cautiously, ensuring that

such assessments are conducted within a comprehensive, multidisciplinary framework (Altinay & Anand, 2020).

- **Establish ethical review structures:** Create internal panels to review cases involving minors, complex co-morbidities, or contested diagnoses, ensuring multidisciplinary oversight. Such structures are essential to navigate the ethical complexities inherent in gender-affirming care for youth (Kimberly et al., 2018).
- **Train clinicians in comprehensive assessment protocols:** Ensure that practitioners involved in gender-affirming care are trained to conduct thorough evaluations, integrating psychological assessments and, where appropriate, considering neurobiological research findings. This approach aligns with the Endocrine Society's emphasis on comprehensive assessments involving qualified mental health professionals (Hembree et al., 2017).
- **Protect exploratory care pathways:** Safeguard access to psychological counseling and non-invasive support for individuals who may not pursue medical interventions, ensuring that all individuals receive appropriate care tailored to their unique needs.

5.3 5.3 Public Health Policymakers

- **Codify objective diagnostic standards:** Incorporate neurobiological criteria into public health regulations and legal frameworks governing gender-affirming treatment eligibility to ensure consistency and scientific rigor.
- **Ensure legal consistency across jurisdictions:** Prevent forum-shopping and medical tourism by encouraging national or supranational alignment around evidence-based treatment thresholds. Disparities in state laws regarding gender-affirming care can lead individuals to seek treatment in jurisdictions with more permissive regulations, underscoring the need for standardized policies (FindLaw, 2025).
- **Protect institutions from politicization:** Provide legal support to medical boards, schools, and insurers implementing science-based eligibility frameworks, insulating them from ideological or activist pressure. Several states have enacted shield laws to protect healthcare providers offering gender-affirming care from out-of-state legal actions, highlighting the importance of legal safeguards in maintaining the integrity of medical institutions (Williams Institute, 2025).
- **Mandate data transparency and outcomes reporting:** Require institutions that provide gender-affirming care to collect and publish anonymized data on long-term outcomes, treatment efficacy, and neurodevelopmental correlates. Implementing patient-reported outcome measures (PROMs) can enhance data transparency and inform best practices in gender-affirming treatments (AMA Journal of Ethics, 2023).

This integrated policy approach balances the right to individualized care with the responsibility to ground that care in measurable, developmental, and biological standards. It enables systems-level reform while preserving the dignity, autonomy, and safety of all patients—ensuring that care is both ethically sound and scientifically justified.

6 Ethical and Legal Considerations

The shift toward neurobiologically informed diagnostic standards for gender dysphoria introduces complex ethical and legal questions. These considerations are not secondary to scientific or clinical reasoning—they are foundational to any responsible model of care.

This section outlines how the proposed framework upholds core principles of medical ethics, strengthens legal protections for both patients and practitioners, and promotes a system of care that respects individual rights without compromising clinical accountability or public trust.

6.1 6.1 Respect for Autonomy

The proposed framework affirms the right of individuals to identify with a gender different from their sex assigned at birth. However, in the context of irreversible medical treatment, autonomy must be paired with the principle of informed consent.

Requiring objective evidence of neurodevelopmental maturity and morphological consistency does not negate personal identity—it ensures that treatment decisions are made with a full understanding of their permanence, biological underpinnings, and long-term consequences. This is especially critical when treating young adults, who may be vulnerable to psychological distress or external pressures during identity formation. Research indicates that adolescent brain development continues into early adulthood, affecting decision-making capacities and highlighting the importance of assessing neurodevelopmental maturity in medical contexts (Weithorn, 2020). Additionally, developmental and neuroscientific studies underscore the necessity of evaluating cognitive maturity when considering irreversible medical interventions in minors (Hein et al., 2015). In the realm of gender-affirming care, implementing informed consent models that balance respect for patient autonomy with comprehensive understanding of treatment implications is vital (Cavanaugh et al., 2016).

6.2 6.2 Protection from Harm

The principle of non-maleficence—"first, do no harm"—is a cornerstone of medical ethics. Irreversible interventions, such as hormone therapy or surgery, carry physical, psychological, and reproductive consequences. When such treatments are initiated without adequate diagnostic precision, the potential for regret, complications, or iatrogenic harm increases significantly.

A neurobiologically grounded approach reduces this risk by ensuring that only those with demonstrable, stable dysphoria receive these interventions, and that others are offered supportive care pathways without undergoing unnecessary or harmful procedures. Studies have shown that regret rates after gender-affirming surgeries are approximately 1%, which is lower than regret rates for other surgical procedures such as prostatectomy and bariatric surgery (Narayan et al., 2023). Additionally, research indicates that the vast majority of transgender and nonbinary youth report satisfaction with gender-affirming medical care, with minimal regret, emphasizing the importance of thorough diagnostic assessments (Tordoff et al., 2022). Understanding the neurobiological underpinnings of gender dysphoria can further enhance diagnostic precision and inform treatment approaches that minimize the risk of harm (Gliske, 2019).

6.3 6.3 Developmental Safeguards for Minors

Minors and adolescents are in a critical period of cognitive, emotional, and neurological development. Their capacity to fully understand the long-term implications of gender-affirming medical treatments is limited, even in the presence of significant distress. Research indicates that adolescents may have diminished decision-making abilities due to ongoing neurocognitive development, which can affect their ability to comprehend the consequences of irreversible medical interventions (Ravindranath et al., 2024).

This framework upholds the legal and ethical standard that irreversible medical interventions should be deferred until developmental maturity is reached and sustained. It does not block access to care—it distinguishes between supportive services (e.g., counseling, social affirmation) and irreversible procedures, offering protection while preserving dignity and support. Ethical considerations in gender-affirming care emphasize the importance of balancing respect for autonomy with non-maleficence, particularly when treating youth during critical developmental stages (Kimberly et al., 2018). The World Professional Association for Transgender Health (WPATH) recommends individualized assessments and careful consideration of developmental maturity before initiating irreversible medical interventions in adolescents (Coleman et al., 2022).

6.4 Institutional Integrity and Legal Defense

Medical institutions, insurers, and public agencies require defensible frameworks that shield them from legal liability and politicized decision-making. In the absence of clear, evidence-based diagnostic thresholds, these institutions risk exposure to litigation from both sides—those denied care and those who later regret it. Establishing objective diagnostic standards is essential to navigate the evolving legal landscape surrounding gender-affirming care (Herbst, 2025).

A model based on objective neuroimaging and longitudinal stability provides a scientifically and legally justifiable basis for care decisions. Studies have analyzed malpractice and liability cases related to gender-affirming surgeries over the past five decades, underscoring the importance of standardized diagnostic pathways to reduce institutional risk (Brozynski et al., 2024). Furthermore, recent neuroimaging-based models of gender dysphoria have proposed objective biological criteria that may serve as the basis for consistent, evidence-based decision-making (Altinay & Anand, 2020).

This framework offers a transparent, evidence-based standard for courts, regulatory bodies, and professional associations to assess the appropriateness of gender-affirming care on a case-by-case basis. By grounding treatment eligibility in longitudinal neurobiological evidence and clinical oversight, institutions can reduce exposure to malpractice claims and protect against politically motivated challenges (Brozynski et al., 2024). It strengthens institutional integrity while preserving individualized care, balancing legal defensibility with ethical obligation.

6.5 Avoiding Discrimination While Ensuring Rigor

This proposal does not seek to marginalize transgender individuals or invalidate their experiences. On the contrary, it aims to protect individuals from premature medicalization and ensure that those who require intervention receive it under conditions that are safe, defensible, and oriented toward long-term well-being. The American Psychological Association underscores the importance of individualized, evidence-based care tailored to the needs of transgender and gender-diverse individuals, emphasizing that inclusion and scientific rigor are not mutually exclusive (American Psychological Association, 2024).

Equal treatment does not imply uniform treatment. Across medicine—and especially within psychiatry and neurology—clinical standards routinely differentiate care based on objective biological markers, symptom severity, and prognosis. The application of biomarkers in psychiatric decision-making is a well-established method to ensure accurate diagnosis and individualized treatment (Strimbu & Tavel, 2020). Extending this principle to gender dysphoria allows clinicians to uphold the ethical standards of beneficence

and non-maleficence while safeguarding patient autonomy through clarity and evidence (Kimberly et al., 2018).

This framework therefore seeks not to limit access but to elevate standards, aligning gender-affirming care with best practices already in place across other domains of developmental and behavioral medicine.

7 Conclusion

The diagnosis and treatment of gender dysphoria represent one of the most ethically sensitive and clinically complex challenges in contemporary medicine. As access to gender-affirming interventions has expanded, so too has the responsibility to implement rigorous, evidence-based standards that uphold both patient autonomy and institutional accountability.

This white paper has presented a neurobiologically grounded framework for diagnosing gender dysphoria—one that integrates findings from structural and functional neuroimaging, establishes clear eligibility thresholds, and mandates multidisciplinary oversight for irreversible interventions. By limiting such treatments to adults with stable, biologically substantiated gender incongruence documented over multiple years, this model offers a pathway that is scientifically defensible, ethically sound, and socially responsible.

The framework affirms individual identity while acknowledging the profound risks associated with premature or misdirected medicalization. It distinguishes between supportive care—which should remain accessible to all—and irreversible procedures, which warrant a higher evidentiary standard due to their permanent impact.

We call upon insurers, medical boards, research institutions, and public policymakers to adopt this framework as a foundation for systemic reform. Doing so will not only safeguard vulnerable populations, but also restore public confidence in the integrity of gender-related healthcare, enable sustainable policy development, and elevate diagnostic standards to the level expected in all other domains of long-term medical intervention.

A future of affirming, evidence-based, and ethically defensible care is possible—but it requires that institutions lead with science, not ideology, and that they prioritize developmental caution over political expediency.

References

- Altinay, M., & Anand, A. (2020). Neuroimaging gender dysphoria: A novel psychobiological model. *Brain Imaging and Behavior*, 14(4), 1281–1297. <https://doi.org/10.1007/s11682-019-00121-8>
- AMA Journal of Ethics. (2023). Patient-reported outcome measures in gender-affirming surgery. *AMA Journal of Ethics*. <https://journalofethics.ama-assn.org/article/patient-reported-outcome-measures-gender-affirming-surgery/2023-06>
- American College of Obstetricians and Gynecologists. (2017). Sterilization of women: Ethical issues and considerations [Available at <https://www.acog.org/clinical/clinical-guidance/committee-opinion/articles/2017/04/sterilization-of-women-ethical-issues-and-considerations>]. *ACOG Committee Opinion No. 695*.
- American Psychological Association. (2024). Apa policy statement on affirming evidence-based inclusive care for transgender and gender diverse individuals. <https://www.apa.org/about/policy/transgender-nonbinary-inclusive-care.pdf>
- Bakker, J. (2017). Mri reveals structural brain differences in young people who may be transgender [Available at <https://healthimaging.com/topics/medical-imaging/neuroimaging/mri-reveals-structural-brain-differences-young-people-who-may>]. *Health Imaging*.
- Brozynski, M., Oleru, O., Seyidova, N., Rew, C., Nathaniel, S., & Taub, P. J. (2024). Litigation in gender affirming surgery: Reviewing five decades of the medical malpractice landscape. *Journal of Plastic, Reconstructive & Aesthetic Surgery*, 77(5), 1045–1052. <https://doi.org/10.1016/j.bjps.2024.04.036>
- Burke, S. M., Manzouri, A. H., & Savic, I. (2017). Structural connections in the brain in relation to gender identity and sexual orientation. *Scientific Reports*, 7(1), 17954. <https://doi.org/10.1038/s41598-017-17352-8>
- Burke, S. M., Manzouri, A. H., & Savic, I. (2021). Mapping hypothalamic development in transgender and cisgender youth. *Research Square*. <https://doi.org/10.21203/rs.3.rs-4883317/v1>
- Cavanaugh, T., Hopwood, R., & Lambert, C. (2016). Informed consent in the medical care of transgender and gender-nonconforming patients. *AMA Journal of Ethics*, 18(11), 1147–1155. <https://doi.org/10.1001/journalofethics.2016.18.11.sect1-1611>
- Coleman, E., Bockting, W., Botzer, M., & et al. (2012). Standards of care for the health of transsexual, transgender, and gender nonconforming people, version 7. *International Journal of Transgenderism*, 13(4), 165–232. <https://doi.org/10.1080/15532739.2011.700873>
- Coleman, E., Radix, A. E., Bouman, W. P., Brown, G. R., & et al. (2022). Standards of care for the health of transgender and gender diverse people, version 8. *International*

- Journal of Transgender Health*, 23(S1), S1–S259. <https://doi.org/10.1080/26895269.2022.2100644>
- Feusner, J. D., Lidström, A., Moody, T. M., Dhejne, C., Bookheimer, S., & Savic, I. (2017). Intrinsic network connectivity and own body perception in gender dysphoria. *Brain Imaging and Behavior*, 11(4), 964–976. <https://doi.org/10.1007/s11682-016-9578-6>
- FindLaw. (2025). State laws on gender-affirming care. <https://www.findlaw.com/lgbtq-law/state-laws-on-gender-affirming-care.html>
- Frigerio, A., Ballerini, L., & Valdés Hernández, M. d. C. (2021). Structural, functional, and metabolic brain differences as a function of gender identity or sexual orientation: A systematic review of the human neuroimaging literature. *Archives of Sexual Behavior*, 50(8), 3329–3352. <https://doi.org/10.1007/s10508-021-02005-9>
- Gliske, S. V. (2019). The neurobiological basis of gender dysphoria. *eNeuro*, 6(6). <https://doi.org/10.1523/ENEURO.0183-19.2019>
- Hein, I. M., Troost, P. W., Lindeboom, R., Benninga, M. A., Zwaan, C. M., van Goudoever, J. B., & Lindauer, R. J. L. (2015). Medical decision-making in children and adolescents: Developmental and neuroscientific aspects. *BMC Pediatrics*, 15, 285. <https://doi.org/10.1186/s12887-017-0869-x>
- Hembree, W. C., Cohen-Kettenis, P. T., Gooren, L., Hannema, S. E., Meyer, W. J., Murad, M. H., Rosenthal, S. M., Safer, J. D., Tangpricha, V., & T'Sjoen, G. G. (2017). Endocrine treatment of gender-dysphoric/gender-incongruent persons: An endocrine society clinical practice guideline. *Journal of Clinical Endocrinology & Metabolism*, 102(11), 3869–3903. <https://doi.org/10.1210/jc.2017-01658>
- Herbst, A. (2025). The future of gender-affirming care: Legal, ethical, and practical considerations for providers. *National Law Review*. <https://www.natlawreview.com/article/future-gender-affirming-care-legal-ethical-and-practical-considerations-providers>
- Kimberly, L. L., Folkers, K. M., Friesen, P., Sultan, S., Quinn, G. P., Bateman-House, A., & Parent, B. (2018). Ethical issues in gender-affirming care for youth. *Pediatrics*, 142(6), e20181537. <https://doi.org/10.1542/peds.2018-1537>
- Kranz, G. S., Hahn, A., Kaufmann, U., Küblböck, M., Hummer, A., Ganger, S., Seiger, R., Winkler, D., Swaab, D. F., Windischberger, C., Kasper, S., & Lanzenberger, R. (2014). White matter microstructure in transsexuals and controls investigated by diffusion tensor imaging. *Journal of Neuroscience*, 34(46), 15466–15475. <https://doi.org/10.1523/JNEUROSCI.2488-14.2014>
- Kreukels, B. P. C., & Guillamon, A. (2016). Neuroimaging studies in people with gender incongruence. *International Review of Psychiatry*, 28(1), 120–128. <https://doi.org/10.3109/09540261.2015.1113163>
- Lin, C.-y., Chou, K.-C., Yeh, H.-W., Tsai, Y.-T., Cheng, D.-S., Lin, C.-P., Tseng, W.-Y. I., Chen, J.-H., & Hsieh, C.-P. (2020). Brain network interactions in transgender

- individuals with gender dysphoria. *NeuroImage*, 211, 116613. <https://doi.org/10.1016/j.neuroimage.2020.116613>
- Luna, B., Marek, S., Larsen, B., Tervo-Clemmens, B., & Chahal, R. (2015). An integrative model of the maturation of cognitive control. *Annual Review of Neuroscience*, 38, 151–170. <https://doi.org/10.1146/annurev-neuro-071714-034054>
- Manzouri, A., & Savic, I. (2017). Intrinsic network connectivity and own body perception in gender dysphoria. *Brain Imaging and Behavior*, 11(4), 964–976. <https://doi.org/10.1007/s11682-016-9578-6>
- Narayan, S. K., Hontscharuk, R., Danker, S., & et al. (2023). A systematic review of patient regret after surgery: A common and understudied problem. *The American Journal of Surgery*. <https://doi.org/10.1016/j.amjsurg.2023.02.056>
- Paulsen, D. J., Hallquist, M. N., Geier, C. F., & Luna, B. (2015). Development of the default-mode network from childhood to adulthood: A systematic review and meta-analysis. *Developmental Cognitive Neuroscience*, 12, 92–106. <https://doi.org/10.1016/j.dcn.2015.01.002>
- Paus, T. (2024). Development and maturation of the human brain, from infancy to adolescence. In *Principles and advances in population neuroscience* (pp. 327–348, Vol. 68). Springer. https://doi.org/10.1007/7854_2024_514
- Permanente, K. (2025). Ur 65 gender-affirming procedures medical necessity criteria. <https://healthy.kaiserpermanente.org/content/dam/kporg/final/documents/health-plan-documents/coverage-information/clinical-review-gender-affirming-procedures-nw.pdf>
- Ravindranath, O., Perica, M. I., Parr, A. C., Ojha, A., McKeon, S. D., Montano, G., Ullendorff, N., Luna, B., & Edmiston, E. K. (2024). Adolescent neurocognitive development and decision-making abilities regarding gender-affirming care. *Developmental Cognitive Neuroscience*, 66, 101351. <https://doi.org/10.1016/j.dcn.2024.101351>
- Reed, M. B., Handschuh, P. A., Klöbl, M., Konadu, M. E., Kaufmann, U., Hahn, A., Kranz, G. S., Spies, M., & Lanzenberger, R. (2023). The influence of sex steroid treatment on insular connectivity in gender dysphoria. *Psychoneuroendocrinology*, 155, 106336. <https://doi.org/10.1016/j.psyneuen.2023.106336>
- Savic, I., & Arver, S. (2011). Sex dimorphism of the brain in male-to-female transsexuals. *Cerebral Cortex*, 21(11), 2525–2533. <https://doi.org/10.1093/cercor/bhr032>
- Shaw, P., Kabani, N. J., Lerch, J. P., Eckstrand, K., Lenroot, R., Gogtay, N., Greenstein, D., Clasen, L., Evans, A., Rapoport, J. L., Giedd, J. N., & Wise, S. P. (2008). Neurodevelopmental trajectories of the human cerebral cortex. *Journal of Neuroscience*, 28(14), 3586–3594. <https://doi.org/10.1523/JNEUROSCI.5309-07.2008>
- Spencer, K., & Spencer, M. (2016). Informed consent in the medical care of transgender and gender-nonconforming patients. *AMA Journal of Ethics*, 18(11), 1147–1155. <https://doi.org/10.1001/journalofethics.2016.18.11.sect1-1611>

- Strimbu, K., & Tavel, J. A. (2020). Biomarkers in psychiatry: Concept, definition, types and implications. *Frontiers in Psychiatry*, 11, 432. <https://doi.org/10.3389/fpsy.2020.00432>
- Tordoff, D. M., Wanta, J. W., Collin, A., & et al. (2022). Levels of satisfaction and regret with gender-affirming medical care in transgender and nonbinary youth. *JAMA Pediatrics*, 176(5), 477–482. <https://doi.org/10.1001/jamapediatrics.2022.0041>
- U.S. Department of Health and Human Services. (2023). Sterilization of persons in federally assisted family planning projects [Code of Federal Regulations 42 CFR Part 441 Subpart F].
- Weithorn, L. A. (2020). Adolescent brain development and medical decision-making. *Pediatrics*, 146(Supplement 1), S18–S24. <https://doi.org/10.1542/peds.2020-0818F>
- Williams Institute. (2025). Shield laws for reproductive and gender-affirming health care: A state-by-state guide. <https://williamsinstitute.law.ucla.edu/publications/shield-laws-fact-sheets/>
- World Professional Association for Transgender Health. (2012). Standards of care for the health of transsexual, transgender, and gender nonconforming people (7th). <https://www.wpath.org/publications/soc>