

SYSTEMATIC SEARCH FOR SINGULARITIES

SYSTEMATIC SEARCH FOR SINGULARITIES IN 3D EULER
FLOWS BASED ON THE VOIGT REGULARIZATION

BY

NOAH BENSLER, B.Sc.

A THESIS

SUBMITTED TO THE SCHOOL OF COMPUTATIONAL SCIENCE & ENGINEERING

AND THE SCHOOL OF GRADUATE STUDIES

OF MCMASTER UNIVERSITY

IN PARTIAL FULFILMENT OF THE REQUIREMENTS

FOR THE DEGREE OF

MASTERS OF COMPUTATIONAL SCIENCE & ENGINEERING

© Copyright by Noah Bensler, September 2026

All Rights Reserved

Masters of Computational Science & Engineering (2026) McMaster University
(School of Computational Science & Engineering) Hamilton, Ontario, Canada

TITLE: SYSTEMATIC SEARCH FOR SINGULARITIES IN 3D
 EULER FLOWS BASED ON THE VOIGT REGULAR-
 IZATION

AUTHOR: Noah Bensler
 B.Sc. (Computer Science & Astrophysics),
 University of Calgary, Calgary, Canada

SUPERVISOR: Dr. Bartosz Protas

NUMBER OF PAGES: [xvi](#), [76](#)

Lay Abstract

A lay abstract of not more 150 words must be included explaining the key goals and contributions of the thesis in lay terms that is accessible to the general public.

The Navier-Stokes equations model the behaviour of viscous incompressible fluids. The question of whether or not the Navier-Stokes equations, or the inviscid Euler equations, are a valid description of fluid mechanics remains unanswered. A singularity is the occurrence of an infinite derivative of the velocity at a single point in the fluid flow. If the flow forms a singularity from smooth initial conditions, then the equations are not a valid descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to not form singularities from smooth initial conditions. However, by decreasing the regularization parameter we approximate Euler flows. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations. **We present numerical evidence that, upon optimization, singularities can form in Euler flows corresponding to smooth initial conditions.**

Abstract

Abstract here (no more than 300 words). An abstract of not more than 300 words must be included and indicates the major emphasis of the thesis, new discoveries, and its contribution to knowledge. The style of abstract varies from discipline to discipline but the student should follow an appropriate style. This page must be numbered 'iv'.

The Navier-Stokes equations are a system of partial differential equations used to model the behaviour of viscous incompressible fluids. The question of whether or not unique classical solutions of the Navier-Stokes equations, and inviscid Euler equations, exist for all time, given smooth initial conditions, remains unanswered for the three-dimensional problem.

A singularity is the occurrence of an infinite-derivative of the velocity at a single point in the fluid flow. The formation of a singularity from smooth initial conditions would invalidate the equations as accurate descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to be globally well-posed from smooth initial conditions. However, by decreasing the regularization parameter the solutions of the Euler-Voigt equations converge

to those of the Euler equations. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations.

We present numerical evidence that, upon optimization, singularities can form in Euler flows corresponding to smooth initial conditions.

Acknowledgements

Acknowledgements go here. An expression of thanks for assistance given by the Supervisor of research and by others should be either set forth on a separate page or incorporated into the Preface (if there is one). These and all subsequent preliminary pages listed under (f), (g) and (h) must be numbered in lower case Roman numerals, i.e., ‘iv’, ‘v’, ‘vi’ etc.

Dr. Bartosz Protas, Dr. Xinyu Zhao, Dr. Jörn Davidsen, members of committee,
Parents

Dr. Xinyu Zhao for laying the groundwork for my thesis and for her help with the software.

I would like to thank Dr. Bartosz Protas for his support and guidance over the last two years.

I also want to thank Dr. Jörn Davidsen of the Complexity Science Group for supervising my undergraduate research and inspiring me to pursue computational physics.

my parents, even if they

Contents

Lay Abstract	iii
Abstract	iv
Acknowledgements	vi
Notation, and Abbreviations	xiii
Declaration of Academic Achievement	xv
1 Introduction	1
1.1 Singularity Formation in Models of 3D Fluid Flows	1
1.2 3D Euler-Voigt Equations	4
1.3 Outline of the Thesis	6
2 Properties of the Euler-Voigt Equations	7
2.1 Finite Time Blow-up Criterion	7
2.2 Initial Conditions	11
2.3 Scaling Symmetries of Euler and Euler-Voigt Systems	13
3 The Optimization Problem	16

3.1	Formulation of the Problem	16
3.2	Initial Guesses	17
3.3	Gradient of Objective Function	19
3.4	Riemannian Conjugate Gradient Method	23
4	Numerical Methods	28
4.1	Discretization	28
4.2	Energy Spectra and Validation of Numerical Resolution	32
4.3	Validation of the Gradient of the Objective Functional (3.32)	34
4.4	Determination of the parameter defining the Gevrey space	37
5	Results	39
5.1	Singularity Detection	40
5.2	Results Obtained Using the Initial Guesses	42
6	Conclusion	49
A	Function Spaces	50
B	Proof of Relation (2.25)	52
C	Rescaling the Euler equations	53
D	Rescaling the Euler-Voigt equations	55
E	Evaluation of $\ \eta\ _{\dot{H}^s}$ for the Taylor Green Vortex	57
F	Evaluation of $\ \nabla \times \eta\ _{L^\infty}$ for the Taylor Green Vortex	60

G	Linearization of the Euler-Voigt System	62
H	Derivation of the Adjoint System	64
I	Derivation of the Gâteaux Directional Differential of the Objective Functional	70

List of Figures

2.1	Maximum of vorticity $\ \boldsymbol{\omega}(t)\ _{L^\infty}$. Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), 1024^3 (blue squares).	15
3.1	Isosurfaces of vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{\eta}_{TGV}$ (a) magnitude $ \boldsymbol{\omega}(\boldsymbol{x}) $, and components (b) $ \omega_1 $, (c) $ \omega_2 $, and (d) $ \omega_3 $	19
3.2	Isosurfaces of vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{\eta}_S$ (a) magnitude $ \boldsymbol{\omega}(\boldsymbol{x}) $, and components (b) $ \omega_1 $, (c) $ \omega_2 $, and (d) $ \omega_3 $	20
4.1	$e(k, t)$ vs. k starting from $\boldsymbol{\eta}_S$ for $\alpha = 8/1024$ and $N = 256^3$ at $t = 0$ (orange), $t = 1$ (cyan), $t = 2$ (brown), $t = 3$ (blue), $t = 4$ (green), $t = 5$ (purple).	33
4.2	$e(k, t)$ vs. k starting from $\boldsymbol{\eta}_{TGV}$ for $\alpha = 64/1024$ and $N = 128^3$ at $t = 0$ (orange), $t = 1$ (cyan), $t = 2$ (brown), $t = 3$ (blue), $t = 4$ (green), $t = 5$ (purple).	34
4.3	Schematic illustration of the log-log dependence of $\kappa(\epsilon)$ on ϵ , cf. (4.60).	36
4.4	Dependence of $\kappa(\epsilon)$ on ϵ for $\alpha = 8/1024$	36
4.5	Dependence of $\kappa(\epsilon)$ on ϵ for $\alpha = 64/1024$	37
4.6	$\Phi_{\alpha, T}(\boldsymbol{\eta}_{\alpha, T}^{(n)})$ vs iteration number n for $\alpha = 16/1024$, $T = 5$, and different indicated values of σ	38

5.1	Isosurfaces of vorticity $\omega = \nabla \times \mathbf{u}_\alpha(T = 3; \boldsymbol{\eta}_{TGV})$. (a) magnitude $ \boldsymbol{\omega}(\mathbf{x}) $, and components (b) $ \omega_1 $, (c) $ \omega_2 $, and (d) $ \omega_3 $	43
5.2	Isosurfaces of vorticity $\omega = \nabla \times \mathbf{u}_\alpha(T = 5; \boldsymbol{\eta}_{TGV})$. (a) magnitude $ \boldsymbol{\omega}(\mathbf{x}) $, and components (b) $ \omega_1 $, (c) $ \omega_2 $, and (d) $ \omega_3 $	44
5.3	Solution of the Euler-Voigt system for $T = 3$ starting from $\boldsymbol{\eta}_{TGV}$. Log-log plot of $\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α at $t = 0$ (blue) to $t = 3$ (red), Steepest slope $p = -0.225$ (black), slope of $p = -1$ (dotted black) . .	44
5.4	Solution of the Euler-Voigt system for $T = 3$ starting from $\boldsymbol{\eta}_{TGV}$. Plot of the slope of $(\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2} \text{ vs } \alpha)$ vs time for each α	45
5.5	Solution of the Euler-Voigt system for $T = 5$ starting from $\boldsymbol{\eta}_{TGV}$. Log-log plot of $\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α at $t = 0$ (blue) to $t = 5$ (red), Steepest slope $p = -0.417$ (black), slope of $p = -1$ (dotted black) . .	45
5.6	Solution of the Euler-Voigt system for $T = 5$ starting from $\boldsymbol{\eta}_{TGV}$. Plot of the slope of $(\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2} \text{ vs } \alpha)$ vs time at each α	46
5.7	TGV for $T = 25$. Log-log plot of $\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2}$ vs α at $t = 0$ (blue) to $t = 25$ (red), Steepest slope $p =$ (black), slope of $p = -1$ (dotted black)	47
5.8	TGV for $T = 25$. Plot of the slope of $(\max_{t \in [0, T]} \ \nabla \mathbf{u}_\alpha(t)\ _{L^2} \text{ vs } \alpha)$ vs time at each α	47

List of Tables

4.1	Results ϵ_Φ for $\alpha = 16/1024, \dots, 64/1024$	30
4.2	Results ϵ_Φ for $\alpha = 8/1024, 12/1024$	30
4.3	The values of ϵ_α for $\alpha = 16/1024, \dots, 64/1024$	31
4.4	The values of ϵ_α for $\alpha = 8/1024, 12/1024$	32

Notation, and Abbreviations

Notation

x	Plain-text to denote scalar x
$\boldsymbol{x}$	Bold-text to denote vector x
$\boldsymbol{x}^\top$	Transpose of vector x
I	Identity Operator
∇	Gradient Operator
Δ	Laplacian Operator
$:=$	Equal to by definition

Abbreviations

PDEs	Partial Differential Equations
3D	three-dimensional or three dimensions
TGV	Taylor Green Vortex

LHS	left-hand side
RHS	right-hand side

Declaration of Academic Achievement

Dr. Zhao developed most of the mathematics for the subspace V (2.26), along with Theorem 2.1.3 and its proof. Dr. Zhao also developed the mathematics for much of the Euler-Voigt optimization problem described in Chapter 3, including the objective functional (3.32), the linearization of the Euler-Voigt equations (3.37), the adjoint system (3.38), and the gradient of the objective functional (3.44). The Riemannian conjugate gradient method for the Euler-Voigt optimization problem in Section 3.4 was developed by Dr. Zhao and Dr. Protas. This includes the tangent space (3.45), the projection operator (3.48), the search direction (3.52), and step size (3.55). Dr. Zhao also provided the data files for two initial guesses described in Chapter 3

The retraction operator (3.49), and the vector transport operation (3.50) were originally developed by Dr. Zhao and Dr. Protas but I modified these formulations to suit the present problem. The formulation of the manifold $\mathcal{M}_C$ (2.28) was based on work by Dr. Zhao, but the calculation of the constraint parameter C was my work. This included the calculations of the $\dot{H}^1$ seminorm (2.30) and vorticity (2.31) of the general Taylor Green Vortex, which are included in Appendices E and F. The results in Appendices B, G, H, and I are my re-derivations of Dr. Zhao's original results. The derivation in Appendices C and D are my re-derivations of the well known Euler and

Euler-Voigt scaling properties.

For the numerical models, Dr. Zhao created the original code base which I modified to suit this project. The determination of the numerical model constants, the time and spatial resolutions, regularization parameter α , and Gevrey parameter σ were also my work. These values were validated through analysis of the stability and accuracy of the model, conservation of the α -energy, comparison of the effect of different values of σ , examination of the solutions' Fourier spectrum, and the kappa test, all of which were my work and are described in Chapter 4. I also validated the results of the numerical models through comparison with the relevant findings of [5] and [20]. I performed all of the computations to obtain the final numerical results. The findings and conclusions in Chapters 5 and 6 are my work. The Python code used to perform the data analysis and produce the graphs is also my own work.

1 Introduction

1.1 Singularity Formation in Models of 3D Fluid Flows

The three-dimensional (3D) Navier-Stokes equations are a set of partial differential equations (PDEs) that describe the motion of homogeneous, viscous, incompressible fluids. The Navier-Stokes equations on the domain $\Omega \subseteq \mathbb{R}^3$ are defined as

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = 0 \quad \text{in } \Omega \times (0, T], \quad (1.1a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times [0, T], \quad (1.1b)$$

$$\mathbf{u}(\mathbf{x}, 0) = \boldsymbol{\eta} \quad \text{in } \Omega \quad (1.1c)$$

where $\mathbf{u} = [u_1, u_2, u_3]^\top$ is the fluid velocity field, p is the scalar pressure, $\nu > 0$ is the coefficient of kinematic viscosity, and T is the length of the time interval. The field $\boldsymbol{\eta}$ is the initial condition assumed to be divergence-free. Finally, the gradient operator $\nabla = [\partial_1, \partial_2, \partial_3]^\top$ is used to define the gradient of the pressure, the tensor $[\nabla \mathbf{u}]_{ij} = \partial_j u_i$ for $i, j = 1, 2, 3$, the divergence of the velocity field, and the Laplacian operator $\Delta \mathbf{u} = \nabla \cdot (\nabla \mathbf{u})$. Because of their generality, the Navier-Stokes equations are utilized to model the behaviour of systems in fields as disparate as aero/hydrodynamics, weather prediction, biophysics, and astrophysics [10, 12]. Despite the numerous fields

that make use of the 3D Navier-Stokes equations, and extensive research examining their properties, it is only known that weak Leray-Hopf solutions exist globally in time [22]. However, these solutions may contain singularities and their uniqueness remains unproven in most cases. It is also yet to be proven whether or not smooth classical solutions, defined in the pointwise sense, exist globally in time [28]. In fact, this is the basis of one of the seven Millennium Problems posed by the Clay Mathematics Institute which offers a substantial monetary prize for the confirmation of the existence, or non-existence, of a classical solution corresponding to smooth initial conditions [14]. The space of smooth functions and all other function spaces utilized in this analysis are defined in Appendix A. It is common for scientific research that focuses the fundamental properties of the Navier-Stokes equations to utilize either the unbounded domain $\Omega := \mathbb{R}^3$, the cylindrical domain $\Omega := \{(r, z) : r < 1, z \in \mathbb{T}\}$, or the periodic 3D torus $\Omega := \mathbb{T}_L^3$ with domain size $L > 0$ [28]. For this analysis, we use the 3D torus of unit size ($L = 1$) with periodic boundary conditions.

A central question in fluid mechanics is whether or not the Navier-Stokes equations are globally well-posed or if singularities can form from smooth initial conditions. A singularity occurs when the derivative of the velocity field becomes infinite at a single point [25]. The formation of a singularity in a fluid flow, from smooth initial conditions, would invalidate the Navier-Stokes equations as an accurate description of fluid flows [28]. The question about singularity formation from smooth initial conditions also remains open for the Euler equations [16]. The Euler equations are the inviscid

form of the Navier-Stokes equations, obtained by setting $\nu = 0$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = 0 \quad \text{in } \Omega \times (0, T], \quad (1.2a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times (0, T], \quad (1.2b)$$

$$\mathbf{u}(\mathbf{x}, 0) = \boldsymbol{\eta} \quad \text{in } \Omega. \quad (1.2c)$$

The Euler equations are often studied because they are simpler to analyse than the Navier-Stokes equations due to the absence of the viscous dissipation. The local existence of classical solutions to the Euler equations was proven in [19] for solutions defined in Sobolev spaces.

Theorem 1.1.1 ([19]). *If $\boldsymbol{\eta} \in H^m(\mathbb{T}^3)$ for some $m > 5/2$ and satisfies $\nabla \cdot \boldsymbol{\eta} = 0$, then there exists a time $T = T(\|\boldsymbol{\eta}\|_{H^m}) > 0$ such that (1.2) has a unique solution $\mathbf{u}(\cdot; \boldsymbol{\eta}) \in C([0, T]; H^m) \cap C^1([0, T]; H^{m-1})$,*

where $\|\cdot\|_{H^m}$ is the standard norm in the Sobolev space H^m defined in Appendix A. If a singularity is to form in the solutions to the Euler equations, it must occur after the minimum local existence time defined in Theorem 1.1.1. There are many different conditional regularity results for determining if a singularity has formed, the best known of which is the Beale-Kato-Majda (BKM) criterion [3] which asserts

Theorem 1.1.2 ([3]). *A smooth solution $\mathbf{u}$ of the Euler system develops a singularity at $t = t_0$ if and only if*

$$\lim_{t \rightarrow t_0} \int_0^t \|\boldsymbol{\omega}(\tau)\|_{L^\infty} d\tau = \infty, \quad \boldsymbol{\omega} = \nabla \times \mathbf{u}, \quad (1.3)$$

where $\|\cdot\|_{L^\infty}$ is the standard norm in the Lebesgue space L^∞ defined in Appendix A.

A key difference between the Navier-Stokes and Euler equations is that there is more numerical evidence that indicates singularity formation in Euler is possible from smooth initial conditions. For example, [4, 5] presented numerical evidence for singularity formation in the Euler equations on a 3D torus ($L = 2\pi$). Hou et al. produced several papers which provide numerical evidence for singularity formation in the 3D axisymmetric Euler equations [24, 18, 17] and which proved singularity formation in the 3D axisymmetric Euler equations with rigid boundaries and periodic boundary conditions in the axial direction [8, 9]. We make special note of the optimization-based approach of [30], which utilized a Riemannian conjugate gradient method to identify ‘extreme’ initial conditions, and reports numerical results consistent with singularity formation in Euler flows on the 3D unit torus. **Two of the optimal initial conditions identified in [30] are used as initial guesses in this analysis and are discussed in more detail in Chapter 3.** Finally, [20] which provides numerical evidence for singularity formation in the Euler equations based on the Voigt regularization. These final two papers are of special significance for us because they form the primary basis for the motivation and methodology of this analysis.

1.2 3D Euler-Voigt Equations

The Euler-Voigt equations are an inviscid regularization of the Euler equations (1.2) which have been shown to be globally well-posed and so do not form singularities

from smooth initial conditions [7]. The Euler-Voigt equations are defined as

$$-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p_\alpha = 0 \quad \text{in } \Omega \times (0, T], \quad (1.4a)$$

$$\nabla \cdot \mathbf{u}_\alpha = 0 \quad \text{in } \Omega \times (0, T], \quad (1.4b)$$

$$\mathbf{u}_\alpha(\mathbf{x}, 0) = \boldsymbol{\eta}_\alpha \quad \text{in } \Omega, \quad (1.4c)$$

where the regularization is accomplished with the addition of the term $-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha$ in which $\alpha > 0$ is the regularization parameter with units of length. This term stabilizes the system without dissipation, which would remove energy from the system [20]. The regularization term acts as a spectral filter and suppresses the development of small scale structures with length-scale smaller than α [26]. We introduce the subscript α to distinguish between solutions of (1.2) and (1.4) i.e., $\mathbf{u}$ and $\mathbf{u}_\alpha$.

For (1.2) the total kinetic energy is conserved so long as the solution remains well-posed in the classical sense

$$E_k(t) := \frac{1}{2} \|\mathbf{u}(t)\|_{L^2}^2 = E_k(0), \quad (1.5)$$

where the norm $\|\cdot\|_{L^2}$ here is the standard Lebesgue L^2 norm. For (1.4) the conserved quantity is known to be the α -energy. The α -energy is defined as

$$E_\alpha(t) := \|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = E_\alpha(0), \quad (1.6)$$

where $\|\mathbf{u}_\alpha(t)\|_{L^2}^2$ is the energy term, and $\|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2$ is the enstrophy term, scaled by α^2 .

1.3 Outline of the Thesis

The goal of this analysis is to search for possible singularities in Euler flows corresponding to smooth initial conditions. This search will be performed based on a PDE-constrained optimization problem in which we will identify initial condition that produce the most 'extreme' behaviour in Euler-Voigt flows. In Chapter 2 we present the properties of the Euler-Voigt equations, including the blow-up criteria, scaling property, and manifold over which we search for initial conditions. In Chapter 3 we present the formulation of the optimization problem. This includes the description of our objective functional, the gradient calculation, and the Riemannian conjugate gradient method used to compute our search direction and iterative relation. In Chapter 4 we describe the numerical methods used to perform the calculations, including approximation in space and time, Gevrey parameters, and the regularization parameter. In addition, we validate the computations using the Fourier spectrum, kappa test, and conservation of α -energy. In Chapter 5 we present our results. We begin with the solutions of the Euler-Voigt equations with the initial conditions given by the initial guesses, without optimization, as $\alpha \rightarrow 0$. These results indicate that singularities do form in Euler flows corresponding to these initial conditions but only when we significantly extend the time window beyond the expected blow-up time. We then present our optimization results as $\alpha \rightarrow 0$, which indicate that RESULTS RESULTS RESULTS. Finally, in Chapter 6 we summarize our findings and present our conclusion.

2 Properties of the Euler-Voigt Equations

In this chapter we prove a theorem which provides the criterion used to determine if a singularity forms in Euler flows from smooth initial conditions. Additionally, we construct a manifold over which we search for optimal smooth initial conditions, which will form the basis for our optimization problem in Chapter 3.

2.1 Finite Time Blow-up Criterion

In order to begin our search for finite-time singularity formation, we must first describe the criteria used to detect singularity formation in Euler flows. In [21] a new criterion for identifying singularity formation in Euler flows was developed which utilizes the Euler-Voigt equations. This criterion relies on the convergence of the solutions of (1.4) to the solutions of (1.2) as $\alpha \rightarrow 0$.

Theorem 2.1.1 ([21]). *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$, let $\boldsymbol{u}, \boldsymbol{u}_\alpha \in C([0, T], H^s(\Omega)) \cap C^1([0, T], H^{s-1}(\Omega))$ be the corresponding solutions to (1.2) and (1.4), respectively, where $0 < T < T_{max}$ and T_{max} is the maximal time for which*

a solution to (1.2) exists and is unique. For all $t \in [0, T]$, we have

$$\begin{aligned} & \|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \leq \\ & (\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2) e^{Ct} + C\alpha^2(e^{Ct} - 1). \end{aligned} \quad (2.7)$$

Based on this theorem we know that if

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (2.8)$$

then for all $t \in [0, T]$,

$$\|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \rightarrow 0. \quad (2.9)$$

Therefore, solutions of the Euler-Voigt system (1.4) converge to the corresponding Euler (1.2) flows in H^1 uniformly in time as $\alpha \rightarrow 0$. Using this result, the following theorem was proved in [20].

Theorem 2.1.2. *Let $\boldsymbol{\eta}_\alpha \in H^s$, $s \geq 3$, with $\nabla \cdot \boldsymbol{\eta}_\alpha = 0$ and let $\mathbf{u}_\alpha$ be the corresponding unique solution of (1.4). Suppose that there exists a time $T^* > 0$ such that*

$$\limsup_{\alpha \rightarrow 0} \left(\alpha \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \right) > 0. \quad (2.10)$$

Then, the corresponding solution of the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}_\alpha$ must develop a singularity within the interval $[0, T^]$.*

We note that this theorem relies on the solutions of (1.2) and (1.4) starting from the same initial condition, i.e., $\boldsymbol{\eta} = \boldsymbol{\eta}_\alpha$. This necessarily satisfies (2.8) as $\alpha \rightarrow 0$. However, in order to use this result in our study, we need to generalize Theorem 2.1.2

for the case where (2.8) is satisfied but $\boldsymbol{\eta} \neq \boldsymbol{\eta}_\alpha$. Therefore, we modify Theorem 5.2 from [21] as follows.

Theorem 2.1.3 (X. Zhao, private communication). *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$ satisfying*

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (2.11)$$

let $\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\mathbb{T}^3)) \cap C^1([0, T], H^{s-1}(\mathbb{T}^3))$ be the corresponding solutions to (1.2) and (1.4), respectively. If there exists a finite time $T^ > 0$ such that the solutions $\mathbf{u}_\alpha$ satisfy*

$$\limsup_{\alpha \rightarrow 0} \left(\alpha^2 \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) > 0, \quad (2.12)$$

then $\mathbf{u}$ develops a singularity in the interval $[0, T^]$.*

Proof. Suppose that the solution $\mathbf{u}$ of the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}$ is regular on $[0, T^*]$. Since the Euler-Voigt equations are globally well-posed, we have, cf. (1.6)

$$\|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2, \quad (2.13)$$

thus

$$\alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2 - \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (2.14)$$

Taking the supremum over $t \in [0, T^*]$ of the above equation, we obtain that

$$\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta}_\alpha)\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (2.15)$$

Taking the limsup as $\alpha \rightarrow 0$, we obtain

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \limsup_{\alpha \rightarrow 0} \left(\|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2 \right). \quad (2.16)$$

We now make use of Theorem 2.1.1, which demonstrates that the convergence of the solutions of the Euler-Voigt equations (1.4) to the solution of the Euler equations (1.2) is uniform on $[0, T^*]$. Therefore we can reduce the above equation to

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \|\boldsymbol{\eta}\|_{L^2}^2 - \|\mathbf{u}(t)\|_{L^2}^2. \quad (2.17)$$

Since the energy of regular solutions of the Euler equations is conserved, we have

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = 0, \quad (2.18)$$

which contradicts the hypothesis in the theorem that the above expression is positive.

Equality (2.17) holds because

$$\begin{aligned} \limsup_{\alpha \rightarrow 0} \left| \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2} - \|\boldsymbol{\eta}\|_{L^2} \right| &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \mathbf{u}(t_\alpha)\|_{L^2}^2 + \|\mathbf{u}(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &= 0, \end{aligned}$$

where the reverse triangle and triangle inequalities were used and t_α is the time at which

$$\inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2} = \|\mathbf{u}_\alpha(t_\alpha)\|_{L^2}. \quad (2.20)$$

□

2.2 Initial Conditions

We have established Theorem 2.1.3 which we will use to identify singularity formation in Euler flows corresponding to smooth initial conditions. In order to formulate our optimization problem we must now describe the function space we perform our search in. Theorems 2.1.1, 2.1.2, and 2.1.3 presented in Section 2.1 require initial conditions $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$. However, the pseudo-spectral methods used in this study to numerically solve system (1.4) require the functions to be analytic in order to have exponential convergence [6]. For the Navier-Stokes equations (1.1), the viscous term acts to instantaneously regularize the solution in space. Therefore, if the initial condition has only Sobolev regularity, the resulting flow becomes real-analytic (with Gevrey regularity defined below) at $t = 0^+$. This regularity allows the solution to achieve exponential convergence.

The Euler equations (1.2) do not include any viscous effects and so the solution simply stays at the same level of smoothness as the initial condition, unless blow-up occurs. For Euler-Voigt the regularization term keeps the solution globally well-posed but does not improve the regularity [21]. Therefore, we must have analytic initial conditions to make use of the pseudo-spectral methods. For all the models considered here, a requirement is that the flow is divergence free at all times, including the initial conditions. Finally, on a periodic domain the quantity $\int_\Omega \boldsymbol{\eta}_\alpha d\mathbf{x}$ is an invariant of motion for solutions of the Euler-Voigt equations. Due to this, we assume without loss of generality that the initial conditions have a mean of zero. In order to identify initial conditions that satisfy these criteria, we first consider analytic initial conditions that

belong to the Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\mathbb{T}^3) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0, \quad (2.21)$$

with the norm defined as

$$\forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} := \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\widehat{\mathbf{u}}_{\mathbf{j}}} \quad (2.22a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}, \quad (2.22b)$$

where hat denotes the Fourier coefficient

$$\widehat{\mathbf{u}}_{\mathbf{k}} = \int_{\Omega} e^{-i\mathbf{k} \cdot \mathbf{x}} \mathbf{u}(\mathbf{x}) \, d\mathbf{x}, \quad (2.23)$$

overbar denotes complex conjugation, and the operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D|\mathbf{v}} \right]_{\mathbf{j}} := 2\pi|\mathbf{j}| \widehat{\mathbf{v}}_{\mathbf{j}}, \quad \left[\widehat{e^{\sigma|D|}\mathbf{v}} \right]_{\mathbf{j}} := e^{2\pi|\mathbf{j}|} \widehat{\mathbf{v}}_{\mathbf{j}}. \quad (2.24)$$

The parameter σ determines the relative 'size' of the Gevrey space

$$G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s, \quad 0 < \sigma_1 < \sigma_2, \quad (2.25)$$

which is proved in Appendix B. The value of σ is chosen in Section 4.4. In order to satisfy the divergence free condition, we introduce the following subspace in which

optimal initial conditions will be sought

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \int_{\mathbb{T}^3} \mathbf{v} \, d\mathbf{x} = 0\}. \quad (2.26)$$

2.3 Scaling Symmetries of Euler and Euler-Voigt Systems

Both (1.2) and (1.4) possess a scaling property. Given a solution $\mathbf{u}$, or $\mathbf{u}_\alpha$, we can construct a new solution

$$\mathbf{v}(\mathbf{y}, \tau) = \frac{1}{\gamma} \mathbf{u}(\beta \mathbf{x}, \lambda t), \quad \mathbf{y} = \beta \mathbf{x}, \quad \tau = \lambda t \quad (2.27)$$

where $\lambda, \gamma, \beta > 0$ must satisfy $\beta/(\lambda\gamma) = 1$, which is proven in Appendices C and D, respectively for the Euler (1.2) and Euler-Voigt (1.4) equations. Our goal is to find a time interval $[0, T]$ that is sufficiently large so that it is outside the interval of local existence established by Kato's theorem 1.1.1 and so allows for the potential of singularity formation in Euler flows. Due to this scaling property of the Euler-Voigt equations (2.27), the length of the time interval will depend on the 'size' of our solution (γ) and the spatial scale (β). Our domain size will not change, and so in order to keep a consistent time scale we must fix γ between all initial conditions. To accomplish this, we restrict our discussion to initial conditions with a fixed $\dot{H}^1$ seminorm which belong to a closed manifold $\mathcal{M}_C \subset V$ defined as cf. (2.26)

$$\mathcal{M}_C = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = C\}, \quad (2.28)$$

where $C > 0$ is a constant. We construct the manifold by fixing the $\dot{H}^1$ seminorm because of the form of our objective functional used in the optimization search, which

will be described in Chapter 3. The question we must answer now is what value of the constraint parameter C we use. To determine this we look to our first initial condition, the 3D Taylor Green Vortex (TGV). The TGV in the domain $\Omega := \mathbb{T}_L^3$ has the form

$$\boldsymbol{\eta}_{TGV}(\mathbf{x}) = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0 \end{bmatrix}, \quad (2.29)$$

for some $V_0 \neq 0$. The $\dot{H}^1$ seminorm of (2.29) is

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3L}\pi V_0, \quad (2.30)$$

which is proven in Appendix E. The BKM criterion (1.1.2) uses the quantity $\|\boldsymbol{\omega}(t)\|_{L^\infty}$ to determine singularity formation. For the TGV

$$\|\boldsymbol{\omega}\|_{L^\infty} = \|\nabla \times \boldsymbol{\eta}_{TGV}\|_{L^\infty} = 4\pi V_0/L, \quad (2.31)$$

which is proven in Appendix F. In [5], the authors provided numerical evidence that blow-up of (1.2) may occur around $T^* \approx 4$ for initial condition (2.29) with parameters $L = 2\pi$ and $V_0 = 1$. This result provides the baseline for our study and so we must match this time window so that singularity formation identified by (2.1.3), if it happens, occurs at approximately the same time as in [5]. Therefore, we set $\lambda = 1$. Our Euler-Voigt system is defined on the domain Ω with $L = 1$ and so we set $\beta = 1/2\pi$. Therefore, from (2.27) we set $\gamma = 1/2\pi$. The model we are aiming to match uses $V_0 = 1$ and so to match the time window we use $V_0 = 1/2\pi$.

These considerations give $\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$ and $\|\nabla \times \boldsymbol{\eta}_{TGV}\|_{L^\infty} = 2$ for the initial condition (2.29). Therefore, we can restrict our discussion to initial conditions with $\dot{H}^1$ seminorm which belong to a closed manifold (2.28) with $C = \sqrt{3}/2$. In order to select initial conditions within this manifold, we will first select initial conditions $\boldsymbol{\eta}_\alpha \in V$ and then apply a retraction function to ensure it has $\|\boldsymbol{\eta}_\alpha\|_{\dot{H}^1} = \sqrt{3}/2$. To verify that this selection of the $\dot{H}^1$ seminorm of the initial condition is correct, we compare this scaling to the results shown in Figure 1b of [5]. This figure shows the evolution of the L^∞ norm of the vorticity over time, starting from $\boldsymbol{\eta}_{TGV}$.




Figure 2.1: Maximum of vorticity $\|\boldsymbol{\omega}(t)\|_{L^\infty}$. Results from runs performed at different resolutions are displayed together: 128^3 (orange plus), 256^3 (cyan star), 512^3 (brown triangles), 1024^3 (blue squares).

We present our reproduction in Figure 2.1 which shows that the evolution of the vorticity at each resolution in our scaled model matches exactly to the corresponding computations shown in Figure 1b in [5]. Therefore, our construction of the manifold $\mathcal{M}_C$ is correct.

3 The Optimization Problem

In this chapter we formulate the main optimization problem and introduce the Riemannian conjugate gradient method used to solve it. Additionally, we examine the structure of the initial conditions including the Taylor Green Vortex, and two optimized initial conditions from the results of [30].

3.1 Formulation of the Problem

With our initial condition and blowup criterion defined by Theorem 2.1.3, we now construct the PDE-constrained optimization problem. We begin with the objective functional $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$, which is defined as

$$\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) := \|\nabla \mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{L^2}^2, \quad (3.32a)$$

$$= \|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{H^1}^2, \quad (3.32b)$$

where $\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)$ is the solution of the Euler-Voigt equations (1.4) obtained at the end of the time interval of size T with the initial condition $\boldsymbol{\eta}_\alpha \in V$ and regularization parameter α . Using this objective functional, we define the optimization problem.

Problem 3.1.1. *Given $\alpha, T \in \mathbb{R}^+$, find*

$$\tilde{\boldsymbol{\eta}}_{\alpha,T} = \operatorname{argmax}_{\boldsymbol{\eta}_{\alpha}^0 \in M_C} \Phi_{\alpha,T}(\boldsymbol{\eta}_{\alpha}^0), \quad (3.33)$$

where $\boldsymbol{\eta}_{\alpha}^0$ is our initial guess. Our goal is to first fix T and solve the optimization problem for a sequence of decreasing values of α that converge to 0. From Figure 1b of [5], the expected time of singularity formation in the Euler flow, with the TGV scaled as discussed in Section 2.3 used as the initial condition, is $T^* \approx 4$. From Theorem 2.1.3, if the initial condition criterion (2.8) and objective functional criteria (2.12) are both satisfied, then the corresponding Euler flow develops a singularity in the interval $[0, T^*]$ from initial condition $\boldsymbol{\eta}$. We examine the results for a time interval slightly shorter and longer than T^* . Therefore, we consider the optimization problem for $T = 3$, and $T = 5$. With optimization, blow-up may occur in both cases. All regularization parameters used in this analysis are a fraction $\alpha = \cdot/1024$. In order to simplify the notation of our optimal solution, we drop the denominator, i.e., the optimal initial condition found by solving problem (3.1.1) for regularization parameter $\alpha = 8/1024$ and time interval $T = 5$ would be denoted $\tilde{\boldsymbol{\eta}}_{8,5}$.

3.2 Initial Guesses

Problem 3.1.1 is non-convex and so may admit a different local maxima depending on the initial guess used at iteration zero, $\boldsymbol{\eta}_{\alpha}^0$. In order to search for optimal initial conditions that may lead to a singularity in the Euler flow, we must solve the optimization problem from different initial guesses. Before we begin our search for singularities, we examine the structure of the initial guesses used in this study. These

include the TGV as well as two optimal initial conditions from the results of [30].

Figure 3.1 shows the isosurfaces of the magnitude of the vorticity $\boldsymbol{\omega} = \nabla \times \boldsymbol{\eta}_{TGV}$. In [30], the authors utilized the Riemannian conjugate gradient method to identify initial conditions that produce 'extreme' behaviour in Euler flows. Their analysis used the same domain Ω as our analysis. However, the manifold in which they searched for optimal initial conditions was formulated by scaling the initial conditions to a fixed $\|\boldsymbol{\eta}\|_{\dot{H}^3} = 1$ and the objective functional was $\Phi_T(\boldsymbol{\eta}) := \|\mathbf{u}(T; \boldsymbol{\eta})\|_{\dot{H}^3}^2$. With the same domain size $L = 1$, this gives their initial conditions a time interval which is scaled by a factor of $6\sqrt{3}\pi^2$ to our model, which is shown in Appendix E. They solved the optimization problem with multiple initial guesses, including the TGV, and time intervals of $T = 25$ and $T = 75$. In our manifold (2.28), this is equivalent to starting from the TGV and optimizing with time intervals of $T \approx 0.2437$ and $T \approx 0.7312$, respectively. The first initial condition from [30] is the optimal initial condition found using the short time window $T = 25$ and a resolution of $N = 128^3$. We denote this initial condition as $\boldsymbol{\eta}_S$. Figure 3.2 shows the isosurfaces of the magnitude of the vorticity for $\boldsymbol{\eta}_S$. The second is the optimal initial condition found using the long time window $T = 75$ and a resolution of $N = 512^3$, which we denote as $\boldsymbol{\eta}_L$. Figure ?? shows the isosurfaces of the magnitude of the vorticity for $\boldsymbol{\eta}_L$.




Figure 3.1: Isosurfaces of vorticity $\omega = \nabla \times \boldsymbol{\eta}_{TGV}$ (a) magnitude $|\omega(\mathbf{x})|$, and components (b) $|\omega_1|$, (c) $|\omega_2|$, and (d) $|\omega_3|$.

3.3 Gradient of Objective Function

The optimization problem (3.1.1) is solved using a Riemannian conjugate gradient method which uses an optimal search direction and distance to create the next approximation of the optimal initial condition. This method is utilized because it accelerates the gradient ascent method which is necessary given the computational cost of each iteration. To solve this problem, we utilize a 'optimize-then-discretize' approach where the optimization problem, and the Riemannian conjugate gradient method, are derived in an infinite-dimensional function space and then discretized to solve numerically. The same method was used in [30] to solve a PDE-constrained optimization problem for the Euler equations to search for initial conditions $\boldsymbol{\eta}$ that produce 'extreme' flows.




Figure 3.2: Isosurfaces of vorticity $\omega = \nabla \times \boldsymbol{\eta}_S$ (a) magnitude $|\omega(\mathbf{x})|$, and components (b) $|\omega_1|$, (c) $|\omega_2|$, and (d) $|\omega_3|$.

We compute the optimal search direction using the gradient of the objection functional $\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)$. It is obtained using an adjoint method which depends on solving a properly defined adjoint system backwards in time. To derive an expression for the gradient, we first introduce the Gâteaux (directional) differential $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) : V \times V \rightarrow \mathbb{R}$. The Gateaux differential is defined as,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)]. \quad (3.34)$$

This represents the change in $\Phi_{\alpha,T}$ from an infinitesimal perturbation in the direction $\boldsymbol{\eta}'_\alpha$ from the initial condition $\boldsymbol{\eta}_\alpha$. By fixing the initial condition $\boldsymbol{\eta}_\alpha$, we can view the Gâteaux differential as a bounded linear functional on V . Therefore, it can also be

represented as an inner product on the Gevrey space using the Riesz representation theorem [30, 23]

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.35)$$

Substituting the second definition of the objective functional (3.32b) into the definition of the Gâteaux differential (3.34), we derive the Gâteaux differential in the form of the $\dot{H}^1$ inner product,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(T), \mathbf{u}'_\alpha(T) \rangle_{\dot{H}^1} \quad (3.36)$$

The derivation for this is shown in Appendix I. In (3.36) the value $\mathbf{u}'_\alpha$ is the solution of the linearization of the Euler-Voigt system (1.4) around its solution $\mathbf{u}_\alpha$ corresponding to the initial condition $\boldsymbol{\eta}_\alpha$. The linearization of the Euler-Voigt system is defined as

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p'_\alpha \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p'_\alpha) \\ \nabla \cdot \mathbf{u}'_\alpha \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (3.37a)$$

$$\mathbf{u}'_\alpha(x, 0) = \boldsymbol{\eta}'_\alpha \quad (3.37b)$$

where $\boldsymbol{\eta}'_\alpha$ and p'_α are the perturbations of the initial condition and pressure, respectively. The derivation of the linearized form of the Euler-Voigt equations is shown in Appendix G. In order to obtain the gradient of the objective functional from the Gâteaux differential we must first introduce the fields $\{\mathbf{u}_\alpha^*, p_\alpha^*\}$ which are the solution

of the adjoint system

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}_\alpha^* - (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}_\alpha^* + (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}_\alpha^*)^\top) \mathbf{u}_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot \mathbf{u}_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (3.38a)$$

$$\mathbf{u}_\alpha^*(\mathbf{x}, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T) \quad (3.38b)$$

which is derived in Appendix H by performing integration by parts, in both space and time, on the following duality-pairing relation

$$\begin{aligned} \left(\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p'_\alpha \end{bmatrix}, \begin{bmatrix} \mathbf{u}_\alpha^* \\ p'_\alpha \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p'_\alpha \end{bmatrix}, \begin{bmatrix} \mathbf{u}_\alpha^* \\ p'_\alpha \end{bmatrix} d\mathbf{x} dt \\ &= \left(\begin{bmatrix} \mathbf{u}'_\alpha \\ p'_\alpha \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p'_\alpha \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, T) \cdot \mathbf{u}_\alpha^*(\mathbf{x}, T) d\mathbf{x} \\ &\quad - \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, 0) \cdot \mathbf{u}_\alpha^*(\mathbf{x}, 0) d\mathbf{x} \\ &= 0. \end{aligned} \quad (3.39)$$

Using the solutions of the adjoint system we can now obtain a convenient expression for the gradient. With the Gâteaux differential (3.36) expressed in terms of the L^2 inner product

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(0), \mathbf{u}'_\alpha(0) \rangle_{L^2}, \quad (3.40)$$

which is derived in Appendix I, and in the Riesz representation form (3.35), we obtain

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(0), \mathbf{u}'_\alpha(0) \rangle_{L^2} = \langle \nabla \Phi_{\alpha, T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.41)$$

Using the definition of the Gevrey inner product (2.22), the L^2 inner product can be converted to the Gevrey inner product,

$$\langle \mathbf{u}_\alpha^*(0), \mathbf{u}'_\alpha(0) \rangle_{L^2} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.42)$$

Therefore, we have two forms of the Gâteaux differential as a Gevrey inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (3.43)$$

From this we obtain the gradient of the objective functional

$$\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0). \quad (3.44)$$

The adjoint system (3.38) is linear and represents a terminal value problem. Its coefficients are determined by evolving the Euler-Voigt system forward from the initial condition $\boldsymbol{\eta}_\alpha$ to compute $\mathbf{u}_\alpha(T)$, from which we obtain the terminal value of (3.38). Finally, we evolve the adjoint system backwards in time to obtain the initial value $\mathbf{u}_\alpha^*(0)$ which is used to compute the gradient (3.44).

3.4 Riemannian Conjugate Gradient Method

When solving the optimization problem, we look for an initial condition $\boldsymbol{\eta}_\alpha \in M_C$ that maximizes our objective functional. Therefore, not only must our initial guess $\boldsymbol{\eta}_\alpha^0$ have a $\dot{H}^1$ seminorm of $\sqrt{3}/2$, but so too must each following iterate $\boldsymbol{\eta}_\alpha^{(n)}$. We ensure this constraint is satisfied using a Riemannian conjugate gradient method modified from the method used in [11, 28, 30]. This method has three steps:

1. Project the gradient $\nabla\Phi_T(\boldsymbol{\eta}_\alpha)$ onto the tangent space to $\mathcal{M}_C$ at $\boldsymbol{\eta}_\alpha$.
2. Perform a suitable vector transport operation in order to construct a Riemannian conjugate ascent direction using the previous search direction.
3. Retract the resulting state back to the constraint manifold $\mathcal{M}_c$.

We first define the space tangent to $\mathcal{M}_C$ at $\mathbf{v}$ as

$$T_{\mathbf{v}}\mathcal{M}_C = \{\mathbf{z} \in V : \langle \mathbf{v}, \mathbf{z} \rangle_{\dot{H}^1} = 0\}. \quad (3.45)$$

For any $\mathbf{z} \in \mathcal{T}_{\mathbf{v}}\mathcal{M}_C$, this condition can be expressed as

$$\langle \mathbf{v}, \mathbf{z} \rangle_{\dot{H}^1} = \int_{\mathbb{T}^3} |D| \mathbf{v} \cdot |D| \mathbf{z} d\mathbf{x} \quad (3.46a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v} \right) \cdot \mathbf{z} d\mathbf{x} \quad (3.46b)$$

$$= \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}, \mathbf{z} \rangle_{G^\sigma} = 0. \quad (3.46c)$$

Thus, at each point $\mathbf{v} \in M_C$, the tangent space $\mathcal{T}_{\mathbf{v}}\mathcal{M}_C$ can be characterized using the unit 'normal' vector given by

$$\mathbf{n}_{\mathbf{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}\|_{G^\sigma}} \quad (3.47)$$

which allows us to construct the projection operator $\mathcal{P}_{\mathcal{T}_{\mathbf{v}}\mathcal{M}_C} : V \rightarrow \mathcal{T}_{\mathbf{v}}\mathcal{M}_C$,

$$\forall \boldsymbol{\omega} \in V, \quad \mathcal{P}_{\mathcal{T}_{\mathbf{v}}\mathcal{M}_C} \boldsymbol{\omega} = \boldsymbol{\omega} - \langle \mathbf{n}_{\mathbf{v}}, \boldsymbol{\omega} \rangle_{G^\sigma} \mathbf{n}_{\mathbf{v}}. \quad (3.48)$$

As stated previously, the initial conditions at each iteration must have a $\dot{H}^1$ seminorm of $\sqrt{3}/2$. In order to ensure this we introduce the retraction operator

$$R(\mathbf{v}) = \frac{\sqrt{3}}{2} \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq \mathbf{0}. \quad (3.49)$$

The conjugate-gradient method uses the search direction obtained at the n -th iteration to compute the search direction at the $(n+1)$ -th iteration. This is done by computing a linear combination of the gradient at the current iteration and the search direction from the previous iteration, weighted with a 'momentum' coefficient. However, the two directions belong to different tangent spaces to $\mathcal{M}_C$ and thus cannot be directly added. In order to allow algebraic operations to be performed on the vectors of the two tangent spaces we introduce the vector transport operation defined by [1]

$$\mathcal{TM}_C \oplus \mathcal{TM}_C \rightarrow \mathcal{TM}_C \quad : \quad (\boldsymbol{\varphi}, \boldsymbol{\xi}) \mapsto \Gamma_{\boldsymbol{\varphi}}(\boldsymbol{\xi}) \in \mathcal{TM}_C,$$

where $\mathcal{TM}_C = \bigcup_{\mathbf{v} \in \mathcal{M}_C} \mathcal{T}_{\mathbf{v}}\mathcal{M}_C$ is the tangent bundle on $\mathcal{M}_C$. The operation transports the vector field $\boldsymbol{\xi}$ along the manifold $\mathcal{M}_C$ in the direction of $\boldsymbol{\varphi}$. This allows us to map search directions between the tangent spaces of two consecutive iterations. This operator is defined via differentiated retraction

$$\Gamma_{\boldsymbol{\varphi}_{\mathbf{u}}}(\boldsymbol{\xi}_{\mathbf{u}}) := \frac{d}{dt} R(\boldsymbol{\varphi}_{\mathbf{u}} + t\boldsymbol{\xi}_{\mathbf{u}}) \Big|_{t=0} \quad (3.50a)$$

$$:= \frac{\sqrt{3}}{2} \frac{1}{\|\mathbf{u} + \boldsymbol{\varphi}_{\mathbf{u}}\|_{\dot{H}^1}} \left[\boldsymbol{\xi}_{\mathbf{u}} - \frac{\langle \mathbf{u} + \boldsymbol{\varphi}_{\mathbf{u}}, \boldsymbol{\xi}_{\mathbf{u}} \rangle_{\dot{H}^1}}{\|\mathbf{u} + \boldsymbol{\varphi}_{\mathbf{u}}\|_{\dot{H}^1}^2} (\mathbf{u} + \boldsymbol{\varphi}_{\mathbf{u}}) \right], \quad \mathbf{u} \in \mathcal{M}_C. \quad (3.50b)$$

With our operators defined we can now construct the iterative relation that will be used to maximize our objective functional (3.32)

$$\boldsymbol{\eta}_\alpha^{(n+1)} = R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau_n \mathbf{d}^{(n)} \right]. \quad (3.51)$$

To simplify notation, let $\mathbf{G}^{(n)} := \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha^{(n)})$ and let $\mathcal{T}_n := \mathcal{T}_{\boldsymbol{\eta}_\alpha^{(n)}} \mathcal{M}_C$. The search direction at iteration n ($\mathbf{d}^{(n)}$) is computed as

$$\begin{aligned} \mathbf{d}^{(0)} &= P_{\mathcal{T}_0} \mathbf{G}^{(0)} \\ \mathbf{d}^{(n)} &= P_{\mathcal{T}_n} \mathbf{G}^{(n)} + \beta_n \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)}(\mathbf{d}^{(n-1)}), \quad n \geq 1, \end{aligned}$$

where the 'momentum' term is computed using the Polak-Ribière approach [27].

$$\beta_n = \frac{\left\langle P_{\mathcal{T}_n} \mathbf{G}^{(n)}, \left(P_{\mathcal{T}_n} \mathbf{G}^{(n)} - \Gamma_{\tau_{n-1}} \mathbf{d}^{(n-1)} P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)} \right) \right\rangle_{G^\sigma}}{\|P_{\mathcal{T}_{n-1}} \mathbf{G}^{(n-1)}\|_{G^\sigma}^2}. \quad (3.53)$$

In order to remove old data that may slow down convergence, the algorithm is periodically restarted by setting $\beta_n = 0$. For our optimization search, we restart the algorithm every 20 iterations, and when the search direction is not an ascent direction, i.e, when

$$\frac{\left\langle \mathbf{d}^{(n)}, P_{\mathcal{T}_n} \mathbf{G}^{(n)} \right\rangle_{G^\sigma}}{\|\mathbf{d}^{(n)}\|_{G^\sigma} \|P_{\mathcal{T}_n} \mathbf{G}^{(n)}\|_{G^\sigma}} < 0. \quad (3.54)$$

Finally, our optimal step size is determined by the distance along an arc tangent to $\mathbf{d}^{(n)}$ which maximizes the objective functional, it is computed by performing an arc-search along this direction using a modification of Brent's method

$$\tau_n = \operatorname{argmax}_{\tau > 0} \Phi_T \left(R \left[\boldsymbol{\eta}_\alpha^{(n)} + \tau \mathbf{d}^{(n)} \right] \right). \quad (3.55)$$

The optimization search is terminated when the relative change in the objective functional from (3.51) is below a chosen tolerance.

$$0 \leq \frac{\Phi_{\alpha,T}(\boldsymbol{\eta}_{\alpha}^{(n+1)}) - \Phi_{\alpha,T}(\boldsymbol{\eta}_{\alpha}^{(n)})}{\Phi_{\alpha,T}(\boldsymbol{\eta}_{\alpha}^{(n)})} < tol, \quad (3.56)$$

where $tol = 10^{-10}$. The complete optimization search algorithm is summarized in Algorithm 1.

Algorithm 1 Solution of Problem 3.1.1 for initial guess $\boldsymbol{\eta}_{\alpha}^0$ and fixed α and T .

Input:

- α - regularization parameter
- T - length of time interval
- $\boldsymbol{\eta}_{\alpha}^0$ - initial guess
- N - resolution
- n - iteration number

Output:

- $\tilde{\boldsymbol{\eta}}_{\alpha,T}$ - optimal initial condition

$n \leftarrow 0$

while (3.56) is not satisfied **do**

Solve the Euler-Voigt system with initial condition $\boldsymbol{\eta}_{\alpha}^{(n)}$, see equation (1.4)

Solve the adjoint system to obtain $\mathbf{u}_{\alpha}^*$ and p^* , see equation (3.38)

Compute the gradient $\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_{\alpha}^{(n)})$, see equation (3.44)

Compute the search direction $\mathbf{d}^{(n)}$, see equation (3.52)

Compute the optimal step size τ_n , see equation (3.55)

Compute the next optimal initial condition $\boldsymbol{\eta}_{\alpha}^{(n+1)}$, see equation (3.51)

$n \leftarrow n + 1$

end while

$\tilde{\boldsymbol{\eta}}_{\alpha,T} \leftarrow \boldsymbol{\eta}_{\alpha}^{(n)}$

if the flow corresponding to $\tilde{\boldsymbol{\eta}}_{\alpha,T}$ is under-resolved, see equation (4.59) **then**

$N \leftarrow 2N$

$\boldsymbol{\eta}_{\alpha}^0 \leftarrow \tilde{\boldsymbol{\eta}}_{\alpha,T}$

Repeat Algorithm 1

end if

4 Numerical Methods

In this chapter we describe and validate the numerical techniques used to solve the optimization problem. In addition, we determine the parameters including the resolution, time step size, regularization parameter α , and Gevrey parameters s and σ .

4.1 Discretization

The Euler-Voigt (1.4) and adjoint system (3.38) were discretized in space using a standard pseudospectral Fourier method on a uniform grid with $N = 128^3, 256^3, 512^3, 1024^3$ grid points. Derivatives are evaluated in the Fourier space and nonlinear products are computed in the physical space. The standard 2/3 dealiasing rule is also applied.

Time stepping is performed with the explicit fourth-order Runge-Kutta (RK4) method. The determination of our time step size Δt is based on two factors. First, we consider the accuracy of the solution, determined by the relative error of the objective functional (3.32) relative to the results obtained with a minimum step size for a time

interval $T = 5$. Second, we consider the conservation of the α -energy (1.6), determined by the maximum relative error of $E_\alpha(t)$ relative to $E_\alpha(0)$, for $t \in [0, 5]$. These factors are analysed for each resolution and value of α for the TGV.

The goal of our optimization search is to identify initial conditions $\tilde{\eta}_{\alpha,T}$ that produce 'extreme' behaviour in solutions to the Euler-Voigt equations. Therefore, the TGV as the initial guess will by definition lead to a less extreme flow evolution than the initial conditions constructed at each iteration of the optimization search. To ensure stability, accuracy, and conservation of α -energy upon optimization, we must select a step size smaller than the minimum step size identified by the above criteria.

4.1.1 Numerical Stability & Accuracy

We begin by analysing the accuracy of the time discretization. To determine the accuracy of the discretization with a given time step size we solve the Euler-Voigt system for $T = 5$ with the TGV used as the initial condition and evaluate the objective functional (3.32). First, we do this for $\Delta t = 2^{-8}$ and designate this result as our 'correct' value Φ_c . We then compute the relative error of the objective functional for each value of Δt , denoted $\Phi_{\Delta t}$, to $\Phi_{\min}$

$$\epsilon_\Phi = \left| \frac{\Phi_{\min} - \Phi_{\Delta t}}{\Phi_{\min}} \right|. \quad (4.57)$$

In [20], the regularization parameter was in the range $\alpha = 12/1024, \dots, 36/1024$. We extend this range in both directions and use the regularization parameters $\alpha = 8/1024, \dots, 64/1024$. Table 4.1 shows the values for ϵ_Φ for $\alpha = 16/1024, \dots, 64/1024$. We include only these values of α in this table because they led to identical results

		α				
		16/1024	24/1024	32/1024	48/1024	64/1024
Δt	2^{-2}	7.00E-05	1.45E-05	4.80E-06	8.70E-07	1.58E-07
	2^{-3}	3.64E-06	7.89E-07	2.63E-07	5.03E-08	9.29E-09
	2^{-4}	2.02E-07	4.53E-08	1.52E-08	3.01E-09	5.63E-10
	2^{-5}	1.21E-08	2.70E-09	9.16E-10	1.83E-10	3.48E-11
	2^{-6}	8.63E-10	1.65E-10	5.99E-11	8.33E-12	2.05E-12
	2^{-7}	9.44E-11	9.42E-12	5.99E-12	0	0

Table 4.1: Results ϵ_Φ for $\alpha = 16/1024, \dots, 64/1024$.

for all four resolutions. This is due to the fact that the regularization term in (1.4) kept the solution well-resolved even at the lowest resolution, and so the accuracy did not increase with the higher resolutions. Table 4.2 shows the results of ϵ_Φ for $\alpha = 8/1024, 12/1024$. For these values the model becomes under-resolved at the lowest resolution and so accuracy does increase at the finer resolutions.

		α	
		8/1024	12/1024
Δt	2^{-2}	0.0045	2.66E-04
	2^{-3}	0.0011	1.46E-05
	2^{-4}	4.85E-04	1.30E-06
	2^{-5}	2.33E-04	3.19E-07
	2^{-6}	1.02E-04	1.23E-07
	2^{-7}	3.45E-05	4.05E-08

(a) $N = 128^3$

		α	
		8/1024	12/1024
Δt	2^{-2}	0.003	2.64E-04
	2^{-3}	1.82E-04	1.34E-05
	2^{-4}	9.46E-06	7.05E-07
	2^{-5}	5.12E-07	3.95E-08
	2^{-6}	3.00E-08	2.32E-09
	2^{-7}	1.96E-09	1.31E-10

(b) $N = 256^3, 512^3, 1024^3$

Table 4.2: Results ϵ_Φ for $\alpha = 8/1024, 12/1024$.

		α				
		16/1024	24/1024	32/1024	48/1024	64/1024
Δt	2^{-2}	8.73E-06	1.90E-06	5.95E-07	9.75E-08	1.64E-08
	2^{-3}	3.72E-07	7.68E-08	2.29E-08	3.81E-09	6.05E-10
	2^{-4}	1.76E-08	3.47E-09	9.74E-10	1.67E-10	2.45E-11
	2^{-5}	9.59E-10	1.74E-10	4.71E-11	9.79E-12	2.80E-12
	2^{-6}	7.55E-11	9.29E-12	3.27E-12	2.41E-12	1.96E-12
	2^{-7}	1.47E-11	2.45E-12	2.86E-12	3.12E-12	2.01E-12
	2^{-8}	6.00E-12	3.07E-12	3.12E-12	3.55E-12	2.21E-12

Table 4.3: The values of ϵ_α for $\alpha = 16/1024, \dots, 64/1024$.

We now examine the conservation of α -energy for the Euler-Voigt model. To determine the accuracy of discretization with a given time step size we solve the Euler-Voigt system for $T = 5$ with the TGV used as the initial condition and evaluate the maximum relative error of the α -energy

$$\epsilon_\alpha = \max_{t \in [0, T]} \left| \frac{E_\alpha(t) - E_\alpha(0)}{E_\alpha(0)} \right|. \quad (4.58)$$

Table 4.3 shows the values of ϵ_α for $\alpha = 16/1024, \dots, 64/1024$. Just as with the values of ϵ_Φ , we include only these values of α because they led to identical values for all four resolutions. Table 4.4 shows the values of ϵ_α for $\alpha = 8/1024, 12/1024$. The time step size of each model run was selected based on the results in these tables to ensure accuracy and conservation of the α -energy for the chosen value of α and resolution.

		α	
		8/1024	12/1024
Δt	2^{-2}	3.82E-04	3.35E-05
	2^{-3}	6.91E-05	1.60E-06
	2^{-4}	3.05E-05	1.26E-07
	2^{-5}	1.57E-05	3.14E-08
	2^{-6}	8.05E-06	1.41E-08
	2^{-7}	4.08E-06	6.99E-09
	2^{-8}	2.05E-06	3.50E-09

(a) $N = 128^3$

		α	
		8/1024	12/1024
Δt	2^{-2}	2.91E-04	3.33E-05
	2^{-3}	1.49E-05	1.50E-06
	2^{-4}	6.85E-07	7.21E-08
	2^{-5}	3.40E-08	3.80E-09
	2^{-6}	1.89E-09	2.14E-10
	2^{-7}	1.38E-10	1.20E-11
	2^{-8}	2.37E-11	2.80E-12

(b) $N = 256^3, 512^3, 1024^3$

Table 4.4: The values of ϵ_α for $\alpha = 8/1024, 12/1024$.

4.2 Energy Spectra and Validation of Numerical Resolution

The energy spectrum of the solution of (1.4) is defined as,

$$e(k, t) := \frac{1}{2} \sum_{j \leq |\mathbf{j}| < j+1} |\hat{\mathbf{u}}_{\mathbf{j}}(t)|^2, \quad k = 2\pi j, \quad \mathbf{j} \in \mathbb{Z}^3. \quad (4.59)$$

The regularization term in (1.4) acts as a filter which suppresses the growth of small-scale structures with length-scales smaller than the regularization parameter α [26]. Due to this property, the solution to (1.4) can remain well-resolved with relatively low resolution models. This is ideal for our analysis because it ensures that the model can fully capture the small scale structures of the fluid and so we can be certain that our objective functional (3.32) is accurately approximated.

We know that for smooth solutions with Gevrey regularity, cf. (2.21), the energy

spectrum (4.59) decays exponentially as the wavenumber $j = |\mathbf{j}|$ increases. Thus, whether or not a solution is well resolved can be assessed by examining the energy spectrum. Figures 4.1 and 4.2 show the energy spectrum of the solution corresponding to $\boldsymbol{\eta}_S$ with $\alpha = 8/1024$ and $\boldsymbol{\eta}_{TGV}$ with $\alpha = 64/1024$, respectively. The spectrum is plotted over the time interval $T = 5$. These figures demonstrate the exponential decay in the Fourier coefficients expected of a solution with Gevrey regularity. As the solutions evolve in time the rate of decay decreases as energy is transferred to small-scale structures. In Figure 4.1 a portion of the spectrum is cut-off by the 2/3 dealiasing filter. Due to the regularization term in (1.4), the solution in Figure 4.2 remains well resolved.




Figure 4.1: $e(k, t)$ vs. k starting from $\boldsymbol{\eta}_S$ for $\alpha = 8/1024$ and $N = 256^3$ at $t = 0$ (orange), $t = 1$ (cyan), $t = 2$ (brown), $t = 3$ (blue), $t = 4$ (green), $t = 5$ (purple).

In order to minimize the computational cost, the optimization searches were started from the initial guess at the lowest resolution, and ran until condition (3.56) was satisfied. If the flow over $[0, T]$ corresponding to the optimal initial condition $\tilde{\boldsymbol{\eta}}_{\alpha, T}$ was under-resolved, then the initial condition was up-scaled to the next higher resolution




Figure 4.2: $e(k, t)$ vs. k starting from $\boldsymbol{\eta}_{TGV}$ for $\alpha = 64/1024$ and $N = 128^3$ at $t = 0$ (orange), $t = 1$ (cyan), $t = 2$ (brown), $t = 3$ (blue), $t = 4$ (green), $t = 5$ (purple).

$2N$ by padding with zeros in Fourier space and then converting back to physical space. The optimization search was restarted, now with the up-scaled $\tilde{\boldsymbol{\eta}}_{\alpha, T}$ as the initial guess $\boldsymbol{\eta}_{\alpha}^0$. Optimization was stopped when criterion (3.56) was satisfied and the flow corresponding to $\tilde{\boldsymbol{\eta}}_{\alpha, T}$ remained well resolved, as verified by inspecting the energy spectrum (4.59). Using this method we can ensure that the final optimal initial conditions and the corresponding flows were all well-resolved, and so we can be certain our evaluation of the objective functional is accurate for all of the cases discussed in Chapter 5.

4.3 Validation of the Gradient of the Objective Functional (3.32)

In the previous sections we examined the accuracy of the objective functional (3.32) and α -energy (1.6) based on the regularization parameter α , resolution N , and time

step size Δt . In order to assess how accurately the gradient (3.44) of the objective functional is evaluated, we introduce the quantity [2, 30].

$$\kappa(\epsilon) = \frac{\epsilon^{-1} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)]}{\langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}}, \quad (4.60)$$

where the numerator is the first-order finite difference approximation of the Gâteaux differential (3.34), and the denominator is the Reisz form (3.35) of the differential.

A log-log plot of $|\kappa - 1|$ vs ϵ can be broken up into three regions, which are dominated by roundoff errors, truncation errors, and discretization errors. The roundoff errors are caused by subtractive cancellation errors due to finite-precision arithmetics in the numerator of 4.60 and so are approximately of order $\mathcal{O}(\epsilon^{-1})$. Therefore, (4.60) will be dominated by round off errors for small ϵ values. The truncation errors are caused by the finite difference formula to approximate the Gâteaux differential in the numerator and are of order $\mathcal{O}(\epsilon)$. Therefore, (4.60) will be dominated by the truncation errors for large values of ϵ . Between these two regions (4.60) will be dominated by the discretization errors in the approximation of the PDEs, which depend on the spatial resolution N and time step size Δt . Refining the discretization in both space and time will reduce these errors. Therefore, if the calculation of the gradient is accurate, the log-log plot of $|\kappa - 1|$ vs ϵ will be of the general form shown in Figure 4.3.

The quantity (4.60) was evaluated by fixing the spacial resolution at $N = 512^3$, to keep the model well resolved, and varying the time step size $\Delta t \in [2^{-2}, 2^{-8}]$. The simulations were run for the time window of $T = 5$ and $\epsilon \in [1\text{E-}14, 1]$. The values for $\alpha = 8/1024$ and $\alpha = 64/1024$ are shown in Figures 4.4 and 4.5, respectively. The results from the κ test reveal behaviour consistent with the expected behaviour




Figure 4.3: Schematic illustration of the log-log dependence of $\kappa(\epsilon)$ on ϵ , cf. (4.60).

illustrated in Figure 4.3. In particular, the values of $\kappa(\epsilon)$ obtained for intermediate values of ϵ , the plateaux in the figures, approach unity as the time step is refined. Therefore, we conclude that the numerical techniques used to compute the gradient are accurate.




Figure 4.4: Dependence of $\kappa(\epsilon)$ on ϵ for $\alpha = 8/1024$.




Figure 4.5: Dependence of $\kappa(\epsilon)$ on ϵ for $\alpha = 64/1024$.

4.4 Determination of the parameter defining the Gevrey space

In Section 2.2 we introduced the Gevrey space (2.21) in which we search for optimal initial conditions. The relative 'size' of the space is determined by two parameters, $s \geq 3$ and $\sigma > 0$. We use a fixed value of $s = 3$. In order to determine an optimal value of σ , we solve the optimization problem for $\alpha = 16/1024$, $T = 5$, and different values of $\sigma = 10^{-1}, \dots, 10^{-5}$. The resulting values of the objective functional are compared in Figure 4.6.

The results of Figure 4.6 show that the number of iterations it took to reach the maximum value does change depending on the value of σ , but there was not a significant variation in the value of the objective functional. This was also the result of the tests conducted for different values of α . These results are consistent with those of [30] which found that the optimization results did not significantly change so long as $\sigma < 10^{-1}$. Therefore, we make the same selection as in [30] and choose $\sigma = 10^{-5}$.




Figure 4.6: $\Phi_{\alpha,T}(\eta_{\alpha,T}^{(n)})$ vs iteration number n for $\alpha = 16/1024$, $T = 5$, and different indicated values of σ .

5 Results

Utilizing the methods described in the previous chapters, we present numerical results consistent with singularity formation in the Euler flow starting from $\boldsymbol{\eta}_{TGV}$, $\boldsymbol{\eta}_S$ and $\boldsymbol{\eta}_L$. However, to obtain these results we must extend the time interval to $T = 25$. We also present numerical results that a singularity does form in the Euler flow corresponding to the optimal initial conditions $\tilde{\boldsymbol{\eta}}_{8,3}$ and $\tilde{\boldsymbol{\eta}}_{8,5}$, within the corresponding time intervals, using $\boldsymbol{\eta}_{TGV}$ as the initial guess. Additionally, our numerical results indicate that a singularity does not form in the flow corresponding to the optimal initial conditions $\tilde{\boldsymbol{\eta}}_{8,3}$ or $\tilde{\boldsymbol{\eta}}_{8,5}$ within the corresponding time intervals, using $\boldsymbol{\eta}_S$ as the initial guess. However, by starting from $\tilde{\boldsymbol{\eta}}_{8,5}$ and evolving the solution over the time interval $T = 25$, we obtain results consistent with singularity formation in the Euler flow. Finally, we present numerical evidence that a singularity DOES(NT) form in the Euler flow corresponding to the optimal initial conditions $\boldsymbol{\eta}_{8,3}$ or $\boldsymbol{\eta}_{8,5}$, using $\boldsymbol{\eta}_L$ as the initial guess.

5.1 Singularity Detection

Theorem 2.1.3 provides the criteria we use to determine if a singularity forms in the Euler flow corresponding to initial condition $\boldsymbol{\eta}$, where $\boldsymbol{\eta}$ is the optimal initial condition found in the limit $\alpha \rightarrow 0$ by solving Problem 3.1.1 as described in Chapter 3. In practice, we take $\boldsymbol{\eta}$ to be the optimal initial condition found for the smallest value of α for which the calculations were well resolved, i.e, $\tilde{\boldsymbol{\eta}}_{8,T}$. First, we check if the sequence of optimal initial conditions, found by solving Problem 3.1.1 for decreasing α , converges to a well-defined limit, i.e, if they satisfy (2.8). If this condition is satisfied, we then check whether (2.12) is satisfied and $\alpha^2 \Phi_{\alpha,T}$ converges to a positive limit as $\alpha \rightarrow 0$.

Simply plotting the left-hand side (LHS) in (2.8) and (2.12) as functions of α will not be sufficient for determining if each condition is satisfied. This is due to the fact that it is difficult to detect if these expressions are converging to zero or to some small positive limit. In order to make this determination we use the method of [20], which proposes the following ansatz,

$$\sup_{t \in [0,T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \approx \mathcal{O}(\alpha^p), \quad (5.61)$$

for $T > 0$ sufficiently large and for some power p . From this and Theorem 2.1.3, in order for a singularity to form in the Euler flow, it must be the case that the increase in the objective functional offsets the decrease of α , which can only occur if $p \leq -1$. However, if $p < -1$, then the gradient term in (1.6) will become unbounded as $\alpha \rightarrow 0$. This would mean the α -energy is not conserved. Therefore, for a singularity to form,

it must be the case that strictly $p = -1$. To find the value of p , we create a log-log plot of $\max_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α . In order to simplify the visualization of the evolution of this quantity over time, the figures included here will only include the results at $t = 0, 1, \dots, T$. The value of p is defined as the steepest slope between any two values of α at any point in the time window.

We utilize the same method to determine if (2.8) is satisfied. However, in this case the condition is satisfied if the quantity does converge to zero. The condition is a sum of a energy term, $\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2$ and a enstrophy term, $\|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2$, which is scaled by α^2 . From Poincare's inequality, if the enstrophy term diminishes to zero as $\alpha \rightarrow 0$, then so too does the energy term and so (2.8) is satisfied [13]. However, if the enstrophy term diverges, but does so more slowly than α diminishes, then (2.8) is only satisfied if the energy term also goes to zero with α . Finally, if the enstrophy term diverges more quickly than α diminishes then (2.8) is not satisfied. Therefore, we propose the following two ansatze.

$$\|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2} \approx \mathcal{O}(\alpha^q). \quad (5.62)$$

and

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2} \approx \mathcal{O}(\alpha^r), \quad (5.63)$$

By creating log-log plots of the LHS in (5.62) and (5.63) vs α , the values of q and r are given by the steepest slope between any two values of α at any point in time for (5.62) and (5.63), respectively. For (2.8) to be satisfied, it must be the case that either $r > 0$, or $-1 < r \leq 0$ and $q > 0$.

5.2 Results Obtained Using the Initial Guesses

We begin by presenting our results obtained using the initial guesses, $\boldsymbol{\eta}_{TGV}$, $\boldsymbol{\eta}_S$, and $\boldsymbol{\eta}_L$. These serve as a baseline comparison with the results obtained by solving Problem 3.1.1. Without optimization, the initial conditions for all values of α are simply the initial guess. In this case, it is trivially true that condition (2.8) is satisfied, and so we only need to compute the value of p to determine if a singularity forms in the Euler flow corresponding to the initial guess.

5.2.1 Results Obtained Using $\boldsymbol{\eta}_{TGV}$

A visualization of the vorticity of $\boldsymbol{\eta}_{TGV}$ is shown in Figure 3.1. Figure 5.1 shows the vorticity of $\omega = \nabla \times \mathbf{u}_\alpha(T = 3; \boldsymbol{\eta}_{TGV})$ for $\alpha = 8/1024$ and Figure 5.2 shows the vorticity of $\omega = \nabla \times \mathbf{u}_\alpha(T = 5; \boldsymbol{\eta}_{TGV})$. The figures show that the areas of high vorticity are concentrated into smaller area as the flow evolves and that the peak of the vorticity increases in magnitude.

Figures 5.3 and 5.4 show that for $\boldsymbol{\eta}_{TGV}$ and $T = 3$, the steepest slope achieved at any point in the time interval is $p = -0.225$ and this slope occurs at $t = 3$. Therefore, these results indicate that a singularity does not form in the Euler flow within the time interval $[0, 3]$ from $\boldsymbol{\eta}_{TGV}$. They also show that the slope is still decreasing at the end of the time window, indicating that they may reach a slope of $p = -1$ at some later point in time. This result is consistent with the results of [5, 20] as the time interval is too short to reach the expected blow-up time of $T^* = 4$.




Figure 5.1: Isosurfaces of vorticity $\omega = \nabla \times \mathbf{u}_\alpha(T = 3; \boldsymbol{\eta}_{TGV})$. (a) magnitude $|\omega(\mathbf{x})|$, and components (b) $|\omega_1|$, (c) $|\omega_2|$, and (d) $|\omega_3|$.

Figures 5.5 and 5.6 show that for $\boldsymbol{\eta}_{TGV}$ and $T = 5$, the steepest slope achieved at any point in the time is $p = -0.417$ and that this slope occurs at $t = 5$. This slope is larger than the p value for the shorter time window. However, it is still not close to the $p = -1$ required by (5.61). Therefore, these results indicate that a singularity does not form in the Euler flow within the time interval $[0, 5]$ from $\boldsymbol{\eta}_{TGV}$. This result conflicts with the results of [5, 20], which indicate a singularity should form at $T^* \approx 4$.

5.2.2 Delayed Singularity Formation

Our results indicate that a singularity does not form in the Euler flow corresponding to $\boldsymbol{\eta}_{TGV}$ within the time interval $[0, 5]$, which conflicts with the results of [20]. Their analysis also examined the Euler-Voigt equations in order to search for singularity




Figure 5.2: Isosurfaces of vorticity $\omega = \nabla \times \mathbf{u}_\alpha(T = 5; \boldsymbol{\eta}_{TGV})$. (a) magnitude $|\omega(\mathbf{x})|$, and components (b) $|\omega_1|$, (c) $|\omega_2|$, and (d) $|\omega_3|$.




Figure 5.3: Solution of the Euler-Voigt system for $T = 3$ starting from $\boldsymbol{\eta}_{TGV}$. Log-log plot of $\max_{t \in [0, T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α at $t = 0$ (blue) to $t = 3$ (red), Steepest slope $p = -0.225$ (black), slope of $p = -1$ (dotted black)




Figure 5.4: Solution of the Euler-Voigt system for $T = 3$ starting from $\boldsymbol{\eta}_{TGV}$. Plot of the slope of $(\max_{t \in [0, T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2})$ vs α vs time for each α .




Figure 5.5: Solution of the Euler-Voigt system for $T = 5$ starting from $\boldsymbol{\eta}_{TGV}$. Log-log plot of $\max_{t \in [0, T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α at $t = 0$ (blue) to $t = 5$ (red), Steepest slope $p = -0.417$ (black), slope of $p = -1$ (dotted black)




Figure 5.6: Solution of the Euler-Voigt system for $T = 5$ starting from $\boldsymbol{\eta}_{TGV}$. Plot of the slope of $(\max_{t \in [0, T']} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \text{ vs } \alpha)$ vs time at each α .

formation in Euler from $\boldsymbol{\eta}_{TGV}$. As shown in Figure 3 of [20], they obtained a result of $p \approx 1$, which indicates that blow-up does occur in Euler flows starting from $\boldsymbol{\eta}_{TGV}$. Additionally, the minimum of p occurred at a time consistent with the results of [5] at $T^* \approx 4.2$. However, their model used a 3D unit torus ($L = 1$) and a coefficient of $V_0 = 1$ in (2.29), rather than the $V_0 = 1/2\pi$ we use to match the scaling of [5]. Because of this, the time interval of their model scaled by a factor of 2π to the time interval of [5]. Therefore, the results of [20] do indicate that a singularity forms in the Euler flow corresponding to $\boldsymbol{\eta}_{TGV}$. However, the blow-up time, in our model, should be $T^* \approx 26.3894$.

limited to the range $\alpha = 16/1024, \dots, 64/1024$ in order to keep the results well resolved.

Figures 5.7 and 5.8 show that for $\boldsymbol{\eta}_{TGV}$ and $T = 25$, the steepest slope achieved at any point in the time is $p = -1.026$ and that this slope occurs at $t \approx 21.25$. This slope satisfies (5.61). Therefore, these results indicate that a singularity does form in the Euler flow within the time interval $[0, 25]$ from $\boldsymbol{\eta}_{TGV}$.




Figure 5.7: TGV for $T = 25$. Log-log plot of $\max_{t \in [0, T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}$ vs α at $t = 0$ (blue) to $t = 25$ (red), Steepest slope $p =$ (black), slope of $p = -1$ (dotted black)




Figure 5.8: TGV for $T = 25$. Plot of the slope of $(\max_{t \in [0, T]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \text{ vs } \alpha)$ vs time at each α .

Theorem 2.1.3 states that if conditions (2.8) and (2.12) are satisfied, then a singularity forms in the Euler flow corresponding to $\boldsymbol{\eta}$ in the interval $[0, T^*]$. This theorem makes no prediction as to the exact moment the singularity forms, only that it occurs somewhere within the interval $[0, T^*]$. Our results indicate that a singularity does form in the Euler flow corresponding to $\boldsymbol{\eta}_{TGV}$ in the interval $[0, 25]$. The results of [5] indicate a singularity forms in the Euler flow corresponding to $\boldsymbol{\eta}_{TGV}$ at $T^* \approx 4$, which is in the interval $[0, 25]$. Therefore, while it is unexpected that the time interval required to satisfy Theorem 2.1.3 is much larger than the results of [5] would suggest, these results do not contradict one another.

6 Conclusion

Every thesis also needs a concluding chapter

Some as of yet known process delaying the formation of a singularity.

A Function Spaces

In this Appendix we introduce the Lebesgue space, **the space of continuous functions**, and the Sobolev space. These function spaces are used to define the initial conditions and solutions of the Euler and Euler-Voigt equations, the criteria for singularity formation, the objective functional, the gradient of the objective functional, and the Riemannian conjugate gradient method.

The Lebesgue space $L^p(\Omega)$ consists of real-valued measurable functions $\mathbf{u}$, defined on Ω , for which the norm

$$\|\mathbf{u}\|_{L^p} = \left(\int_{\Omega} |\mathbf{u}|^p d\mathbf{x} \right)^{1/p},$$

is bounded, where $1 \leq p < \infty$. For $p = \infty$, the norm is given by

$$\|\mathbf{u}\|_{L^\infty} = \operatorname{ess\,sup}_{\mathbf{x} \in \Omega} |\mathbf{u}(\mathbf{x})|,$$

where $\operatorname{ess\,sup}$ is the essential supremum of $|\mathbf{u}|$ on Ω [15, 29].

Let $\alpha = (\alpha_1, \dots, \alpha_n)$ be an n -tuple of nonnegative integers α_i with the magnitude $|\alpha| = \sum_{i=1}^n \alpha_i$. For any nonnegative integer k , the space $C^k(\Omega)$ is the space of

functions $\mathbf{u}$ which, together with all partial derivatives $\partial^\alpha \mathbf{u}$ of orders $|\alpha| \leq k$, are continuous on Ω [15].

For $s \in \mathbb{R}^+$, the L^2 -based Sobolev space $H^s(\Omega)$ is the space of all functions for which

$$\|\mathbf{u}\|_{H^s}^2 := (2\pi)^3 \sum_{\mathbf{k} \in \mathbb{Z}^3} (1 + |\mathbf{k}|^{2s}) |\widehat{\mathbf{u}}_{\mathbf{k}}|^2.$$

is finite [15, 29]. In physical space, the norm is defined as

$$\|\mathbf{u}\|_{H^s}^2 := \sum_{|\alpha| \leq s} \frac{s!}{\alpha!} \|d^\alpha \mathbf{u}\|_{L^2}^2,$$

where $\alpha! = \alpha_1! \alpha_2! \alpha_3!$, and $d^\alpha \mathbf{u}$ is defined as

$$d^\alpha \mathbf{u} = \frac{\partial^{|\alpha|}}{\partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}} \mathbf{u}.$$

For any function $\mathbf{u} \in H^s(\mathbb{T}^3)$ the $\dot{H}^s$ seminorm is defined by

$$\|\mathbf{u}\|_{\dot{H}^s}^2 := (2\pi)^3 \sum_{\mathbf{k} \in \mathbb{Z}^3} |\mathbf{k}|^{2s} |\widehat{\mathbf{u}}_{\mathbf{k}}|^2.$$

The seminorm in physical space is defined as

$$\|\mathbf{u}\|_{\dot{H}^s}^2 := \sum_{|\alpha|=s} \frac{s!}{\alpha!} \|d^\alpha \mathbf{u}\|_{L^2}^2.$$

B Proof of Relation (2.25)

In this Appendix we prove relation (2.25) which shows that the parameter σ determines the relative 'size' of the Gevrey space in which we search for optimal initial conditions.

Proof. Let $\sigma_1, \sigma_2 \in \mathbb{R}$ be defined such that $0 < \sigma_1 < \sigma_2$. Therefore, from (2.22)

$$\begin{aligned} \|\mathbf{v}\|_{G^{\sigma_2}} &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_2|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} \\ &\geq \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_1|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^{\sigma_1}} \\ &\geq \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^0 \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^0} \\ &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{H^s}. \end{aligned}$$

Thus,

$$\|\mathbf{v}\|_{G^{\sigma_2}} \geq \|\mathbf{v}\|_{G^{\sigma_1}} \geq \|\mathbf{v}\|_{G^0} = \|\mathbf{v}\|_{H^s}.$$

It follows from this that if $\|\mathbf{v}\|_{G^{\sigma_2}} < \infty$ then $\|\mathbf{v}\|_{G^{\sigma_1}} < \infty$ and so if $v \in G^{\sigma_2}$ then $v \in G^{\sigma_1}$. However, the converse is not true. The same argument shows that if $v \in G^{\sigma_1}$ then $v \in G^0$, which is equivalent to $v \in H^s$. Therefore, $G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s$.

□

C Rescaling the Euler equations

In this Appendix we discuss the scaling symmetry of the Euler-Voigt equation (1.4). Given a solution $\mathbf{u}$ of the Euler equations (1.2), we construct a new scaled solution with

$$\tau = \lambda t, \quad \text{and} \quad \mathbf{y} = \beta \mathbf{x}.$$

where $\lambda, \beta > 0$ are constants. Our new solution $\mathbf{v}$ is defined as

$$\mathbf{v} = \mathbf{v}(\mathbf{y}, \tau) = \gamma \mathbf{u}(\mathbf{y}/\beta, \tau/\lambda).$$

Therefore

$$\partial_t \mathbf{u} = \frac{1}{\gamma} \frac{\partial \mathbf{v}}{\partial \tau} \frac{d\tau}{dt} = \frac{\lambda}{\gamma} \partial_\tau \mathbf{v}.$$

and

$$\nabla \mathbf{u} = \frac{1}{\gamma} \nabla \mathbf{v} \frac{d\mathbf{y}}{d\mathbf{x}} = \frac{\beta}{\gamma} \nabla \mathbf{v}.$$

Substituting $\mathbf{v}$ into equation (1.2a)

$$\frac{\lambda}{\gamma} \partial_\tau \mathbf{v} + \frac{\beta}{\gamma^2} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q = 0,$$

where q is defined as whatever pressure is required to satisfy the incompressibility condition. This can be rearranged to the form,

$$\frac{\lambda\gamma}{\beta}\partial_\tau\mathbf{v} + (\mathbf{v} \cdot \nabla)\mathbf{v} + \nabla q' = 0.$$

Therefore, our function $\mathbf{v}$ can only be a solution to the Euler equations if

$$\frac{\lambda\gamma}{\beta} = 1.$$

For the Taylor Green Vortex with $L = 2\pi$ and $V_0 = 1$, let

$$\gamma = \lambda = \beta = 1.$$

In order to rescale this to the domain with $L = 1$ we must set $\beta = 1/2\pi$. Additionally, we want our rescaled solution to have the same time scale as the original solution and so we must set $\lambda = 1$. Therefore, for our scaled solution $\mathbf{v}$ to be a solution of the Euler equations we must set $\gamma = V_0 = 1/2\pi$.

D Rescaling the Euler-Voigt equations

In this Appendix we discuss the scaling symmetry of the Euler-Voigt equation (1.4). Given a solution $\mathbf{u}_\alpha$ of the Euler-Voigt equations (1.4), we construct a new scaled solution with the same domain size by constructing a new scaled solution with

$$\tau = \lambda t, \quad \kappa = \beta \alpha, \quad \mathbf{y} = \beta \mathbf{x}.$$

where $\lambda, \beta > 0$ are constants. The variables $\mathbf{x}$ and α are both scaled by β because they are both spatial and we want to keep the relative size of the filter to the domain size fixed. Our new solution $\mathbf{v}$ is defined as

$$\mathbf{v} = \mathbf{v}(\mathbf{y}, \tau) = \gamma \mathbf{u}_\alpha(\mathbf{y}/\beta, \tau/\lambda).$$

Therefore

$$\partial_t \mathbf{u}_\alpha = \frac{1}{\gamma} \frac{\partial \mathbf{v}}{\partial \tau} \frac{d\tau}{dt} = \frac{\lambda}{\gamma} \partial_\tau \mathbf{v}.$$

and

$$\nabla \mathbf{u}_\alpha = \frac{1}{\gamma} \nabla \mathbf{v} \frac{d\mathbf{y}}{d\mathbf{x}} = \frac{\beta}{\gamma} \nabla \mathbf{v}.$$

and

$$\partial_t \Delta \mathbf{u}_\alpha = \partial_t \frac{\beta^2}{\gamma} \Delta \mathbf{v} = \frac{\lambda \beta^2}{\gamma} \partial_\tau \Delta \mathbf{v}.$$

Substituting $\mathbf{v}$ into equation (1.4a)

$$-\alpha^2 \frac{\lambda \beta^2}{\gamma} \partial_\tau \Delta \mathbf{v} + \frac{\lambda}{\gamma} \partial_\tau \mathbf{v} + \frac{\beta}{\gamma^2} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q = 0,$$

where q is defined as whatever pressure is required to satisfy the incompressibility condition. This can be rearranged to the form,

$$-\kappa^2 \partial_t \Delta \mathbf{v} + \partial_\tau \mathbf{v} + \frac{\beta}{\lambda \gamma} (\mathbf{v} \cdot \nabla) \mathbf{v} + \nabla q' = 0.$$

Therefore, our function $\mathbf{v}$ can only be a solution to the Euler-Voigt equations if

$$\frac{\beta}{\lambda \gamma} = 1.$$

For the Taylor Green Vortex with $L = 2\pi$ and $V_0 = 1$, let $\gamma = \lambda = \beta = 1$. In order to rescale this to the domain with $L = 1$ we must set $\beta = 1/2\pi$. Additionally, we want our rescaled solution to have the same time scale as the original solution and so we must set $\lambda = 1$. Therefore, for our scaled solution $\mathbf{v}$ to be a solution of the Euler-Voigt equations we must set $\gamma = V_0 = 1/2\pi$.

E Evaluation of $\|\boldsymbol{\eta}\|_{\dot{H}^s}$ for the Taylor Green Vortex

In this Appendix we compute the $\dot{H}^s$ seminorm for the Taylor Green Vortex. The $\dot{H}^1$ seminorm is required for properly scaling the initial conditions of the Euler and Euler-Voigt equations and the $\dot{H}^3$ is required for comparing our results to those of [30].

The 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is given in (2.29). To simplify the notation we drop the subscript TGV . The formula for $\|\boldsymbol{\eta}\|_{\dot{H}^s}$ is

$$\|\boldsymbol{\eta}\|_{\dot{H}^s}^2 := \sum_{|\alpha|=s} \frac{s!}{\alpha!} \|d^\alpha \boldsymbol{\eta}\|_{L^2}^2.$$

This can be expressed in integral form as

$$\|\boldsymbol{\eta}\|_{\dot{H}^s}^2 = \sum_{|\alpha|=s} \frac{s!}{\alpha!} \left[\sum_{i=1}^3 \int_{\Omega} \left(\frac{\partial^{|\alpha|}}{\partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}} \eta_i \right)^2 d\mathbf{x} \right].$$

For the TGV, the third term η_3 is zero and the partial derivative of the other two terms η_1 and η_2 are

$$\frac{\partial^{|\alpha|}}{\partial_1^{\alpha_1} \partial_2^{\alpha_2} \partial_3^{\alpha_3}} \eta_i = V_0 \left(\frac{2\pi}{L} \right)^{|\alpha|} f(x_1) g(x_2) h(x_3), \quad i = 1, 2,$$

where f , g , and h are either $\sin(2\pi x_j/L)$ or $\cos(2\pi x_j/L)$. This allows us to simplify the equation to

$$\|\boldsymbol{\eta}\|_{\dot{H}^s}^2 = \sum_{|\alpha|=s} \frac{s!}{\alpha!} \left[\sum_{i=1}^2 V_0^2 \left(\frac{2\pi}{L} \right)^{2|\alpha|} \int_{\Omega} f(x_1)^2 g(x_2)^2 h(x_3)^2 d\mathbf{x} \right].$$

Because the integrand is separable, the integral is equal to the product of,

$$\int_0^L \cos^2(2\pi x_j/L) dx_j = \frac{L}{2}, \quad \text{and} \quad \int_0^L \sin^2(2\pi x_j/L) dx_j = \frac{L}{2}.$$

Therefore, the equation can be simplified to

$$\|\boldsymbol{\eta}\|_{\dot{H}^s}^2 = \sum_{|\alpha|=s} \frac{s!}{\alpha!} \left[2V_0^2 \left(\frac{2\pi}{L} \right)^{2|\alpha|} \left(\frac{L}{2} \right)^3 \right] = \sum_{|\alpha|=s} \frac{s!}{\alpha!} \left[2^{2|\alpha|-2} V_0^2 \pi^{2|\alpha|} L^{3-2|\alpha|} \right].$$

For $s = 1$, there are three tuples α with degree $|\alpha| = 1$. These are the three permutations of $\alpha = (1, 0, 0)$ with $\alpha! = 1$. Therefore, we obtain the result

$$\|\boldsymbol{\eta}\|_{\dot{H}^1} = \sqrt{3LV_0}\pi.$$

For $s = 3$, there are ten tuples α with degree $|\alpha| = 3$. These are $\alpha = (1, 1, 1)$ with $\alpha! = 1$, the three permutations of $\alpha = (3, 0, 0)$ with $\alpha! = 6$, and the six permutations

of $\alpha = (1, 2, 0)$ with $\alpha! = 2$. Therefore,

$$\|\boldsymbol{\eta}\|_{\dot{H}^3}^2 = \left[\frac{3!}{1} + 3\frac{3!}{6} + 6\frac{3!}{2} \right] 2^4 V_0^2 \pi^6 L^{-3}.$$

From this, we obtain the result

$$\|\boldsymbol{\eta}\|_{\dot{H}^3} = \frac{12\sqrt{3}V_0\pi^3}{\sqrt{L^3}}.$$

In Appendices C and D we derived the scaling property of the Euler and Euler-Voigt equations, respectively. This property requires that $\beta/\lambda\gamma = 1$, where β is the scaling factor in space, λ is the scaling factor in time, and γ is the scaling factor of the 'size' of the solution. The optimization problem of [30] searched for optimal initial conditions with a fixed $\|\boldsymbol{\eta}\|_{\dot{H}^3} = 1$ on a domain with 3D periodic cube with side length $L = 1$. Therefore, in this model, $V_0 = 1/(12\sqrt{3}\pi^3)$. Let $\beta = \gamma = \lambda = 1$.

Our optimization problem searches for optimal initial conditions with a fixed $\|\boldsymbol{\eta}\|_{\dot{H}^1} = \sqrt{3}/2$. Our search is also performed on a domain with $L = 1$. Using these results, we obtain the relative scaling of our model to that of [30]. These are $\beta = 1$ and $\gamma = 1/6\sqrt{3}\pi^2$. Thus, to satisfy the scaling property, $\gamma = 6\sqrt{3}\pi^2$. Therefore, one unit of time in our model is equivalent to $6\sqrt{3}\pi^2$ units of time in the model of [30].

F Evaluation of $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ for the Taylor Green Vortex

In this Appendix we compute the L^∞ norm of the vorticity field for the Taylor Green Vortex. It is required for properly scaling the initial conditions of the Euler and Euler-Voigt equations.

The 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is given in (2.29). To simplify the notation we drop the subscript TGV and substitute in

$$x' = 2\pi x/L, \quad y' = 2\pi y/L, \quad z' = 2\pi z/L.$$

The vorticity of $\boldsymbol{\eta}$ is

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} \partial_2 \eta_3 - \partial_3 \eta_2 \\ \partial_3 \eta_1 - \partial_1 \eta_3 \\ \partial_1 \eta_2 - \partial_2 \eta_1 \end{bmatrix}$$

where ∂_i is the partial derivative with respect to the i -th component and η_j is the j -th component of $\boldsymbol{\eta}$. Substituting in the partial derivatives

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} 0 - (-(2\pi V_0/L) \cos(x') \sin(y') \sin(z')) \\ -(2\pi V_0/L) \sin(x') \cos(y') \sin(z') - 0 \\ (2\pi V_0/L) \sin(x') \sin(y') \cos(z') + (2\pi V_0) \sin(x') \sin(y') \cos(z'), \end{bmatrix}$$

which simplifies to

$$\nabla \times \boldsymbol{\eta}_{TG} = \begin{bmatrix} (2\pi V_0/L) \cos(x') \sin(y') \sin(z') \\ -(2\pi V_0/L) \sin(x') \cos(y') \sin(z') \\ (4\pi V_0/L) \sin(x') \sin(y') \cos(z') \end{bmatrix}.$$

The $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ norm is the maximum absolute value of the components of $\nabla \times \boldsymbol{\eta}$.

$$\begin{aligned} \|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} &= \max(|(2\pi V_0/L) \cos(x') \sin(y') \sin(z')| \\ &\quad , |-(2\pi V_0/L) \sin(x') \cos(y') \sin(z')| \\ &\quad , |(4\pi V_0/L) \sin(x') \sin(y') \cos(z')|) \end{aligned}$$

The maximums are

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = \max((2\pi V_0/L), (2\pi V_0/L), (4\pi V_0/L))$$

Therefore,

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = 4\pi V_0/L.$$

G Linearization of the Euler-Voigt System

In this Appendix we derive the linearization of the Euler-Voigt equations. It is used to obtain the adjoint system and the gradient of the objective functional.

The Euler-Voigt equations are defined as,

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p_\alpha = 0 \\ \nabla \cdot \mathbf{u}_\alpha = 0 \\ \mathbf{u}_\alpha|_{t=0} = \boldsymbol{\eta}_\alpha. \end{cases}$$

Substituting in $\mathbf{u}_\alpha = \mathbf{u}_\alpha + \mathbf{u}'_\alpha$, $\boldsymbol{\eta}_\alpha = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha$, and $p_\alpha = p_\alpha + p'_\alpha$,

$$\begin{cases} -\alpha^2 \partial_t \Delta (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \partial_t (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + ((\mathbf{u}_\alpha + \mathbf{u}'_\alpha) \cdot \nabla) (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \nabla (p_\alpha + p'_\alpha) = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha. \end{cases}$$

By expanding each term we obtain,

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha - \alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \\ \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p_\alpha + \nabla p'_\alpha = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta}_\alpha + \boldsymbol{\eta}'_\alpha. \end{cases}$$

Substituting in the Euler-Voigt equations we simplify to

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p'_\alpha = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

Linearizing the system by removing non-linear terms, we derive the linearized Euler-Voigt system

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p'_\alpha = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

This can be rearranged to the form

$$\begin{cases} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p'_\alpha) = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'_\alpha. \end{cases}$$

H Derivation of the Adjoint System

In this Appendix we derive the adjoint system (3.38) using the duality-pairing relation (3.39). We drop the subscript α from $\mathbf{u}_\alpha$, $\mathbf{u}_\alpha^*$, and $\mathbf{u}'_\alpha$ to simplify the notation. Additionally, we use a simplified integral notation where $\Theta = (\Omega \times [0, T])$, i.e.

$$\int_0^T \int_\Omega \mathbf{u} \, d\mathbf{x} \, dt = \int_\Theta \mathbf{u} \, d\mathbf{x} \, dt.$$

We begin by expanding (3.39) into individual integrals which will be evaluated separately

$$\left(\mathcal{L} \begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} \right) = \int_\Theta \partial_t \mathbf{u}' \cdot \mathbf{u}^* \, d\mathbf{x} \, dt \tag{H.64a}$$

$$+ \int_\Theta (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* \, d\mathbf{x} \, dt \tag{H.64b}$$

$$+ \int_\Theta (I - \alpha^2 \Delta)^{-1} (\mathbf{u}' \cdot \nabla \mathbf{u}) \cdot \mathbf{u}^* \, d\mathbf{x} \, dt \tag{H.64c}$$

$$+ \int_\Theta \nabla p' \cdot \mathbf{u}^* \, d\mathbf{x} \, dt \tag{H.64d}$$

$$+ \int_\Theta (\nabla \cdot \mathbf{u}') p^* \, d\mathbf{x} \, dt. \tag{H.64e}$$

Integrating (H.64a) by part with respect to time gives

$$\int_{\Theta} \partial_t \mathbf{u}' \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Omega} \mathbf{u}^* \cdot \mathbf{u}'|_{t=0}^{t=T} d\mathbf{x} - \int_{\Theta} \partial_t \mathbf{u}^* \cdot \mathbf{u}' d\mathbf{x} dt.$$

To evaluate (H.64b) we first split the integrand into its components, and use Einstein's notation to simplify.

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Theta} (I - \alpha^2 \Delta)^{-1} u_i \partial_i u'_j u_j^* d\mathbf{x} dt.$$

Integrating (H.64b) by part with respect to space gives

$$\begin{aligned} \int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt &= \int_0^T (I - \alpha^2 \Delta)^{-1} u_i u_j^* u'_j|_{\partial\Omega} dt - \\ &\quad \int_{\Theta} u'_j \partial_i ((I - \alpha^2 \Delta)^{-1} u_i u_j^*) d\mathbf{x} dt. \end{aligned}$$

Because Ω has periodic boundaries, the first integral is zero, and we can expand the second integral.

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Theta} u'_j [u_i \partial_i ((I - \alpha^2 \Delta)^{-1} u_i) - (I - \alpha^2 \Delta)^{-1} u_j^* \partial_i u_i] d\mathbf{x} dt.$$

The flow is divergence free and so $\partial_i u_i = 0$. We can convert the second integral back to vector notation to obtain

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = - \int_{\Theta} (\mathbf{u} \cdot \nabla (I - \alpha^2 \Delta)^{-1} \mathbf{u}^*) \cdot \mathbf{u}' d\mathbf{x} dt.$$

To evaluate (H.64c) we first split the integrand into its components, and use Einstein's notation to simplify.

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Theta} (I - \alpha^2 \Delta)^{-1} u'_i (\partial_i u_j) u_j^* d\mathbf{x} dt.$$

The flow is divergence free and so we can convert express this as

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Theta} (I - \alpha^2 \Delta)^{-1} \partial_i (u'_i u_j) u_j^* d\mathbf{x} dt.$$

Integrating (H.64c) by part with respect to space gives.

$$\begin{aligned} \int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt &= \int_0^T \int_{\Theta} (I - \alpha^2 \Delta)^{-1} u_j^* u'_i u_j dt - \\ &\quad \int_{\Theta} u'_i u_j \partial_i (I - \alpha^2 \Delta)^{-1} u_j^* d\mathbf{x} dt \end{aligned}$$

Because Ω has periodic boundaries, the first integral is zero. We can convert the second integral back to vector notation to obtain

$$\int_{\Theta} (I - \alpha^2 \Delta)^{-1} (\mathbf{u} \cdot \nabla \mathbf{u}') \cdot \mathbf{u}^* d\mathbf{x} dt = - \int_{\Theta} (\nabla (I - \alpha^2 \Delta)^{-1} \mathbf{u}^*)^\top \mathbf{u} \cdot \mathbf{u}' d\mathbf{x} dt.$$

To evaluate (H.64d) we first split the integrand into its components, and use Einstein's notation to simplify.

$$\int_{\Theta} \nabla p' \cdot \mathbf{u}^* d\mathbf{x} dt = \int_{\Theta} \partial_i p' u_i^* d\mathbf{x} dt.$$

Integrating (H.64d) by part with respect to space gives

$$\int_{\Theta} \nabla p' \cdot \mathbf{u}^* d\mathbf{x} dt = \int_0^T \mathbf{u}^* p'|_{\partial\Omega} dt - \int_{\Theta} p' \partial_i u_i^* d\mathbf{x} dt.$$

Because Ω has periodic boundaries, the first integral is zero, and we can convert the second integral back to vector notation to obtain

$$\int_{\Theta} \nabla p' \cdot \mathbf{u}^* d\mathbf{x} dt = - \int_{\Theta} p' (\nabla \cdot \mathbf{u}^*) d\mathbf{x} dt.$$

To evaluate (H.64e) we first split the integrand into its components, and use Einstein's notation to simplify.

$$\int_{\Theta} (\nabla \cdot \mathbf{u}') p^* d\mathbf{x} dt = \int_{\Theta} (\partial_i u'_i) p^* d\mathbf{x} dt$$

Integrating (H.64e) by part with respect to space

$$\int_{\Theta} (\nabla \cdot \mathbf{u}') p^* d\mathbf{x} dt = \int_0^T p^* \mathbf{u}'|_{\partial\Omega} dt - \int_{\Theta} \mathbf{u}'_i \partial_i p^* d\mathbf{x} dt.$$

Because Ω has periodic boundaries, the first integral is zero, and we can convert the second integral back to vector notation to obtain

$$\int_{\Theta} (\nabla \cdot \mathbf{u}') p^* d\mathbf{x} dt = - \int_{\Theta} \mathbf{u}' \cdot \nabla p^* d\mathbf{x} dt.$$

Combining the integrals, we can express the LHS of (3.39) as

$$\left(\mathcal{L} \begin{bmatrix} \mathbf{u}' \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^* \\ p^* \end{bmatrix} \right) = \int_{\Omega} \mathbf{u}^* \cdot \mathbf{u}'|_{t=0}^{t=T} d\mathbf{x} - \int_{\Theta} \mathbf{u}' \cdot \partial_t \mathbf{u}^* d\mathbf{x} dt. \quad (\text{H.67a})$$

$$- \int_{\Theta} (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}^* \cdot \mathbf{u}) \cdot \mathbf{u}' d\mathbf{x} dt. \quad (\text{H.67b})$$

$$- \int_{\Theta} (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}^*)^\top \mathbf{u} \cdot \mathbf{u}' d\mathbf{x} dt \quad (\text{H.67c})$$

$$- \int_{\Theta} p' (\nabla \cdot \mathbf{u}^*) d\mathbf{x} dt \quad (\text{H.67d})$$

$$- \int_{\Theta} \mathbf{u}' \cdot \nabla p^* d\mathbf{x} dt. \quad (\text{H.67e})$$

Using this and the right-hand side (RHS) of (3.39) we obtain the adjoint system,

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}_\alpha^* - (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}_\alpha^* + (\nabla(I - \alpha^2 \Delta)^{-1} \mathbf{u}_\alpha^*)^T) \mathbf{u}_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot \mathbf{u}_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

To derive the terminal condition of the adjoint system we use the RHS of (3.39).

$$\int_{\Omega} \mathbf{u}_\alpha^*(T) \cdot \mathbf{u}'_\alpha(T) d\mathbf{x} = \int_{\Omega} \mathbf{u}_\alpha^*(0) \cdot \mathbf{u}'_\alpha(0) d\mathbf{x}.$$

From the two forms of the Gâteaux differential, (3.36) and (3.40), we obtain

$$\int_{\Omega} \mathbf{u}_{\alpha}^*(T) \cdot \mathbf{u}_{\alpha}'(T) \, dx = 2 \int_{\Omega} |D| \mathbf{u}_{\alpha}^*(T) \cdot |D| \mathbf{u}_{\alpha}'(T) \, d\mathbf{x}.$$

Therefore,

$$\mathbf{u}_{\alpha}^*(T) = 2|D|^2 \mathbf{u}_{\alpha}(T).$$

I Derivation of the Gâteaux Directional Differential of the Objective Functional

In this Appendix we derive an expression for the Gâteaux directional differential of the objective functional, $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha)$ in terms of the $\dot{H}^1$ inner product and L^2 inner product. This differential is used to obtain the gradient of the objective functional.

We begin by substituting the second definition of (3.32) into (3.34) to obtain

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha)\|_{\dot{H}^1}^2 - \|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{\dot{H}^1}^2].$$

Using the $\dot{H}^1$ seminorm as defined in Appedix A, and taking the limit $\epsilon \rightarrow 0$, we can express $\Phi'_{\alpha,T}$ in the form of the $\dot{H}^1$ inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(T), \mathbf{u}'_\alpha(T) \rangle_{\dot{H}^1}.$$

Using the integral form of the $\dot{H}^s$ seminorm from Appendix A, the Gâteaux differential can be expressed in physical space as

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T) \cdot \mathbf{u}'_\alpha(\mathbf{x}, T) d\mathbf{x}.$$

We now make use of the terminal condition in the adjoint system imposed in Appendix

[H](#)

$$\mathbf{u}_\alpha^*(x, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T).$$

Therefore,

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} \mathbf{u}_\alpha^*(\mathbf{x}, T) \cdot \mathbf{u}'_\alpha(\mathbf{x}, T) d\mathbf{x}.$$

Now using the duality-pairing [3.39](#)

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \int_{\mathbb{T}^3} \mathbf{u}_\alpha^*(\mathbf{x}, 0) \cdot \boldsymbol{\eta}'_\alpha(\mathbf{x}) d\mathbf{x},$$

which lets us express the value of $\Phi'_{\alpha, T}$ with the L^2 inner product

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{L^2}.$$

Therefore,

$$\Phi'_{\alpha, T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}_\alpha(T), \mathbf{u}'_\alpha(T) \rangle_{\dot{H}^1} = \langle \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{L^2}.$$

As described in Section [3.3](#), the differential can also be expressed as a Gevrey inner product using the Riesz representation theorem ([3.35](#)). Using the definition of the Gevrey inner product ([2.22](#)), the L^2 inner product can be converted to the Gevrey inner product,

$$\langle \mathbf{u}_\alpha^*(0), \mathbf{u}'_\alpha(0) \rangle_{L^2} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}.$$

Therefore, we have two forms of the Gâteaux differential as a Gevrey inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}.$$

From this we obtain the gradient of the objective functional

$$\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(0).$$

Bibliography

- [1] P. Absil. *Optimization Algorithms on Matrix Manifolds*. 2008.
- [2] Diego Ayala and Bartosz Protas. Extreme Vortex States and the Growth of Enstrophy in 3D Incompressible Flows. *Journal of Fluid Mechanics*, 818:772–806, May 2017. arXiv:1605.05742 [physics.flu-dyn].
- [3] J. T. Beale, T. Kato, and A. Majda. Remarks on the breakdown of smooth solutions for the 3-D Euler equations. *Communications in Mathematical Physics*, 94(1):61–66, March 1984.
- [4] Marc E. Brachet, Daniel I. Meiron, Steven A. Orszag, B. G. Nickel, Rudolf H. Morf, and Uriel Frisch. Small-scale structure of the Taylor–Green vortex. *Journal of Fluid Mechanics*, 130:411–452, May 1983.
- [5] Miguel D. Bustamante and Marc Brachet. Interplay between the Beale-Kato-Majda theorem and the analyticity-strip method to investigate numerically the incompressible Euler singularity problem. *Physical Review E*, 86(6):066302, December 2012. arXiv:1112.1571 [physics].
- [6] Claudio Canuto, M. Youssuff Hussaini, Alfio Quarteroni, and Thomas A. Zang.

- Spectral Methods: Fundamentals in Single Domains.* Scientific Computation. Springer, Berlin, Heidelberg, 2006.
- [7] Yanping Cao, Evelyn M. Lunasin, and Edriss S. Titi. Global well-posedness of the three-dimensional viscous and inviscid simplified Bardina turbulence models. *Communications in Mathematical Sciences*, 4(4):823–848, December 2006.
- [8] Jiajie Chen and Thomas Y. Hou. Finite time blowup of 2D Boussinesq and 3D Euler equations with $C^{1,\alpha}$ velocity and boundary. *Communications in Mathematical Physics*, 383(3):1559–1667, May 2021. arXiv:1910.00173 [math.AP].
- [9] Jiajie Chen and Thomas Y. Hou. Stable nearly self-similar blowup of the 2D Boussinesq and 3D Euler equations with smooth data I: Analysis, May 2023. arXiv:2210.07191 [math.AP].
- [10] Hyuntae Cho, Yesom Park, and Myungjoo Kang. Solving incompressible Navier–Stokes equations on irregular domains and quadtrees by monolithic approach. *Journal of Computational Physics*, 463:111304, August 2022.
- [11] Ionut Danaila and Bartosz Protas. Computation of Ground States of the Gross-Pitaevskii Functional via Riemannian Optimization. *SIAM Journal on Scientific Computing*, 39(6):B1102–B1129, January 2017. arXiv:1703.07693 [math].
- [12] Charles R. Doering. The 3D Navier-Stokes Problem. *Annual Review of Fluid Mechanics*, 41(Volume 41, 2009):109–128, January 2009.
- [13] Charles R. Doering and J. D. Gibbon. *Applied Analysis of the Navier-Stokes*

- Equations*. Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 1995.
- [14] Charles L Fefferman. Existence and Smoothness of the Navier-Stokes equation, 2022.
- [15] John J. F. Fournier and Robert A. Adams. *Sobolev Spaces*. 2 edition, June 2003.
- [16] J. D. Gibbon. The three-dimensional Euler equations: Where do we stand? *Physica D: Nonlinear Phenomena*, 237(14):1894–1904, August 2008.
- [17] Thomas Y. Hou. Potential Singularity of the 3D Euler Equations in the Interior Domain. *Foundations of Computational Mathematics*, 23(6):2203–2249, December 2023.
- [18] Thomas Y. Hou and De Huang. Potential singularity formation of incompressible axisymmetric Euler equations with degenerate viscosity coefficients, May 2022. arXiv:2102.06663 [math.AP].
- [19] Tosio Kato. Nonstationary flows of viscous and ideal fluids in R^3 . *Journal of Functional Analysis*, 9(3):296–305, March 1972.
- [20] Adam Larios, Mark R. Petersen, Edriss S. Titi, and Beth Wingate. A computational investigation of the finite-time blow-up of the 3D incompressible Euler equations based on the Voigt regularization. *Theoretical and Computational Fluid Dynamics*, 32(1):23–34, February 2018.
- [21] Adam Larios and Edriss S. Titi. On the Higher-Order Global Regularity of the Inviscid Voigt-Regularization of Three-Dimensional Hydrodynamic Models, February 2010. arXiv:0910.3354 [math].

- [22] Jean Leray. Sur le mouvement d'un liquide visqueux emplissant l'espace. *Acta Mathematica*, 63(1):193–248, December 1934.
- [23] David G. Luenberger. *Optimization by Vector Space Methods*. 1969.
- [24] Guo Luo and Thomas Y. Hou. Potentially Singular Solutions of the 3D Incompressible Euler Equations. *Proceedings of the National Academy of Sciences*, 111(36):12968–12973, September 2014. arXiv:1310.0497 [physics.flu-dyn].
- [25] H. K. Moffatt. Singularities in Fluid Mechanics, July 2019. arXiv:1907.06400 [physics].
- [26] Giuseppe Di Molfetta, Giorgio Krstulovic, and Marc Brachet. Self-truncation and scaling in Euler-Voigt-alpha and related fluid models. *Physical Review E*, 92(1):013020, July 2015. arXiv:1502.05544 [physics].
- [27] Jorge Nocedal and Stephen J. Wright. *Numerical Optimization*. Second edition edition, 2000.
- [28] Bartosz Protas. Systematic Search For Extreme and Singular Behavior in Some Fundamental Models of Fluid Mechanics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 380(2225):20210035, June 2022. arXiv:2112.14917 [math].
- [29] James C. Robinson, Jose L. Rodrigo, and Witold Sadowski. *The Three-Dimensional Navier-Stokes Equations*. 2016.
- [30] Xinyu Zhao and Bartosz Protas. Systematic Search for Singularities in 3D Euler Flows. *Journal of Nonlinear Science*, 33(6):120, October 2023.