

SYSTEMATIC SEARCH FOR SINGULARITIES

SYSTEMATIC SEARCH FOR SINGULARITIES IN MODELS OF
3D EULER-VOIGT FLOWS

BY
NOAH BENSLER, B.Sc.

A THESIS
SUBMITTED TO THE DEPARTMENT OF MATHEMATICS & STATISTICS
AND THE SCHOOL OF GRADUATE STUDIES
OF MCMASTER UNIVERSITY
IN PARTIAL FULFILMENT OF THE REQUIREMENTS
FOR THE DEGREE OF
MASTERS OF COMPUTATIONAL SCIENCE & ENGINEERING

© Copyright by Noah Bensler, MONTH 2026

All Rights Reserved

Masters of Computational Science & Engineering (2026)
(Department of Mathematics & Statistics)

McMaster University
Hamilton, Ontario, Canada

TITLE: SYSTEMATIC SEARCH FOR SINGULARITIES IN
MODELS OF 3D EULER-VOIGT FLOWS

AUTHOR: Noah Bensler
B.Sc. (Computer Science & Astrophysics),
University of Calgary, Calgary, Canada

SUPERVISOR: Dr. Bartosz Protas

NUMBER OF PAGES: xiv, 51

Lay Abstract

A lay abstract of not more 150 words must be included explaining the key goals and contributions of the thesis in lay terms that is accessible to the general public.

The Navier-Stokes equations model the behaviour of viscous incompressible fluids. The question of whether or not the Navier-Stokes equations, or the inviscid Euler equations, are a valid description of fluid mechanics remains unanswered. A singularity is the occurrence of an infinite derivative of the velocity at a single point in the fluid flow. If the equations form a singularity from smooth initial conditions, then they are not valid descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to not form singularities from smooth initial conditions. However, by decreasing the regularization parameter we approximate Euler flows. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations. ## We present numerical evidence that CONCLUSION ##

Abstract

Abstract here (no more than 300 words). An abstract of not more than 300 words must be included and indicates the major emphasis of the thesis, new discoveries, and its contribution to knowledge. The style of abstract varies from discipline to discipline but the student should follow an appropriate style. This page must be numbered 'iv'.

The Navier-Stokes equations are a system of partial differential equations used to model the behaviour of viscous incompressible fluids. The question of whether or not unique classical solutions of the Navier-Stokes equations, and inviscid Euler equations, exist for all time, given smooth initial conditions, remains unanswered for the three-dimensional problem.

A singularity is the occurrence of an infinite derivative of the velocity at a single point in the fluid flow. The formation of a singularity from smooth initial conditions would invalidate the equations as accurate descriptions of fluid mechanics. The Euler-Voigt equations are an inviscid regularization of the Euler equations which are known to be globally well-posed from smooth initial conditions. However, by decreasing the regularization parameter the solutions of the Euler-Voigt equations converge

to those of the Euler equations. We examine the Euler-Voigt equations with a PDE-constrained optimization problem to identify smooth initial conditions that produce 'extreme' behaviour in numerical simulations. ## We present numerical evidence that CONCLUSION ##

Your Dedication
Optional second line

Acknowledgements

Acknowledgements go here. An expression of thanks for assistance given by the Supervisor of research and by others should be either set forth on a separate page or incorporated into the Preface (if there is one). These and all subsequent preliminary pages listed under (f), (g) and (h) must be numbered in lower case Roman numerals, i.e., ‘iv’, ‘v’, ‘vi’ etc.

Dr. Bartosz Protas, Dr. Xinyu Zhao, Dr. Jörn Davidsen, members of committee,
Parents

I would like to thank Dr. Bartosz Protas for his support and guidance on my research project. I also want to thank Dr. Jörn Davidsen of the Complexity Science Group for supervising my undergraduate research and inspiring me to pursue computational physics.

Contents

Lay Abstract	iii
Abstract	iv
Acknowledgements	vii
Notation, Definitions, and Abbreviations	xii
Declaration of Academic Achievement	xiv
1 Introduction	1
1.1 Singularity Formation in Models of 3D Fluid Flows	1
1.2 3D Euler-Voigt Equations	3
1.3 Finite Time Blow-up Criteria	4
1.4 Initial Conditions	7
1.5 Outline	11
2 Statement of the Optimization Problem	12
2.1 Formulation of the Problem	12
2.2 Gradient of Objective Function	13

2.3	Riemannian Conjugate Gradient Method	16
2.4	Spatial Discretization & Spectral Methods	18
2.5	Spectrum Validation	18
2.6	Kappa Test Validation	19
2.7	Conservation of the α -Energy	25
3	Results	26
3.1	Pre-optimization Results	26
3.2	Optimization Results	26
4	Conclusion	31
A	Derivation of the Gateaux Directional Differential	32
B	Linearization of the Euler-Voigt System	36
C	Derivation of the Adjoint System	38
D	Proof of Gevrey Class size	43
E	$\ \boldsymbol{\eta}\ _{\dot{H}^1}$ of General Taylor Green Vortex	45
F	$\ \nabla \times \boldsymbol{\eta}\ _{L^\infty}$ of General Taylor Green Vortex	48

List of Figures

- 2.1 Plot showing the expected κ behaviour for different spatial and temporal resolution where δ represents a small arbitrary constant. . . . 21

List of Tables

Notation, Definitions, and Abbreviations

Notation

$A \leq B$	A is less than or equal to B
x	Plain-text to denote scalar x
$\boldsymbol{x}$	Bold-text to denote vector x
$\boldsymbol{x}^\top$	Transpose of vector x
I	Identity Operator

Definitions

Challenge	With respect to video games, a challenge is a set of goals presented to the player that they are tasks with completing; challenges can test a variety of player skills, including accuracy, logical reasoning, and creative problem solving
------------------	---

Abbreviations

PDE	Partial Differential Equation
------------	-------------------------------

3D	three-dimensional
TGV	Taylor Green Vortex

Declaration of Academic Achievement

The student will declare his/her research contribution and, as appropriate, those of colleagues or other contributors to the contents of the thesis.

Xinyu developed most of the math and code

Appendix D-J are mine

1 Introduction

1.1 Singularity Formation in Models of 3D Fluid Flows

The Navier-Stokes equations are a set of partial differential equations (PDEs) that describe the motion of viscous incompressible fluids. The fluids described by the Navier-Stokes equations can be modelled in domains with a wide variety of shape, dimension, and boundary conditions [CITATION]. Because of their flexibility they are utilized to model the behaviour of systems in fields as disparate as aero/hydrodynamics, weather prediction, biophysics, and astrophysics [2][4]. Despite the numerous fields that make use of the 3D Navier-Stokes equations, and extensive research examining their properties, there is no known unique classical solution [9]. In fact, this is the basis of the well known millennium problem from the Clay Institute which offers a million dollar award for the confirmation of the existence, or non-existence, of a solution for smooth initial conditions [10].

The Navier-Stokes system on the domain $\Omega \subseteq \mathbb{R}^3$ is defined as

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p - \nu \Delta \mathbf{u} = 0 \quad \text{in } \Omega \times (0, T], \quad (1.1a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times [0, T], \quad (1.1b)$$

$$\mathbf{u}(x, 0) = \boldsymbol{\eta} \quad (1.1c)$$

where $\mathbf{u} = [u_1, u_2, u_3]^\top$ is the fluid velocity field, p is the scalar pressure, $\nu > 0$ is the coefficient of kinematic viscosity, and T is the time-scale. The value $\boldsymbol{\eta}$ is the initial divergence free velocity field of the fluid at time $t = 0$. Finally, the $\nabla = [\partial_1, \partial_2, \partial_3]^T$ operator is used to define the gradient of the pressure, the tensor $[\nabla \mathbf{u}]_{ij} = \partial_j u_i$ for $i, j = 1, 2, 3$, and the divergence of the velocity field. It is common for scientific research that focuses the fundamental properties of the Navier-Stokes equations to utilize either the unbounded domain $\Omega := \mathbb{R}^3$, or the periodic 3D torus $\Omega := \mathbb{T}_L^3$ with $L > 0$ [9]. For this analysis, we use the unit torus.

A central question in fluid mechanics is whether or not the Navier-Stokes equations are globally well-posed or if singularities can form from smooth initial conditions. A singularity occurs when the derivative of the velocity field becomes infinite at a single point [8]. The formation of a singularity in a fluid flow, from smooth initial conditions, would invalidate the Navier-Stokes equations as an accurate description of fluid flows [9]. The question of singularity formation from smooth initial conditions has also not been answered for the Euler equations [CITATION]. The Euler equations

are the inviscid form of the Navier-Stokes equations, obtained by setting $\nu = 0$

$$\partial_t \mathbf{u} + \mathbf{u} \cdot \nabla \mathbf{u} + \nabla p = 0 \quad \text{in } \Omega \times (0, T], \quad (1.2a)$$

$$\nabla \cdot \mathbf{u} = 0 \quad \text{in } \Omega \times [0, T], \quad (1.2b)$$

$$\mathbf{u}(x, 0) = \boldsymbol{\eta} \quad (1.2c)$$

The Euler are often studied because they are simpler to analyze than the Navier-Stokes equations [Citation]. Additionally, many people believe the Eulers equations are more likely to form singularities than the Navier-Stokes due to the lack of a viscosity term dampening the fluid flow [Citation].

1.2 3D Euler-Voigt Equations

The Euler-Voigt equations are an inviscid regularization of the Euler equations (1.2) which have been shown to be globally well-posed and so do not form singularities from smooth initial conditions [1]. The Euler-Voigt equations are defined as

$$-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p = 0 \quad \text{in } \Omega \times (0, T], \quad (1.3a)$$

$$\nabla \cdot \mathbf{u}_\alpha = 0 \quad \text{in } \Omega \times (0, T], \quad (1.3b)$$

$$\mathbf{u}_\alpha(x, 0) = \boldsymbol{\eta}_\alpha. \quad (1.3c)$$

where we introduce the subscript α to distinguish between solutions of (1.2) and (1.3). The regularization of the Euler-Voigt equations is accomplished with the addition of the term $-\alpha^2 \partial_t \Delta \mathbf{u}_\alpha$ where $\alpha > 0$ is the regularization parameter. This term is used to stabilize simulations without adding artificial viscosity, which would remove energy

from the system [5]. For (1.2) the quantity $\|\mathbf{u}\|_{L^2}^2$ is conserved so long as the solution remains well-posed [CITATION]. For (1.3) the conserved quantity is known as the α -energy [CITATION]. The α -energy is defined as

$$E_\alpha(t) := \|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = E_\alpha(0). \quad (1.4)$$

We note that both (1.2) and (1.3) possess the following time scaling symmetry

$$\tilde{\mathbf{u}}(\mathbf{x}, t) = \lambda \mathbf{u}(\mathbf{x}, \lambda t), \quad (1.5)$$

which will become relevant in our discussion of initial conditions.

1.3 Finite Time Blow-up Criteria

The solutions of (1.3) converge to the solutions of (1.2) as $\alpha \rightarrow 0$. Therefore, the behaviour of the solutions of (1.3) as $\alpha \rightarrow 0$ can be used to determine whether the solutions of the Euler equations will develop a finite time singularity. This was proved by the following theorem from [5].

Theorem 1.3.1. *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$, let $\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\mathbb{T}^3)) \cap C^1([0, T], H^{s-1}(\mathbb{T}^3))$ be the corresponding solutions to (1.2) and (1.3), respectively, where $0 < T < T_{\max}$ and $T_{\max}$ is the maximal time for which a solution to (1.2) exists and is unique. For all $t \in [0, T]$, we have*

$$\begin{aligned} & \|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \leq \\ & (\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2) e^{Ct} + C\alpha^2(e^{Ct} - 1). \end{aligned} \quad (1.6)$$

Based on this theorem we know that if

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (1.7)$$

then for all $t \in [0, T]$,

$$\|\mathbf{u}(t) - \mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla(\mathbf{u}(t) - \mathbf{u}_\alpha(t))\|_{L^2}^2 \rightarrow 0. \quad (1.8)$$

Therefore, if (1.7) is satisfied then $\mathbf{u}_\alpha$ converges to $\mathbf{u}$ as $\alpha \rightarrow 0$ [5]. The convergence of Euler-Voigt to Euler can be used to determine if blow-up occurs in Euler. This was proved by the following theorem from [5].

Theorem 1.3.2. *Let $\boldsymbol{\eta}_\alpha \in H^s$, $s \geq 3$ with $\nabla \cdot \boldsymbol{\eta}_\alpha = 0$ and let $\mathbf{u}_\alpha$ be the corresponding unique solution of (1.3). Suppose that there exists a time $T^* > 0$ such that*

$$\limsup_{\alpha \rightarrow 0^+} \left(\alpha \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2} \right) > 0. \quad (1.9)$$

Then the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}_\alpha$ must develop a singularity within the interval $[0, T^]$.*

This theorem only works in the case where $\boldsymbol{\eta} = \boldsymbol{\eta}_\alpha$, which does satisfy (1.7). However, in order to use this in our optimization search we must generalize to the initial conditions $\boldsymbol{\eta} \neq \boldsymbol{\eta}_\alpha$ that satisfy criteria (1.7). Therefore, we slightly modify Theorem 5.2 from [6] as follows. Special thanks to Dr. Xinyu Zhao for developing the following proof.

Theorem 1.3.3. *Given initial data $\boldsymbol{\eta}, \boldsymbol{\eta}_\alpha \in H^s(\mathbb{T}^3)$ for some $s \geq 3$ satisfying*

$$\|\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta} - \boldsymbol{\eta}_\alpha)\|_{L^2}^2 \rightarrow 0, \quad \text{as } \alpha \rightarrow 0, \quad (1.10)$$

let $\mathbf{u}, \mathbf{u}_\alpha \in C([0, T], H^s(\mathbb{T}^3)) \cap C^1([0, T], H^{s-1}(\mathbb{T}^3))$ be the corresponding solutions to (1.2) and (1.3), respectively. If there exists a finite time $T^ > 0$ such that the solutions $\mathbf{u}_\alpha$ satisfies*

$$\limsup_{\alpha \rightarrow 0} \left(\alpha^2 \sup_{t \in [0, T^*]} \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) > 0, \quad (1.11)$$

then $\mathbf{u}$ develops a singularity in the interval $[0, T^]$.*

Proof. Suppose that the solution $\mathbf{u}$ of the Euler equations (1.2) with the initial condition $\boldsymbol{\eta}$ is regular on $[0, T^*]$. Since the Euler-Voigt equations are globally well-posed, we have

$$\|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2, \quad (1.12)$$

thus

$$\alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla \boldsymbol{\eta}_\alpha\|_{L^2}^2 - \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (1.13)$$

Taking the supremum over $t \in [0, T^*]$ of the above equation, we obtain that

$$\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = \|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta}_\alpha)\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2. \quad (1.14)$$

Taking the limsup as $\alpha \rightarrow 0$, we obtain that

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \limsup_{\alpha \rightarrow 0} \left(\|\boldsymbol{\eta}_\alpha\|_{L^2}^2 + \alpha^2 \|\nabla(\boldsymbol{\eta}_\alpha)\|_{L^2}^2 - \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2}^2 \right). \quad (1.15)$$

We now make use of Theorem (1.3.1), which demonstrates that the convergence of

the solutions of the Euler-Voigt equations (1.3) to the solution of the Euler equations (1.2) is uniform on $[0, T^*]$. Therefore we can reduce the above equation to

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = \|\boldsymbol{\eta}\|_{L^2}^2 - \|\mathbf{u}(t)\|_{L^2}^2. \quad (1.16)$$

Since the energy of regular solutions of the Euler equations is conserved, we have

$$\limsup_{\alpha \rightarrow 0} \left(\sup_{t \in [0, T^*]} \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 \right) = 0, \quad (1.17)$$

which contradicts the hypothesis in the theorem that the above expression is positive.

The second equality holds because

$$\begin{aligned} \limsup_{\alpha \rightarrow 0} \left| \inf_{t \in [0, T^*]} \|\mathbf{u}_\alpha(t)\|_{L^2} - \|\boldsymbol{\eta}\|_{L^2} \right| &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &\leq \limsup_{\alpha \rightarrow 0} \|\mathbf{u}_\alpha(t_\alpha) - \mathbf{u}(t_\alpha)\|_{L^2}^2 + \|\mathbf{u}(t_\alpha) - \boldsymbol{\eta}\|_{L^2}^2 \\ &= 0. \end{aligned}$$

□

1.4 Initial Conditions

In order to formulate our optimization problem we must first describe the properties that each initial condition must have. Firstly, the pseudo-spectral methods used in this study require the solutions of Euler-Voigt to be smooth and real-analytic in order to have exponential convergence [CITATION]. For the Navier-Stokes equations, the viscosity component acts to dissipate small scale structures. Due to this, if the initial

condition is in H^3 then the solution immediately reaches Gevrey regularity [CITATION].

For Euler there is no dissipative term to smooth the initial condition and so for the solution to be analytic, the initial condition must also be [CITATION]. The regularization term of Euler-Voigt ensures global well-posedness but it is not dissipative and so also does not smooth the initial condition [CITATION]. Therefore, to make use of the pseudo-spectral methods, the initial conditions of our optimization search must be analytic. A requirement of the Euler-Voigt equations is that the initial conditions are divergence free. Because of this, we assume without loss of generality that the initial conditions have a mean of zero.

In order to identify initial conditions that satisfy these criteria, we first consider analytic initial conditions that belong to the following Gevrey class

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\mathbb{T}^3) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0. \quad (1.19)$$

We select the following inner product, expressed in both summation and integral form,

$$\forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} := \sum_{\mathbf{j} \in \mathbb{Z}^3} (1 + |2\pi \mathbf{j}|^2)^s e^{4\pi\sigma|\mathbf{j}|} \hat{\mathbf{v}}_{\mathbf{j}} \cdot \overline{\hat{\mathbf{u}}_{\mathbf{j}}} \quad (1.20a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}, \quad (1.20b)$$

where hat denotes the Fourier coefficient, overbar denotes complex conjugation, and the operators $|D|$ and $e^{\sigma|D|}$ are defined via

$$\left[\widehat{|D|\mathbf{v}}\right]_j := 2\pi|j|\hat{\mathbf{v}}_j, \quad \left[\widehat{e^{\sigma|D|}\mathbf{v}}\right]_j := e^{2\pi|j|\sigma}\hat{\mathbf{v}}_j. \quad (1.21)$$

The parameter σ determines the size of the Gevrey space

$$G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s, \quad 0 < \sigma_1 < \sigma_2, \quad (1.22)$$

which is proved in Appendix (D). The value of σ is determined in our discussion of the numerical methods (SECTION REF). As previously mentioned, the initial conditions must be divergence free and have a mean of zero. Therefore, we introduce the following subspace in which optimal initial conditions will be sought

$$V := \{\mathbf{v} \in G^\sigma : \nabla \cdot \mathbf{v} = 0, \quad \int_{\mathbb{T}^3} \mathbf{v} d\mathbf{x} = 0\}. \quad (1.23)$$

Finally, due to the time scaling property of the Euler-Voigt equations (1.5), the characteristic time-scale of our solution will depend on the 'size' of the initial data and the characteristic length scale of the domain ($L=1$). Therefore, we can restrict our discussion to initial conditions which belong to a closed manifold $\mathcal{M}_1 \subset V$ defined as

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = C\}, \quad (1.24)$$

where C is some constant. In order to best compare our results to the available literature we must choose C such that our characteristic time-scale is the same. We can compute the the value of C by analysing our first initial condition, the 3D Taylor

Green Vortex (TGV). The TGV in the domain 3D torus of length scale L is defined as

$$\eta_{TGV}(x) = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0. \end{bmatrix} \quad (1.25)$$

The characteristic time-scale of (1.25) is defined as

$$t_c = \frac{L}{V_0}. \quad (1.26)$$

The $\|\cdot\|_{\dot{H}^1}$ of $\boldsymbol{\eta}_{TGV}$ is

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}L\pi V_0, \quad (1.27)$$

which is derived in Appendix E. The $\|\cdot\|_{L^\infty}$ of the vorticity $\boldsymbol{\omega}_{TGV} = \nabla \times \boldsymbol{\eta}_{TGV}$ is

$$\|\boldsymbol{\omega}_{TGV}\|_{L^\infty} = 4\pi V_0/L, \quad (1.28)$$

which is derived in Appendix F.

Bustamante et al. found numerical evidence that blow-up of the (1.2) occurs at $T^* \approx 4.4$ from (1.25)[CITATION]. Their analysis used parameters $L = 2\pi$ and $V_0 = 1$, and so the characteristic time-scale of their model is $t_c = 2\pi$. Our calculations are performed on the domain with length scale $L = 1$. Thus to match the characteristic time-scale we use $V_0 = 1/2\pi$. These parameters give our model an initial seminorm $\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = \sqrt{3}/2$ and initial vorticity $\|\nabla \times \boldsymbol{\eta}_{TGV}\|_{L^\infty} = 2$.

Therefore, we can restrict our discussion of the TGV to initial conditions with $\dot{H}^1$

seminorm which belong to a closed manifold $\mathcal{M}_1 \subset V$ defined as

$$\mathcal{M}_1 = \{\mathbf{v} \in V : \|\mathbf{v}\|_{\dot{H}^1} = \sqrt{3}/2\}. \quad (1.29)$$

In order to select initial conditions within this manifold, we will first select initial conditions $\boldsymbol{\eta} \in V$ and then apply a retraction operator.

1.5 Outline

outline. In section 2 we provide the description of the optimization problem, the Riemannian conjugate gradient method used to perform iterations, and the numerical methods used to perform the calculations. Additionally, we describe methods used to determine the various parameters of the models including resolution, time step, sigma, etc. Finally, we validate our model using a spectrum resolution, Kappa test, and conservation of $\alpha - energy$.

In section 3 we present the pre-optimization results, Results of optimization at T=3 and results of optimization at T=5.

In section 4 we summarize our results.

2 Statement of the Optimization Problem

2.1 Formulation of the Problem

With our initial conditions and blowup criteria defined, we now define the objection function $\Phi_{\alpha,T} : V \rightarrow \mathbb{R}^+$ as

$$\Phi_{\alpha,T}(\boldsymbol{\eta}) := \|\nabla \mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{L^2}^2 = \|\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)\|_{H^1}^2, \quad (2.30)$$

where $\mathbf{u}_\alpha(T; \boldsymbol{\eta}_\alpha)$ is the solution of the Euler-Voigt equations obtained at time T with the initial condition $\boldsymbol{\eta}_\alpha \in V$ and regularization parameter α . We can now formulate our optimization problem. Given $\alpha, T \in \mathbb{R}_+$, find

$$\tilde{\boldsymbol{\eta}}_{\alpha,T} = \operatorname{argmax}_{\boldsymbol{\eta}_\alpha \in M_1} \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha). \quad (2.31)$$

The objective function is maximized using an iterative relation described in Section (2.3). The solution to the problem is the value of $\boldsymbol{\eta}_{\alpha,T}^n$ as the iteration number $n \rightarrow \infty$. Our goal is to first fix T and solve the optimization problem for a sequence of decrease values of α that converge to 0.

If the initial condition causes a singularity to form in the time interval $[0, T]$, then

$\Phi_{\alpha,T}(\tilde{\boldsymbol{\eta}}_{\alpha,T})$ should diverge as $\alpha \rightarrow 0$.

2.2 Gradient of Objective Function

The optimization problem is solved using an iterative relation which uses an optimal search direction and distance to create the next iteration's initial condition. We compute the optimal search direction using the gradient of the objection functional $\nabla\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)$. This value is computed using an adjoint method which depends on solving a properly defined adjoint system backwards in time. We first introduce the Gâteaux (directional) differential $\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) : V \times V \rightarrow \mathbb{R}$. The first order finite difference equation for the Gateaux differential is,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha + \epsilon \boldsymbol{\eta}'_\alpha) - \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha)]. \quad (2.32)$$

This can also be represented as an inner product on the Gevrey space using the Riesz representation theorem [7]

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = \langle \nabla\Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (2.33)$$

Substituting the second definition of the objective functional (2.30) into the finite difference form of the Gâteaux differential, we derive the Gâteaux differential in the form of the $\dot{H}^1$ inner product and the L^2 inner product,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \langle \mathbf{u}_\alpha(\cdot, T), \mathbf{u}'_\alpha(\cdot, T) \rangle_{\dot{H}^1} = 2 \langle \mathbf{u}_\alpha^*(\cdot, 0), \mathbf{u}'_\alpha(\cdot, 0) \rangle_{L^2} \quad (2.34)$$

The complete derivations for both forms are shown in Appendix (A). In (2.34) the value $\mathbf{u}'_\alpha$ is the solution of the linearization of the Euler-Voigt system (1.3) around its solution $\mathbf{u}_\alpha$. The linearization of the Euler-Voigt system is defined as

$$\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix} := \begin{bmatrix} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (2.35a)$$

$$\mathbf{u}'_\alpha(x, 0) = \boldsymbol{\eta}'_\alpha \quad (2.35b)$$

where $\boldsymbol{\eta}'_\alpha$ and p' are the perturbations of the initial condition and pressure, respectively. The derivation of the linearized form of the Euler-Voigt equations is shown in Appendix (B). The value $\mathbf{u}_\alpha^*$ in (2.34) is the solution of the adjoint system

$$\mathcal{L}^* \begin{bmatrix} \mathbf{u}_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t \mathbf{u}_\alpha^* - (\nabla(I - \alpha^2)^{-1} \mathbf{u}_\alpha^* + (\nabla(I - \alpha^2)^{-1} \mathbf{u}_\alpha^*)^\top) \mathbf{u}_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot \mathbf{u}_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix} \quad (2.36a)$$

$$\mathbf{u}_\alpha^*(\mathbf{x}, T) = 2|D|^2 \mathbf{u}_\alpha(\mathbf{x}, T) \quad (2.36b)$$

which is derived in Appendix (C) by performing integration by parts, in both space and time, on the following duality-pairing relation

$$\begin{aligned}
\left(\mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} \right) &:= \int_0^T \int_{\mathbb{T}^3} \mathcal{L} \begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} d\mathbf{x} dt \\
&= \left(\begin{bmatrix} \mathbf{u}'_\alpha \\ p' \end{bmatrix}, \mathcal{L}^* \begin{bmatrix} \mathbf{u}^*_\alpha \\ p^* \end{bmatrix} \right) + \int_{\mathbb{T}^3} \mathbf{u}'_\alpha(\mathbf{x}, T) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, T) d\mathbf{x} \quad (2.37) \\
&\quad - \int_{\mathbb{T}^3} \boldsymbol{\eta}'(\mathbf{x}) \cdot \mathbf{u}^*_\alpha(\mathbf{x}, 0) d\mathbf{x} \\
&= 0.
\end{aligned}$$

Using the adjoint system we can now define the gradient. We take the Gâteaux differential in the form of the L^2 inner product (2.34) and in the Riesz representation form (2.33)

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}_\alpha; \boldsymbol{\eta}'_\alpha) = 2 \langle \mathbf{u}^*_\alpha(\cdot, 0), \mathbf{u}'_\alpha(\cdot, 0) \rangle_{L^2} = \langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha), \boldsymbol{\eta}'_\alpha \rangle_{G^\sigma}. \quad (2.38)$$

Expressing the L^2 inner product in its integral form,

$$\langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle_{G^\sigma} = \int_{\mathbb{T}^3} \mathbf{u}^*_\alpha(x, 0) \cdot \boldsymbol{\eta}'(x) dx \quad (2.39a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} [(1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}^*_\alpha(x, 0)] \cdot \boldsymbol{\eta}'(x) dx \quad (2.39b)$$

$$= \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}^*_\alpha(x, 0), \boldsymbol{\eta}'(x) \rangle_{G^\sigma} \quad (2.39c)$$

This gives us the Gâteaux differential in the form of the Gevrey inner product

$$\langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle_{G^\sigma} = \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(x, 0), \boldsymbol{\eta}'(x) \rangle_{G^\sigma}. \quad (2.40)$$

Therefore, the gradient of the objective functional is

$$\nabla \Phi_{\alpha,T}(\boldsymbol{\eta}_\alpha) = (1 + |D|^2)^{-3} e^{-2\sigma|D|} \mathbf{u}_\alpha^*(x, 0) \quad (2.41)$$

2.3 Riemannian Conjugate Gradient Method

The optimization solution looks for an initial condition $\boldsymbol{\eta}_\alpha \in M_1$ that maximizes our objective functional. Therefore, not only must our first initial condition $\boldsymbol{\eta}_{\alpha,T}^0$ have a unit $\dot{H}^1$ seminorm, but so to must each following iteration $\boldsymbol{\eta}_{\alpha,T}^{n+1}$. We ensure this constraint is upheld using a Riemannian conjugate gradient method modified from the method used in [9] and [3]. This method has three steps:

1. Project the gradient $\nabla \Phi_T(\boldsymbol{\eta}_\alpha)$ onto the tangent space to $\mathcal{M}_1$ at $\boldsymbol{\eta}_\alpha$
2. Perform a suitable vector transport operation in order to construct a Riemannian conjugate ascent direction using the previous search direction
3. Retract the resulting state back to the constraint manifold $\mathcal{M}_1$

We first define the tangent space to $\mathcal{M}_1$ as

$$\mathcal{T}_v \mathcal{M}_1 = \{ \mathbf{z} \in V : \langle \mathbf{v}, \mathbf{z} \rangle_{\dot{H}^1} = 0 \}.$$

. For any $\mathbf{z} \in \mathcal{T}_{\mathbf{v}}\mathcal{M}_1$, this condition can be expressed as

$$\langle \mathbf{v}, \mathbf{z} \rangle_{\dot{H}^1} = \int_{\mathbb{T}^3} |D| \mathbf{v} \cdot |D| \mathbf{z} d\mathbf{x} \quad (2.42a)$$

$$= \int_{\mathbb{T}^3} (1 + |D|^2)^3 e^{2\sigma|D|} \left((1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v} \right) \cdot \mathbf{z} d\mathbf{x} \quad (2.42b)$$

$$= \langle (1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}, \mathbf{z} \rangle_{G^\sigma} = 0. \quad (2.42c)$$

Thus, at each point $\mathbf{v} \in M_1$, the tangent space $\mathcal{T}_{\mathbf{v}}\mathcal{M}_1$ can be characterized using the unit 'normal' vector given by

$$\mathbf{n}_{\mathbf{v}} = \frac{(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}}{\|(1 + |D|^2)^{-3} e^{-2\sigma|D|} |D|^2 \mathbf{v}\|_{G^\sigma}} \quad (2.43)$$

which allows us to construct the projection operator $\mathbf{P}_{\mathcal{T}_{\mathbf{v}}\mathcal{M}_1} : V \rightarrow \mathcal{T}_{\mathbf{v}}\mathcal{M}_1$,

$$\forall \mathbf{w} \in V, \quad \mathbf{P}_{\mathcal{T}_{\mathbf{v}}\mathcal{M}_1} \mathbf{w} = \mathbf{w} - \langle \mathbf{n}_{\mathbf{v}}, \mathbf{w} \rangle_{G^\sigma} \mathbf{n}_{\mathbf{v}}. \quad (2.44)$$

As stated previously, the initial conditions at each iteration must have a unit $\dot{H}^1$. In order to ensure this we introduce the Retraction operator

$$\mathbf{R}(\mathbf{v}) = \frac{\mathbf{v}}{\|\mathbf{v}\|_{\dot{H}^1}}, \quad \mathbf{v} \neq \mathbf{0}. \quad (2.45)$$

The Reimannian method uses the search direction of the n -th iteration to compute the search direction of the $(n+1)$ -th iteration. This is done by a linear combination of the current direction and a 'momentum' term based on the previous iteration's search direction. However, the two search directions belong to different tangent spaces to $\mathcal{M}_1$ and thus cannot be directly added. In order to allow algebraic operations to be

performed on the vectors of the two tangent spaces we introduce the vector transport operation

$$\mathcal{T}\mathcal{M}_1 \oplus \mathcal{T}\mathcal{M}_1 \rightarrow \mathcal{T}\mathcal{M}_1 : (\varphi, \xi) \mapsto \Gamma_\varphi(\xi) \in \mathcal{T}\mathcal{M}_1,$$

where $\mathcal{T}\mathcal{M}_1 = \bigcup_{v \in \mathcal{M}_1} \mathcal{T}_v \mathcal{M}_1$ is the tangent bundle on $\mathcal{M}_1$. The operation transports the vector field ξ along the manifold $\mathcal{M}_1$ in the direction of φ [CITATION]. This allows us to map search directions between the tangent spaces of two consecutive iterations.

2.4 Spatial Discretization & Spectral Methods

define the domain and mathematical tools used

2.4.1 Time Stepping

what time stepping methods are there. which time stepping method will we use. Stability vs time step size for each alpha.

2.5 Spectrum Validation

USING SPECTRUM VALIDATE THE FUNCTION REMAINS SMOOTH AND REAL ANALYTIC.

The Energy vs wavenumber graph of the initial condition shown below demonstrates the exponential decay expected by a analytic function. As the solution evolves in time the rate of decay decreases due to aliasing. While this can be reduced by anti-aliasing techniques, it cannot be entirely prevented

Low resolution and small alpha simulations are especially vulnerable to aliasing errors and the solutions stop being analytic before the chosen timescale.

The chosen timescales, alphas, and resolutions are validated in the graphs below which show that the spectral decay remains exponential and thus the solutions remain analytic.

2.6 Kappa Test Validation

The simulations are discretized in space using a psuedospectral method and time using a fourth order Runge Kutta method. The kappa test is a means of validating these discretization methods based on the validation method of [?]. We define the value κ as

$$\kappa(\epsilon) = \frac{\epsilon^{-1} [\Phi_{\alpha,T}(\boldsymbol{\eta} + \epsilon\boldsymbol{\eta}') - \Phi_{\alpha,T}(\boldsymbol{\eta})]}{\langle \nabla \Phi_{\alpha,T}(\boldsymbol{\eta}), \boldsymbol{\eta}' \rangle} \quad (2.46)$$

where $\Phi_{\alpha,T}$ is the objective functional (2.30), $\boldsymbol{\eta}$ is the initial condition, and $\nabla \Phi_{\alpha,T}$ is the gradient of the objective functional (2.41). The numerator is the first order finite difference approximation of the Gateaux differential. The denominator is the Reisz form of the differential that utilizes the adjoint system (2.36). If the numerical approximation methods are accurate, then we expect $\kappa \approx 1$.

A log-log plot of $|\kappa - 1|$ vs ϵ can be broken up into three regions, each dominated by different error sources. These are the discretization errors, round off errors, and

truncation errors. The round off errors are caused by finite precision numerical representation and so are approximately of order $O(\epsilon^{-1})$. Therefore, we expect the value of κ to be dominated by round off errors for small ϵ values. The truncation errors are caused by using a first order finite difference formula to calculate the derivative. The truncation errors are of order $O(\epsilon)$. Therefore, we expect the value of κ to be dominated by the truncation errors for large values of ϵ . Between these two regions the value of κ will be determined by the discretization errors in the numerical solutions, which are themselves a product of the spatial and temporal resolution. With smaller values of Δx and Δt causing the error to decrease. This will cause the value of $\log_{10} |\kappa - 1|$ to decrease as the resolution increases. Additionally, an increase in the resolution will decrease the range of ϵ over which the discretization errors dominate. Therefore, we expect the plot of κ vs ϵ to be of the general form shown in the graph below.

The following κ test graphs were generated using a fixed spacial resolution of 128 and with varying temporal resolutions $\Delta t \in [2^{-4}, 2^{-12}]$. The initial conditions used are the 3D Taylor Green, 3D ABC, and 3D ABC Mixture, each with a random initial perturbation. The simulations were run for timescales of $T = 1$ and $T = 3$. The results are shown below.

The results from the 3D Taylor-Green κ test show the expected behaviour from the theoretical kappa test graph. The value of kappa increases at very small and large







Figure 2.1: Plot showing the expected κ behaviour for different spatial and temporal resolution where δ represents a small arbitrary constant.



values of epsilon. The value of kappa approaches 1 as Δt decreases, and so the plateau region lowers and decreases in width. The $T=1$ time scale results follow very closely to the expected results while the $T=3$ time scale simulations show deviations from the expected behaviour. These differences are due to the random initial direction of each simulation having more time to deviate from one another.



The kappa test results for the 3D ABC initial condition exhibit the same lowering of kappa as the time step decreases. However, unlike the Taylor Green initial condition, the results from the 3D ABC κ test have little to no plateau region. This indicates that the 3D ABC initial condition simulations are much more sensitive to the round off and truncation errors.



The results from the 3D ABC Mixture κ test show the same expected behaviour as the 3D Taylor Green kappa test.

The three kappa test results align well with the expected theoretical result. Therefore, we conclude that the discretization and time stepping techniques used in this analysis are valid and accurate.

2.7 Conservation of the α -Energy

Plots showing the conservation of α -Energy at different values of α

As discussed in the introduction, the conserved quantity of the Euler-Voigt (1.3) is known as the α - *Energy* and is defined as

$$E_\alpha(t) := \|\mathbf{u}_\alpha(t)\|_{L^2}^2 + \alpha^2 \|\nabla \mathbf{u}_\alpha(t)\|_{L^2}^2 = E_\alpha(0).$$

This quantity can be used to verify the model by computing the maximum relative error

$$\epsilon_\alpha := \max_{t \in [0, T]} \left| \frac{E_\alpha(t) - E_\alpha(0)}{E_\alpha(0)} \right|, \quad (2.47)$$

for each value of α .

3 Results

3.1 Pre-optimization Results

ALPHAS, RESOLUTIONS, DT, T, SIGMA, PRE-OPTIMIZATION RESULTS INDICATE BLOWUP WILL NOT OCCUR

3.2 Optimization Results

TEXT

3.2.1 $\alpha = 16/1024$

TEXT

Resolution - 128

TEXT

Resolution - 256

TEXT

Resolution - 512

TEXT

Resolution - 1024

TEXT

3.2.2 $\alpha = 8/1024$

TEXT

Resolution - 128

TEXT

Resolution - 256

TEXT

Resolution - 512

TEXT

Resolution - 1024

TEXT

3.2.3 $\alpha = 4/1024$

TEXT

Resolution - 128

TEXT

Resolution - 256

TEXT

Resolution - 512

TEXT

Resolution - 1024

TEXT

3.2.4 $\alpha = 2/1024$

TEXT

Resolution - 128

TEXT

Resolution - 256

TEXT

Resolution - 512

TEXT

Resolution - 1024

TEXT

3.2.5 $\alpha = 1/1024$

TEXT

Resolution - 128

TEXT

Resolution - 256

TEXT

Resolution - 512

TEXT

Resolution - 1024

TEXT

3.2.6 $\alpha \rightarrow 0$

TEXT

$$\alpha^2 \Phi_{\alpha,T}(\tilde{\eta}_{\alpha,T})$$

TEXT

$$\boldsymbol{\eta}_\alpha^{vs} \tilde{\boldsymbol{\eta}}_{\alpha,T}$$

TEXT

4 Conclusion

Every thesis also needs a concluding chapter

A Derivation of the Gateaux Directional Differential

The equation for the Gateaux differential (2.32) is

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') := \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\Phi_{\alpha,T}(\boldsymbol{\eta} + \epsilon \boldsymbol{\eta}') - \Phi_{\alpha,T}(\boldsymbol{\eta})].$$

The objective functional (2.30) is defined as,

$$\Phi_{\alpha,T}(\boldsymbol{\eta}) := \|\nabla \mathbf{u}_{\alpha}(T; \boldsymbol{\eta})\|_{L^2}^2 = \|\mathbf{u}_{\alpha}(T; \boldsymbol{\eta})\|_{\dot{H}^1}^2.$$

Substituting the second definition of (2.30) into (2.32),

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [\|\mathbf{u}_{\alpha}(T; \boldsymbol{\eta} + \epsilon \boldsymbol{\eta}')\|_{\dot{H}^1}^2 - \|\mathbf{u}_{\alpha}(T; \boldsymbol{\eta})\|_{\dot{H}^1}^2].$$

The equation for the $\dot{H}^1$ seminorm is

$$\|u\|_{\dot{H}^1}^2 = (2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 |\hat{u}_{\mathbf{k}}|^2.$$

Substituting in the equation for the $\dot{H}^1$ seminorm

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [(2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 |\hat{\mathbf{u}}_k + \epsilon \hat{\mathbf{u}}'_k|^2 - (2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 |\hat{\mathbf{u}}_k|^2].$$

This simplifies to

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} [(2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 2\epsilon |\hat{\mathbf{u}}_k \hat{\mathbf{u}}'_k| + (2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 \epsilon^2 |\hat{\mathbf{u}}'_k|^2].$$

Taking the limit $\epsilon \rightarrow 0$

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2(2\pi)^3 \sum_{k \in \mathbb{Z}^3} |k|^2 |\hat{\mathbf{u}}_k \hat{\mathbf{u}}'_k|.$$

This can be expressed in the integral form

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \int_{\mathbb{T}^3} \sum_{k \in \mathbb{Z}^3} |k|^2 \hat{\mathbf{u}}_k \hat{\mathbf{u}}'_k e^{2ikx} dx.$$

Which is equivalent to the inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \left\langle \sum_{k \in \mathbb{Z}^3} |k| \hat{\mathbf{u}}_k e^{ikx}, \sum_{k \in \mathbb{Z}^3} |k| \hat{\mathbf{u}}'_k e^{ikx} \right\rangle_{L^2}.$$

This gives us the Gateaux differential in the form of the $\dot{H}^1$ inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \langle \mathbf{u}_\alpha(\cdot, T), \mathbf{u}'_\alpha(\cdot, T) \rangle_{\dot{H}^1}.$$

The integral form of the $\dot{H}^1$ inner product is

$$\langle \mathbf{u}, \mathbf{v} \rangle_{\dot{H}^1} = \int_{\mathbb{T}^3} |D| \mathbf{u} \cdot |D| \mathbf{v} \, dx,$$

where $|D|$ is the operator defined in (1.21). Therefore, the Gateaux differential can be expressed as

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \int_{\mathbb{T}^3} |D| \mathbf{u}_\alpha(x, T) \cdot |D| \mathbf{u}'_\alpha(x, T) \, dx,$$

which can be rearranged to the form

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \int_{\mathbb{T}^3} 2|D|^2 \mathbf{u}_\alpha(x, T) \cdot \mathbf{u}'_\alpha(x, T) \, dx.$$

We now make use of the adjoint system terminal condition derived in Appendix (C)

$$\mathbf{u}_\alpha^*(x, T) = 2|D|^2 \mathbf{u}_\alpha(x, T).$$

Therefore

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \int_{\mathbb{T}^3} \mathbf{u}_\alpha^*(x, T) \cdot \mathbf{u}'_\alpha(x, T) \, dx.$$

Now using the duality-pairing defined in Appendix (C)

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \int_{\mathbb{T}^3} \boldsymbol{\eta}'(x) \cdot \mathbf{u}_\alpha^*(x, 0) \, dx.$$

This is the inner product

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = \langle \mathbf{u}_\alpha^*(x, 0), \boldsymbol{\eta}'(x) \rangle_{L^2}.$$

Therefore,

$$\Phi'_{\alpha,T}(\boldsymbol{\eta}; \boldsymbol{\eta}') = 2 \langle \mathbf{u}_{\alpha}(\cdot, T), \mathbf{u}'_{\alpha}(\cdot, T) \rangle_{\dot{H}^1} = \langle \mathbf{u}_{\alpha}^*(x, 0), \boldsymbol{\eta}'(x) \rangle_{L^2}.$$

B Linearization of the Euler-Voigt System

Euler-Voigt Equations,

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha + \partial_t \mathbf{u}_\alpha + (\mathbf{u}_\alpha \cdot \nabla) \mathbf{u}_\alpha + \nabla p = 0 \\ \nabla \cdot \mathbf{u}_\alpha = 0 \\ \mathbf{u}_\alpha|_{t=0} = \boldsymbol{\eta}. \end{cases}$$

Substituting in $\mathbf{u}_\alpha = \mathbf{u}_\alpha + \mathbf{u}'_\alpha$, $\boldsymbol{\eta} = \boldsymbol{\eta} + \boldsymbol{\eta}'$, and $p = p + p'$,

$$\begin{cases} -\alpha^2 \partial_t \Delta (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \partial_t (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + ((\mathbf{u}_\alpha + \mathbf{u}'_\alpha) \cdot \nabla) (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) + \nabla (p + p') = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta} + \boldsymbol{\eta}'. \end{cases}$$

Expanding

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}_\alpha - \alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}_\alpha + \\ \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p + \nabla p' = 0 \\ \nabla \cdot (\mathbf{u}_\alpha + \mathbf{u}'_\alpha) = 0 \\ (\mathbf{u}_\alpha + \mathbf{u}'_\alpha)|_{t=0} = \boldsymbol{\eta} + \boldsymbol{\eta}'. \end{cases}$$

Substituting in the Euler-Voigt equations we simplify to

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}'_\alpha + \nabla p' = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'. \end{cases}$$

Linearizing the system by removing non-linear terms, we derive the Linearized Euler-Voigt system

$$\begin{cases} -\alpha^2 \partial_t \Delta \mathbf{u}'_\alpha + \partial_t \mathbf{u}'_\alpha + \mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p' = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'. \end{cases}$$

This can be rearranged to the form

$$\begin{cases} \partial_t \mathbf{u}'_\alpha + (I - \alpha^2 \Delta)^{-1} (\mathbf{u}_\alpha \cdot \nabla \mathbf{u}'_\alpha + \mathbf{u}'_\alpha \cdot \nabla \mathbf{u}_\alpha + \nabla p') = 0 \\ \nabla \cdot \mathbf{u}'_\alpha = 0 \\ \mathbf{u}'_\alpha|_{t=0} = \boldsymbol{\eta}'. \end{cases} \tag{B.48}$$

C Derivation of the Adjoint System

For this derivation we will exclude the α subscript on the u_α , u'_α , and u^*_α terms for the sake of simplicity. Additionally, we substitute the integration with the following notation

$$\int_0^T \int_{\mathbb{T}^3} u \, dt = \int_D u \, dt$$

We derive the adjoint system using the following duality pairing relation,

$$(L[u'], u^*) := \int_D L[u'] u^* \, dx \, dt = (u', L[u^*]) + \int_a^b u' u^*|_{t=T} - \int_a^b u' u^*|_{t=0} \, dx = 0.$$

Substituting in the equation from the Euler-Voigt perturbation system,

$$(L[u'], u^*) := \int_D (\partial_t u' + (I - \alpha^2 \Delta)^{-1} (u \cdot \nabla u' + u' \cdot \nabla u + \nabla p')) u^* + (\nabla \cdot u') p^* \, dx \, dt.$$

This integral can be split into five parts. Solving the first part with integration by parts, with $u = u^*$ and $v' = \partial_t u'$

$$\int_D u^* \partial_t u' \, dx \, dt = \int_{\mathbb{T}^3} u^* u'|_{t=0}^{t=T} \, dx - \int_D u' \partial_t u^* \, dx \, dt.$$

Starting with the second part and expressing it in Einstein Summation notation,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} u_j \frac{\partial u'_i}{\partial x_j} \right] u_i^* dx dt.$$

Using the derivative product rule and the fact that $\nabla \cdot u = 0$,

$$\frac{\partial}{\partial x_j} (u_j u'_i) = \frac{\partial u_j}{\partial x_j} u'_i + u_j \frac{\partial u'_i}{\partial x_j} = u_j \frac{\partial u'_i}{\partial x_j}.$$

Substituting this into the integral,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} \frac{\partial}{\partial x_j} (u_j u'_i) \right] u_i^* dx dt.$$

Using the periodic boundaries of the system, we can now apply integration by parts with $u = [(I - \alpha^2 \Delta)^{-1} u_i^*]$ and $v' = \frac{\partial}{\partial x_j} (u_j u'_i)$,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = -u_j u'_i \frac{\partial}{\partial x_j} [(I - \alpha^2 \Delta)^{-1} u_i^*] dx dt.$$

Converting back to vector notation,

$$[(I - \alpha^2 \Delta)^{-1} u \cdot \nabla u'] \cdot u^* dx dt = -u' \cdot [u \cdot \nabla [(I - \alpha^2 \Delta)^{-1} u^*]] dx dt.$$

Starting with the third part and expressing it in Einstein Summation notation,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = \left[(I - \alpha^2 \Delta)^{-1} u'_j \frac{\partial u_i}{\partial x_j} \right] u_i^* dx dt.$$

Applying integration by parts with $u = [(I - \alpha^2 \Delta)^{-1} u_i^*]$ and $v' = \frac{\partial}{\partial x_j} (u'_j u_i)$

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u'_j u_i \frac{\partial}{\partial x_j} [(I - \alpha^2 \Delta)^{-1} u_i^*] dx dt.$$

Because the goal is to extract u' and $v \cdot w = v^T \cdot w^T$, we can apply a transpose to the above equation and so swap the indices to,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u'_i u_j \frac{\partial}{\partial x_i} [(I - \alpha^2 \Delta)^{-1} u_j^*] dx dt.$$

Converting back to vector notation,

$$[(I - \alpha^2 \Delta)^{-1} u' \cdot \nabla u] \cdot u^* dx dt = -u' \cdot \left[u \cdot [\nabla (I - \alpha^2 \Delta)^{-1} u^*]^T \right] dx dt.$$

Starting with the fourth part of the integral and expressing it in Einstein Summation notation,

$$\nabla p' \cdot u^* dx dt = \frac{\partial p'}{\partial x_i} u_i^* dx dt.$$

Applying integration by parts with $u = u_i^*$ and $v' = \partial x_i p'$,

$$\nabla p' \cdot u^* dx dt = -p' \frac{\partial u_i^*}{\partial x_i} dx dt.$$

Converting back to vector notation,

$$\nabla p' \cdot u^* dx dt = -p' (\nabla \cdot u^*) dx dt.$$

Starting with the final part of the integral and expressing it in Einstein Summation notation,

$$(\nabla \cdot u') p^* dx dt = \frac{\partial u'_i}{\partial x_i} p^* dx dt.$$

Applying integration by parts with $u = p^*$ and $v' = \partial x_i u'_i$,

$$(\nabla \cdot u') p^* dx dt = -\frac{\partial p^*}{\partial x_i} u'_i dx dt.$$

Converting back to vector notation,

$$(\nabla \cdot u') p^* dx dt = -u' \cdot \nabla p^* dx dt.$$

Combining the integrals,

$$\int_{\mathbb{T}^3} u^* u' \Big|_{t=0}^{t=T} dx - u' \left[\partial_t u^* - [u \cdot \nabla [(I - \alpha^2 \Delta)^{-1} u^*]] - \left[u \cdot [\nabla (I - \alpha^2 \Delta)^{-1} u^*]^T \right] - \nabla p^* \right] - p' (\nabla \cdot u^*)$$

This gives us our adjoint system,

$$\mathbb{L}^* \begin{bmatrix} u_\alpha^* \\ p_\alpha^* \end{bmatrix} := \begin{bmatrix} -\partial_t u^* - (\nabla (I - \alpha^2)^{-1} u_\alpha^* + (\nabla (I - \alpha^2)^{-1} u_\alpha^*)^T) u_\alpha - \nabla p_\alpha^* \\ -\nabla \cdot u_\alpha^* \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}$$

To derive the terminal condition of the adjoint system we compare the second term of the duality pairing to the objective functional.

$$\begin{aligned} \int_{\mathbb{T}^3} u'(x, T) \mathbf{u}^*(x, T) dx &= 2 \langle u(x, T), u'(x, T) \rangle_{\dot{H}^1} \\ &= 2 \int_{\mathbb{T}^3} |D| u(x, T) \mathbf{u}'(x, T) dx \end{aligned}$$

$$= \int_{\mathbb{T}^3} 2|D|^2 u(x, T) \mathbf{u}'(x, T) dx$$

Therefore,

$$\mathbf{u}^*(x, T) = 2|D|^2 u(x, T)$$

D Proof of Gevrey Class size

Proof. The definition of the Gevrey class (1.19) is

$$G^\sigma := \left\{ \mathbf{v} \in H^s(\mathbb{T}^3) : \|\mathbf{v}\|_{G^\sigma} = \langle \mathbf{v}, \mathbf{v} \rangle_{G^\sigma}^{1/2} < \infty \right\}, \quad s \geq 3, \quad \sigma > 0.$$

The definition of the Gevrey inner product (1.20) is

$$\forall \mathbf{v}, \mathbf{u} \in G^\sigma, \quad \langle \mathbf{v}, \mathbf{u} \rangle_{G^\sigma} := \int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma|D|} \mathbf{v} \cdot \mathbf{u} \, d\mathbf{x}.$$

Let σ_1, σ_2 be defined such that

$$0 < \sigma_1 < \sigma_2.$$

Therefore,

$$\begin{aligned} \|\mathbf{v}\|_{G^{\sigma_2}} &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_2|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} \\ &> \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^{2\sigma_1|D|} \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^{\sigma_1}} \\ &> \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s e^0 \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{G^0} \\ &= \left[\int_{\mathbb{T}^3} (1 + |D|^2)^s \mathbf{v} \cdot \mathbf{v} \, d\mathbf{x} \right]^{1/2} = \|\mathbf{v}\|_{H^s} \end{aligned} \tag{D.49}$$

Thus,

$$\|\mathbf{v}\|_{G^{\sigma_2}} > \|\mathbf{v}\|_{G^{\sigma_1}} > \|\mathbf{v}\|_{G^0}.$$

It follows from this that if $\|\mathbf{v}\|_{G^{\sigma_2}} < \infty$ then $\|\mathbf{v}\|_{G^{\sigma_1}} < \infty$ and so if $v \in G^{\sigma_2}$ then $v \in G^{\sigma_1}$. However it does not follow from this that if $\|\mathbf{v}\|_{G^{\sigma_1}} < \infty$ then $\|\mathbf{v}\|_{G^{\sigma_2}} < \infty$, and so it is not required that if $v \in G^{\sigma_1}$ then $v \in G^{\sigma_2}$. The same argument shows that if $v \in G^{\sigma_1}$ then $v \in G^0$ which is equivalent to $v \in H^s$. Therefore,

$$G^{\sigma_2} \subset G^{\sigma_1} \subset G^0 = H^s$$

□

E $\|\boldsymbol{\eta}\|_{\dot{H}^1}$ of General Taylor Green Vortex

The general form of the 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is

$$\boldsymbol{\eta}_{TGV} = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0. \end{bmatrix}$$

To simplify the notation we drop the subscript TGV and substitute in

$$x' = 2\pi x/L, \quad y' = 2\pi y/L, \quad z' = 2\pi z/L.$$

The formula for $\|\boldsymbol{\eta}\|_{\dot{H}^1}$ is

$$\|\boldsymbol{\eta}\|_{\dot{H}^1} = \|\nabla \boldsymbol{\eta}\|_{L^2} = \left[\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} \right]^{1/2},$$

where ∂_i is the partial derivative with respect to the i -th component and $\boldsymbol{\eta}_j$ is the j -th component of $\boldsymbol{\eta}$. Because the 3rd component of $\boldsymbol{\eta}$ is zero, the partial derivative

is also zero. This leaves us with six integrals to evaluate.

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= \int_{\Omega} (\partial_1 \boldsymbol{\eta}_1)^2 d\mathbf{x} + \int_{\Omega} (\partial_2 \boldsymbol{\eta}_1)^2 d\mathbf{x} + \int_{\Omega} (\partial_3 \boldsymbol{\eta}_1)^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (\partial_1 \boldsymbol{\eta}_2)^2 d\mathbf{x} + \int_{\Omega} (\partial_2 \boldsymbol{\eta}_2)^2 d\mathbf{x} + \int_{\Omega} (\partial_3 \boldsymbol{\eta}_2)^2 d\mathbf{x}. \end{aligned}$$

Evaluating each partial derivative

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= \int_{\Omega} (2\pi V_0/L \cos(x') \cos(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \sin(x') \sin(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \sin(x') \cos(y') \sin(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (2\pi V_0/L \sin(x') \sin(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (-2\pi V_0/L \cos(x') \cos(y') \cos(z'))^2 d\mathbf{x} \\ &\quad + \int_{\Omega} (2\pi V_0/L \cos(x') \sin(y') \sin(z'))^2 d\mathbf{x}. \end{aligned}$$

Which simplifies to

$$\begin{aligned} \sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} &= 4\pi^2 V_0^2 / L^2 \int_{\Omega} \cos^2(x') \cos^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2 / L^2 \int_{\Omega} \sin^2(x') \sin^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2 / L^2 \int_{\Omega} \sin^2(x') \cos^2(y') \sin^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2 / L^2 \int_{\Omega} \sin^2(x') \sin^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2 / L^2 \int_{\Omega} \cos^2(x') \cos^2(y') \cos^2(z') d\mathbf{x} \\ &\quad + 4\pi^2 V_0^2 / L^2 \int_{\Omega} \cos^2(x') \sin^2(y') \sin^2(z') d\mathbf{x}. \end{aligned}$$

Because the integrand is separable, the integrals can be reduced to products of,

$$\int_0^L \cos^2(x') dx = \frac{L}{2}, \quad \text{and} \quad \int_0^L \sin^2(x') dx = \frac{L}{2}.$$

Therefore, each of the six integrals is equal to

$$\frac{4\pi^2 V_0^2}{L^2} \frac{L^3}{8}.$$

The sum of the six integrals is

$$\sum_{i,j=1}^3 \int_{\Omega} (\partial_i \boldsymbol{\eta}_j)^2 d\mathbf{x} = 6 \frac{4\pi^2 V_0^2}{L^2} \frac{L^3}{8} = 3\pi^2 V_0^2 L.$$

Substituting into the formula for $\|\cdot\|_{\dot{H}^1}$

$$\|\boldsymbol{\eta}_{TGV}\|_{\dot{H}^1} = [3\pi^2 V_0^2 L]^{1/2} = \sqrt{3L}\pi V_0.$$

F $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ of General Taylor Green Vortex

The general form of the 3D Taylor Green Vortex on the 3D torus Ω with characteristic length L is

$$\boldsymbol{\eta}_{TGV} = \begin{bmatrix} V_0 \sin(2\pi x/L) \cos(2\pi y/L) \cos(2\pi z/L) \\ -V_0 \cos(2\pi x/L) \sin(2\pi y/L) \cos(2\pi z/L) \\ 0. \end{bmatrix}$$

To simplify the notation we drop the subscript TGV and substitute in

$$x' = 2\pi x/L, \quad y' = 2\pi y/L, \quad z' = 2\pi z/L.$$

The vorticity of $\boldsymbol{\eta}$ is

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} \partial_2 \boldsymbol{\eta}_3 - \partial_3 \boldsymbol{\eta}_2 \\ \partial_3 \boldsymbol{\eta}_1 - \partial_1 \boldsymbol{\eta}_3 \\ \partial_1 \boldsymbol{\eta}_2 - \partial_2 \boldsymbol{\eta}_1. \end{bmatrix}$$

where ∂_i is the partial derivative with respect to the i -th component and $\boldsymbol{\eta}_j$ is the j -th component of $\boldsymbol{\eta}$. Substituting in the partial derivatives

$$\nabla \times \boldsymbol{\eta} = \begin{bmatrix} 0 - (-(2\pi V_0/L) \cos(x'/L) \sin(y'/L) \sin(z'/L)) \\ -(2\pi V_0/L) \sin(x'/L) \cos(y'/L) \sin(z'/L) - 0 \\ (2\pi V_0/L) \sin(x'/L) \sin(y'/L) \cos(z'/L) + (2\pi V_0/L) \sin(x'/L) \sin(y'/L) \cos(z'/L). \end{bmatrix}$$

which simplifies to

$$\nabla \times \boldsymbol{\eta}_{TG} = \begin{bmatrix} (2\pi V_0/L) \cos(x'/L) \sin(y'/L) \sin(z'/L) \\ -(2\pi V_0/L) \sin(x'/L) \cos(y'/L) \sin(z'/L) \\ (4\pi V_0/L) \sin(x'/L) \sin(y'/L) \cos(z'/L) \end{bmatrix}$$

The $\|\nabla \times \boldsymbol{\eta}\|_{L^\infty}$ norm is the maximum absolute value of the components of $\nabla \times \boldsymbol{\eta}$.

$$\begin{aligned} \|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = \max(& |(2\pi V_0/L) \cos(x'/L) \sin(y'/L) \sin(z'/L)| \\ & , |-(2\pi V_0/L) \sin(x'/L) \cos(y'/L) \sin(z'/L)| \\ & , |(4\pi V_0/L) \sin(x'/L) \sin(y'/L) \cos(z'/L)|) \end{aligned}$$

The maximums are

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = \max((2\pi V_0/L), (2\pi V_0/L), (4\pi V_0/L))$$

Therefore,

$$\|\nabla \times \boldsymbol{\eta}_{TG}\|_{L^\infty} = 4\pi V_0/L.$$

Bibliography

- [1] Yanping Cao, Evelyn M. Lunasin, and Edriss S. Titi. Global well-posedness of the three-dimensional viscous and inviscid simplified Bardina turbulence models. *Communications in Mathematical Sciences*, 4(4):823–848, December 2006. Publisher: International Press of Boston.
- [2] Hyuntae Cho, Yesom Park, and Myungjoo Kang. Solving incompressible Navier–Stokes equations on irregular domains and quadtrees by monolithic approach. *Journal of Computational Physics*, 463:111304, August 2022.
- [3] Ionut Danaila and Bartosz Protas. Computation of Ground States of the Gross-Pitaevskii Functional via Riemannian Optimization. *SIAM Journal on Scientific Computing*, 39(6):B1102–B1129, January 2017. arXiv:1703.07693 [math].
- [4] Charles R. Doering. The 3D Navier-Stokes Problem. *Annual Review of Fluid Mechanics*, 41(Volume 41, 2009):109–128, January 2009. Publisher: Annual Reviews.
- [5] Adam Larios, Mark R. Petersen, Edriss S. Titi, and Beth Wingate. A computational investigation of the finite-time blow-up of the 3D incompressible Euler equations based on the Voigt regularization. *Theoretical and Computational Fluid Dynamics*, 32(1):23–34, February 2018.

-
- [6] Adam Larios and Edriss S. Titi. On the Higher-Order Global Regularity of the Inviscid Voigt-Regularization of Three-Dimensional Hydrodynamic Models, February 2010. arXiv:0910.3354 [math].
- [7] David G. Luenberger. *Optimization by Vector Space Methods*.
- [8] H. K. Moffatt. Singularities in Fluid Mechanics, July 2019. arXiv:1907.06400 [physics].
- [9] Bartosz Protas. Systematic Search For Extreme and Singular Behavior in Some Fundamental Models of Fluid Mechanics. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 380(2225):20210035, June 2022. arXiv:2112.14917 [math].
- [10] James C. Robinson. The Navier–Stokes regularity problem. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 378(2174):20190526, June 2020. Publisher: Royal Society.